

Combinatorial Speculation and Combinatorial Conjecture for Mathematics

Linfan Mao

(Chinese Academy of Mathematics and System Sciences, Beijing 100080, P.R.China)

Email: maolinfan@163.com

Abstract Extended: This survey was widely spread after reported at a combinatorial conference of China in 2006. As a powerful tool for dealing with relations among objectives, combinatorics mushroomed in the past century, particularly in catering to the need of computer science and children games. However, an even more important work for mathematician is to apply it to other mathematics and other sciences besides just to find combinatorial behavior for objectives. *How can it contributes more to the entirely mathematical science, not just in various games, but in metric mathematics? What is a right mathematical theory for the original face of our world?* I presented a well-known proverb, i.e., the six blind men and an elephant in the *3th Northwest Conference on Number Theory and Smarandache's Notion of China* and answered the second question to be Smarandache multi-spaces in logic. Prior to that explaining, I have brought a heartening conjecture for advancing mathematics in 2005, i.e., *mathematical science can be reconstructed from or made by combinatorialization* after a long time speculation, also a bringing about Smarandache multi-space for mathematics. This conjecture is not just like an open problem, but more like a deeply thought for advancing the modern mathematics. The main trend of modern sciences is overlap and hybrid. Whence the mathematics of 21st century should be consistency with the science development in the 21st century, i.e., the *mathematical combinatorics* resulting in the combinatorial conjecture for mathematics. For introducing more readers known this heartening mathematical notion for sciences, there would be no simple stopping point if I began to incorporate the more recent development, for example, the *combinatorially differential geometry*, so it being published here in its original form to survey these thinking and ideas for mathematics and cosmological physics, such as those of multi-spaces, map geometries and combinatorial structures of cosmoses. Some open problems are also included for the advance of 21st mathematics by a combinatorial speculation. More recent progresses can be found in papers and books nearly published, for example, in [20]-[23] for details.

Key words: combinatorial speculation, combinatorial conjecture for mathematics, Smarandache multi-space, M-theory, combinatorial cosmos.

AMS(2000): 03C05,05C15,51D20,51H20,51P05,83C05,83E50.

¹Received May 25, 2007. Accepted June 15, 2007

²Reported at the *2nd Conference on Combinatorics and Graph Theory of China*, Aug. 16-19, 2006, Tianjing, P.R.China

§1. The role of classical combinatorics in mathematics

Modern science has so advanced that to find a universal genus in the society of sciences is nearly impossible. Thereby a scientist can only give his or her contribution in one or several fields. The same thing also happens for researchers in combinatorics. Generally, combinatorics deals with twofold:

Question 1.1. *to determine or find structures or properties of configurations, such as those structure results appeared in graph theory, combinatorial maps and design theory, ..., etc..*

Question 1.2. *to enumerate configurations, such as those appeared in the enumeration of graphs, labeled graphs, rooted maps, unrooted maps and combinatorial designs, ..., etc..*

Consider the contribution of a question to science. We can separate mathematical questions into three ranks:

Rank 1 *they contribute to all sciences.*

Rank 2 *they contribute to all or several branches of mathematics.*

Rank 3 *they contribute only to one branch of mathematics, for instance, just to the graph theory or combinatorial theory.*

Classical combinatorics is just a *rank 3 mathematics* by this view. This conclusion is despair for researchers in combinatorics, also for me 5 years ago. *Whether can combinatorics be applied to other mathematics or other sciences? Whether can it contributes to human's lives, not just in games?*

Although become a universal genus in science is nearly impossible, *our world is a combinatorial world*. A combinatorician should stand on all mathematics and all sciences, not just on classical combinatorics and with a real combinatorial notion, i.e., *combining different fields into a unifying field* ([29]-[32]), such as combine different or even anti-branches in mathematics or science into a unifying science for its freedom of research ([28]). This notion requires us answering three questions for solving a combinatorial problem before. *What is this problem working for? What is its objective? What is its contribution to science or human's society?* After these works be well done, modern combinatorics can applied to all sciences and all sciences are combinatorialization.

§2. The metrical combinatorics and mathematics combinatorialization

There is a prerequisite for the application of combinatorics to other mathematics and other sciences, i.e, to introduce various metrics into combinatorics, ignored by the classical combinatorics since they are the fundamental of scientific realization for our world. This speculation was firstly appeared in the beginning of Chapter 5 of my book [16]:

... our world is full of measures. For applying combinatorics to other branch of mathematics, a good idea is pullback measures on combinatorial objects again, ignored by the classical combinatorics and reconstructed or make combinatorial generalization for the classical

mathematics, such as those of algebra, differential geometry, Riemann geometry, Smarandache geometries, \dots and the mechanics, theoretical physics, \dots .

The combinatorial conjecture for mathematics, abbreviated to *CCM* is stated in the following.

Conjecture 2.1(CCM Conjecture) *Mathematical science can be reconstructed from or made by combinatorialization.*

Remark 2.1 We need some further clarifications for this conjecture.

(1) This conjecture assumes that one can select finite combinatorial rulers and axioms to reconstruct or make generalization for classical mathematics.

(2) Classical mathematics is a particular case in the combinatorialization of mathematics, i.e., the later is a combinatorial generalization of the former.

(3) We can make one combinatorialization of different branches in mathematics and find new theorems after then.

Therefore, a branch in mathematics can not be ended if it has not been combinatorialization and all mathematics can not be ended if its combinatorialization has not completed. There is an assumption in one's realization of our world, i.e., *science can be made by mathematicalization*, which enables us get a similar combinatorial conjecture for the science.

Conjecture 2.2(CCS Conjecture) *Science can be reconstructed from or made by combinatorialization.*

A typical example for the combinatorialization of classical mathematics is the *combinatorial map theory*, i.e., a combinatorial theory for surfaces([14]-[15]). Combinatorially, a surface is topological equivalent to a polygon with even number of edges by identifying each pairs of edges along a given direction on it. If label each pair of edges by a letter $e, e \in \mathcal{E}$, a surface S is also identifying to a cyclic permutation such that each edge $e, e \in \mathcal{E}$ just appears two times in S , one is e and another is e^{-1} . Let a, b, c, \dots denote the letters in \mathcal{E} and A, B, C, \dots the sections of successive letters in a linear order on a surface S (or a string of letters on S). Then, a surface can be represented as follows:

$$S = (\dots, A, a, B, a^{-1}, C, \dots),$$

where, $a \in \mathcal{E}, A, B, C$ denote a string of letters. Define three elementary transformations as follows:

$$\begin{aligned} (O_1) \quad & (A, a, a^{-1}, B) \Leftrightarrow (A, B); \\ (O_2) \quad & (i) \quad (A, a, b, B, b^{-1}, a^{-1}) \Leftrightarrow (A, c, B, c^{-1}); \\ & (ii) \quad (A, a, b, B, a, b) \Leftrightarrow (A, c, B, c); \\ (O_3) \quad & (i) \quad (A, a, B, C, a^{-1}, D) \Leftrightarrow (B, a, A, D, a^{-1}, C); \\ & (ii) \quad (A, a, B, C, a, D) \Leftrightarrow (B, a, A, C^{-1}, a, D^{-1}). \end{aligned}$$

If a surface S can be obtained from S_0 by these elementary transformations O_1 - O_3 , we say that S is elementary equivalent with S_0 , denoted by $S \sim_{El} S_0$. Then we can get the classification theorem of compact surface as follows([29]):

Any compact surface is homeomorphic to one of the following standard surfaces:

(P_0) the sphere: aa^{-1} ;

(P_n) the connected sum of $n, n \geq 1$ tori:

$$a_1b_1a_1^{-1}b_1^{-1}a_2b_2a_2^{-1}b_2^{-1} \cdots a_nb_na_n^{-1}b_n^{-1};$$

(Q_n) the connected sum of $n, n \geq 1$ projective planes:

$$a_1a_1a_2a_2 \cdots a_na_n.$$

A map M is a connected topological graph cellularly embedded in a surface S . In 1973, Tutte suggested an algebraic representation for an embedding graph on a locally orientable surface ([16]):

A *combinatorial map* $M = (\mathcal{X}_{\alpha,\beta}, \mathcal{P})$ is defined to be a basic permutation \mathcal{P} , i.e, for any $x \in \mathcal{X}_{\alpha,\beta}$, no integer k exists such that $\mathcal{P}^k x = \alpha x$, acting on $\mathcal{X}_{\alpha,\beta}$, the disjoint union of *quadrlicells* Kx of $x \in X$ (the base set), where $K = \{1, \alpha, \beta, \alpha\beta\}$ is the *Klein group* satisfying the following two conditions:

(i) $\alpha\mathcal{P} = \mathcal{P}^{-1}\alpha$;

(ii) the group $\Psi_J = \langle \alpha, \beta, \mathcal{P} \rangle$ is transitive on $\mathcal{X}_{\alpha,\beta}$.

For a given map $M = (\mathcal{X}_{\alpha,\beta}, \mathcal{P})$, it can be shown that $M^* = (\mathcal{X}_{\beta,\alpha}, \mathcal{P}\alpha\beta)$ is also a map, call it the *dual* of the map M . The vertices of M are defined as the pairs of conjugate orbits of \mathcal{P} action on $\mathcal{X}_{\alpha,\beta}$ by the condition (i) and edges the orbits of K on $\mathcal{X}_{\alpha,\beta}$, for example, for $\forall x \in \mathcal{X}_{\alpha,\beta}$, $\{x, \alpha x, \beta x, \alpha\beta x\}$ is an edge of the map M . Define the faces of M to be the vertices in the dual map M^* . Then the Euler characteristic $\chi(M)$ of the map M is

$$\chi(M) = \nu(M) - \varepsilon(M) + \phi(M)$$

where, $\nu(M), \varepsilon(M), \phi(M)$ are the number of vertices, edges and faces of the map M , respectively. For each vertex of a map M , its valency is defined to be the length of the orbits of \mathcal{P} action on a quadrlicell incident with u .

For example, the graph K_4 on the tours with one face length 4 and another 8 shown in Fig.2.1




Fig.2.1

can be algebraically represented by $(\mathcal{X}_{\alpha,\beta}, \mathcal{P})$ with $\mathcal{X}_{\alpha,\beta} = \{x, y, z, u, v, w, \alpha x, \alpha y, \alpha z, \alpha u, \alpha v, \alpha w, \beta x, \beta y, \beta z, \beta u, \beta v, \beta w, \alpha\beta x, \alpha\beta y, \alpha\beta z, \alpha\beta u, \alpha\beta v, \alpha\beta w\}$ and

$$\begin{aligned} \mathcal{P} &= (x, y, z)(\alpha\beta x, u, w)(\alpha\beta z, \alpha\beta u, v)(\alpha\beta y, \alpha\beta v, \alpha\beta w) \\ &\times (\alpha x, \alpha z, \alpha y)(\beta x, \alpha w, \alpha u)(\beta z, \alpha v, \beta u)(\beta y, \beta w, \beta v) \end{aligned}$$

with 4 vertices, 6 edges and 2 faces on an orientable surface of genus 1.

By the view of combinatorial maps, these standard surfaces P_0, P_n, Q_n for $n \geq 1$ is nothing but the bouquet B_n on a locally orientable surface with just one face. Therefore, combinatorial maps are the combinatorialization of surfaces.

Many open problems are motivated by the *CCM Conjecture*. For example, a *Gauss mapping* among surfaces is defined as follows.

Let $\mathcal{S} \subset R^3$ be a surface with an orientation \mathbf{N} . The mapping $\mathbf{N} : \mathcal{S} \rightarrow R^3$ takes its value in the unit sphere

$$S^2 = \{(x, y, z) \in R^3 \mid x^2 + y^2 + z^2 = 1\}$$

along the orientation \mathbf{N} . The map $\mathbf{N} : \mathcal{S} \rightarrow S^2$, thus defined, is called the *Gauss mapping*.

We know that for a point $P \in \mathcal{S}$ such that the Gaussian curvature $K(P) \neq 0$ and V a connected neighborhood of P with K does not change sign,

$$K(P) = \lim_{A \rightarrow 0} \frac{N(A)}{A},$$

where A is the area of a region $B \subset V$ and $N(A)$ is the area of the image of B by the Gauss mapping $N : \mathcal{S} \rightarrow S^2([2],[4])$. Now the questions are

(i) what is its combinatorial meaning of the Gauss mapping? How to realizes it by combinatorial maps?

(ii) how can we define various curvatures for maps and rebuilt these results in the classical differential geometry?

Let \mathcal{S} be a compact orientable surface. Then the *Gauss-Bonnet theorem* asserts that

$$\int \int_{\mathcal{S}} K d\sigma = 2\pi\chi(\mathcal{S}),$$

where K is the Gaussian curvature of \mathcal{S} .

By the *CCM Conjecture*, the following questions should be considered.

(i) *How can we define various metrics for combinatorial maps, such as those of length, distance, angle, area, curvature,...*?

(ii) *Can we rebuilt the Gauss-Bonnet theorem by maps for dimensional 2 or higher dimensional compact manifolds without boundary?*

One can see references [15] and [16] for more open problems for the classical mathematics motivated by this *CCM Conjecture*, also raise new open problems for his or her research works.

§3. The contribution of combinatorial speculation to mathematics

3.1. The combinatorialization of algebra

By the view of combinatorics, algebra can be seen as a combinatorial mathematics itself. The combinatorial speculation can generalize it by the means of combinatorialization. For this objective, a Smarandache multi-algebraic system is combinatorially defined in the following definition.

Definition 3.1([17],[18]) *For any integers $n, n \geq 1$ and $i, 1 \leq i \leq n$, let A_i be a set with an operation set $O(A_i)$ such that $(A_i, O(A_i))$ is a complete algebraic system. Then the union*

$$\bigcup_{i=1}^n (A_i, O(A_i))$$

is called an n multi-algebra system.

An example of multi-algebra systems is constructed by a finite additive group. Now let n be an integer, $Z_1 = (\{0, 1, 2, \dots, n-1\}, +)$ an additive group (*mod* n) and $P = (0, 1, 2, \dots, n-1)$ a permutation. For any integer $i, 0 \leq i \leq n-1$, define

$$Z_{i+1} = P^i(Z_1)$$

satisfying that if $k + l = m$ in Z_1 , then $P^i(k) +_i P^i(l) = P^i(m)$ in Z_{i+1} , where $+_i$ denotes the binary operation $+_i : (P^i(k), P^i(l)) \rightarrow P^i(m)$. Then we know that

$$\bigcup_{i=1}^n Z_i$$

is an n multi-algebra system .

The conception of multi-algebra systems can be extensively used for generalizing conceptions and results for these existent algebraic structures, such as those of groups, rings, bodies, fields and vector spaces, ..., etc.. Some of them are explained in the following.

Definition 3.2 Let $\tilde{G} = \bigcup_{i=1}^n G_i$ be a closed multi-algebra system with a binary operation set $O(\tilde{G}) = \{\times_i, 1 \leq i \leq n\}$. If for any integer $i, 1 \leq i \leq n$, $(G_i; \times_i)$ is a group and for $\forall x, y, z \in \tilde{G}$ and any two binary operations \times and \circ , $\times \neq \circ$, there is one operation, for example the operation \times satisfying the distribution law to the operation \circ provided their operation results exist, i.e.,

$$x \times (y \circ z) = (x \times y) \circ (x \times z),$$

$$(y \circ z) \times x = (y \times x) \circ (z \times x),$$

then \tilde{G} is called a multi-group.

For a multi-group $(\tilde{G}, O(\tilde{G}))$, $\tilde{G}_1 \subset \tilde{G}$ and $O(\tilde{G}_1) \subset O(\tilde{G})$, call $(\tilde{G}_1, O(\tilde{G}_1))$ a sub-multi-group of $(\tilde{G}, O(\tilde{G}))$ if \tilde{G}_1 is also a multi-group under the operations in $O(\tilde{G}_1)$, denoted by $\tilde{G}_1 \preceq \tilde{G}$. For two sets A and B , if $A \cap B = \emptyset$, we denote the union $A \cup B$ by $A \oplus B$. Then we get a generalization of the *Lagrange theorem* of finite group.

Theorem 3.1([18]) For any sub-multi-group \tilde{H} of a finite multi-group \tilde{G} , there is a representation set T , $T \subset \tilde{G}$, such that

$$\tilde{G} = \bigoplus_{x \in T} x\tilde{H}.$$

For a sub-multi-group \tilde{H} of \tilde{G} , $\times \in O(\tilde{H})$ and $\forall g \in \tilde{G}(\times)$, if for $\forall h \in \tilde{H}$,

$$g \times h \times g^{-1} \in \tilde{H},$$

then call \tilde{H} a normal sub-multi-group of \tilde{G} . An order of operations in $O(\tilde{G})$ is said an oriented operation sequence, denoted by $\vec{O}(\tilde{G})$. We get a generalization of the *Jordan-Hölder theorem* for finite multi-groups.

Theorem 3.2([18]) For a finite multi-group $\tilde{G} = \bigcup_{i=1}^n G_i$ and an oriented operation sequence $\vec{O}(\tilde{G})$, the length of maximal series of normal sub-multi-groups is a constant, only dependent on \tilde{G} itself.

In Definition 2.2, choose $n = 2$, $G_1 = G_2 = \tilde{G}$. Then \tilde{G} is a body. If $(G_1; \times_1)$ and $(G_2; \times_2)$ both are commutative groups, then \tilde{G} is a field. For multi-algebra systems with two or more operations on one set, we introduce the conception of multi-rings and multi-vector spaces in the following.

Definition 3.3 Let $\tilde{R} = \bigcup_{i=1}^m R_i$ be a closed multi-algebra system with double binary operation set $O(\tilde{R}) = \{(+_i, \times_i), 1 \leq i \leq m\}$. If for any integers i, j , $i \neq j$, $1 \leq i, j \leq m$, $(R_i; +_i, \times_i)$ is a ring and for $\forall x, y, z \in \tilde{R}$,

$$(x +_i y) +_j z = x +_i (y +_j z), \quad (x \times_i y) \times_j z = x \times_i (y \times_j z)$$

and

$$x \times_i (y +_j z) = x \times_i y +_j x \times_i z, \quad (y +_j z) \times_i x = y \times_i x +_j z \times_i x$$

provided all their operation results exist, then \tilde{R} is called a multi-ring. If for any integer $1 \leq i \leq m$, $(R; +_i, \times_i)$ is a filed, then \tilde{R} is called a multi-filed.

Definition 3.4 Let $\tilde{V} = \bigcup_{i=1}^k V_i$ be a closed multi-algebra system with binary operation set

$O(\tilde{V}) = \{(\dot{+}_i, \cdot_i) \mid 1 \leq i \leq m\}$ and $\tilde{F} = \bigcup_{i=1}^k F_i$ a multi-filed with double binary operation set

$O(\tilde{F}) = \{(+_i, \times_i) \mid 1 \leq i \leq k\}$. If for any integers i, j , $1 \leq i, j \leq k$ and $\forall \mathbf{a}, \mathbf{b}, \mathbf{c} \in \tilde{V}$, $k_1, k_2 \in \tilde{F}$,

(i) $(V_i; \dot{+}_i, \cdot_i)$ is a vector space on F_i with vector additive $\dot{+}_i$ and scalar multiplication \cdot_i ;

(ii) $(\mathbf{a} \dot{+}_i \mathbf{b}) \dot{+}_j \mathbf{c} = \mathbf{a} \dot{+}_i (\mathbf{b} \dot{+}_j \mathbf{c})$;

(iii) $(k_1 +_i k_2) \cdot_j \mathbf{a} = k_1 +_i (k_2 \cdot_j \mathbf{a})$;

provided all those operation results exist, then \tilde{V} is called a multi-vector space on the multi-filed \tilde{F} with a binary operation set $O(\tilde{V})$, denoted by $(\tilde{V}; \tilde{F})$.

Similar to multi-groups, we can also obtain results for multi-rings and multi-vector spaces to generalize classical results in rings or linear spaces. Certainly, results can be also found in the references [17] and [18].

3.2. The combinatorialization of geometries

First, we generalize classical metric spaces by the combinatorial speculation.

Definition 3.5 A multi-metric space is a union $\tilde{M} = \bigcup_{i=1}^m M_i$ such that each M_i is a space with metric ρ_i for $\forall i, 1 \leq i \leq m$.

We generalized two well-known results in metric spaces.

Theorem 3.3([19]) Let $\tilde{M} = \bigcup_{i=1}^m M_i$ be a completed multi-metric space. For an ϵ -disk sequence $\{B(\epsilon_n, x_n)\}$, where $\epsilon_n > 0$ for $n = 1, 2, 3, \dots$, the following conditions hold:

(i) $B(\epsilon_1, x_1) \supset B(\epsilon_2, x_2) \supset B(\epsilon_3, x_3) \supset \dots \supset B(\epsilon_n, x_n) \supset \dots$;

(ii) $\lim_{n \rightarrow +\infty} \epsilon_n = 0$.

Then $\bigcap_{n=1}^{+\infty} B(\epsilon_n, x_n)$ only has one point.

Theorem 3.4([19]) Let $\tilde{M} = \bigcup_{i=1}^m M_i$ be a completed multi-metric space and T a contraction on \tilde{M} . Then

$$1 \leq \# \Phi(T) \leq m.$$

Particularly, let $m = 1$. We get the *Banach fixed-point theorem* again.

Corollary 3.1(Banach) *Let M be a metric space and T a contraction on M . Then T has just one fixed point.*

Smarandache geometries were proposed by Smarandache in [29] which are generalization of classical geometries, i.e., these *Euclid, Lobachevshy-Bolyai-Gauss* and *Riemann geometries* may be united altogether in a same space, by some Smarandache geometries under the combinatorial speculation. These geometries can be either partially Euclidean and partially Non-Euclidean, or Non-Euclidean. In general, Smarandache geometries are defined in the next.

Definition 3.6 *An axiom is said to be Smarandachely denied if the axiom behaves in at least two different ways within the same space, i.e., validated and invalidated, or only invalidated but in multiple distinct ways.*

A Smarandache geometry is a geometry which has at least one Smarandachely denied axiom(1969).

For example, let us consider an euclidean plane \mathbf{R}^2 and three non-collinear points A, B and C . Define s -points as all usual euclidean points on \mathbf{R}^2 and s -lines as any euclidean line that passes through one and only one of points A, B and C . Then this geometry is a Smarandache geometry because two axioms are Smarandachely denied comparing with an Euclid geometry:

(i) The axiom (A5) that through a point exterior to a given line there is only one parallel passing through it is now replaced by two statements: *one parallel* and *no parallel*. Let L be an s -line passing through C and is parallel in the euclidean sense to AB . Notice that through any s -point not lying on AB there is one s -line parallel to L and through any other s -point lying on AB there is no s -lines parallel to L such as those shown in Fig.3.1(a).




Fig.3.1

(ii) The axiom that through any two distinct points there exists one line passing through them is now replaced by; *one s-line* and *no s-line*. Notice that through any two distinct s -points D, E collinear with one of A, B and C , there is one s -line passing through them and through any two distinct s -points F, G lying on AB or non-collinear with one of A, B and C , there is no s -line passing through them such as those shown in Fig.3.1(b).

A *Smarandache n -manifold* is an n -dimensional manifold that supports a Smarandache geometry. Now there are many approaches to construct Smarandache manifolds for $n = 2$. A

general way is by the so called *map geometries* without or with boundary underlying orientable or non-orientable maps proposed in references [14] and [15] firstly.

Definition 3.7 For a combinatorial map M with each vertex valency ≥ 3 , endow with a real number $\mu(u), 0 < \mu(u) < \frac{4\pi}{\rho_M(u)}$, to each vertex $u, u \in V(M)$. Call (M, μ) a map geometry without boundary, $\mu(u)$ an angle factor of the vertex u and orientable or non-orientable if M is orientable or not.

Definition 3.8 For a map geometry (M, μ) without boundary and faces $f_1, f_2, \dots, f_l \in F(M), 1 \leq l \leq \phi(M)-1$, if $S(M) \setminus \{f_1, f_2, \dots, f_l\}$ is connected, then call $(M, \mu)^{-l} = (S(M) \setminus \{f_1, f_2, \dots, f_l\}, \mu)$ a map geometry with boundary f_1, f_2, \dots, f_l , where $S(M)$ denotes the locally orientable surface underlying map M .

The realization for vertices $u, v, w \in V(M)$ in a space \mathbf{R}^3 is shown in Fig.3.2, where $\rho_M(u)\mu(u) < 2\pi$ for the vertex u , $\rho_M(v)\mu(v) = 2\pi$ for the vertex v and $\rho_M(w)\mu(w) > 2\pi$ for the vertex w , are called to be elliptic, euclidean or hyperbolic, respectively.




Fig.3.2

On an Euclid plane \mathbf{R}^2 , a straight line passing through an elliptic or a hyperbolic point is shown in Fig.3.3.




Fig.3.3

Theorem 3.5([17]) *There are Smarandache geometries, including paradoxist geometries, non-geometries and anti-geometries in map geometries without or with boundary.*

Generally, we can ever generalize the ideas in Definitions 3.7 and 3.8 to a metric space and find new geometries.

Definition 3.9 Let U and W be two metric spaces with metric ρ , $W \subseteq U$. For $\forall u \in U$, if there is a continuous mapping $\omega : u \rightarrow \omega(u)$, where $\omega(u) \in \mathbf{R}^n$ for an integer $n, n \geq 1$ such that for any number $\epsilon > 0$, there exists a number $\delta > 0$ and a point $v \in W$, $\rho(u - v) < \delta$ such that $\rho(\omega(u) - \omega(v)) < \epsilon$, then U is called a metric pseudo-space if $U = W$ or a bounded metric pseudo-space if there is a number $N > 0$ such that $\forall w \in W$, $\rho(w) \leq N$, denoted by (U, ω) or (U^-, ω) , respectively.

For the case $n = 1$, we can also explain $\omega(u)$ being an angle function with $0 < \omega(u) \leq 4\pi$ as in the case of map geometries without or with boundary, i.e.,

$$\omega(u) = \begin{cases} \omega(u)(\text{mod}4\pi), & \text{if } u \in W, \\ 2\pi, & \text{if } u \in U \setminus W \end{cases} \quad (*)$$

and get some interesting metric pseudo-space geometries. For example, let $U = W = \text{Euclid plane} = \Sigma$, then we obtained some interesting results for pseudo-plane geometries (Σ, ω) as shown in the following([17]).

Theorem 3.6 In a pseudo-plane (Σ, ω) , if there are no euclidean points, then all points of (Σ, ω) is either elliptic or hyperbolic.

Theorem 3.7 There are no saddle points and stable knots in a pseudo-plane plane (Σ, ω) .

Theorem 3.8 For two constants ρ_0, θ_0 , $\rho_0 > 0$ and $\theta_0 \neq 0$, there is a pseudo-plane (Σ, ω) with

$$\omega(\rho, \theta) = 2\left(\pi - \frac{\rho_0}{\theta_0\rho}\right) \text{ or } \omega(\rho, \theta) = 2\left(\pi + \frac{\rho_0}{\theta_0\rho}\right)$$

such that

$$\rho = \rho_0$$

is a limiting ring in (Σ, ω) .

Now for an m -manifold M^m and $\forall u \in M^m$, choose $U = W = M^m$ in Definition 3.9 for $n = 1$ and $\omega(u)$ a smooth function. We get a pseudo-manifold geometry (M^m, ω) on M^m . By definitions in the reference [2], a *Minkowski norm* on M^m is a function $F : M^m \rightarrow [0, +\infty)$ such that

- (i) F is smooth on $M^m \setminus \{0\}$;
- (ii) F is 1-homogeneous, i.e., $F(\lambda\bar{u}) = \lambda F(\bar{u})$ for $\bar{u} \in M^m$ and $\lambda > 0$;
- (iii) for $\forall y \in M^m \setminus \{0\}$, the symmetric bilinear form $g_y : M^m \times M^m \rightarrow R$ with

$$g_y(\bar{u}, \bar{v}) = \frac{1}{2} \frac{\partial^2 F^2(y + s\bar{u} + t\bar{v})}{\partial s \partial t} \Big|_{t=s=0}$$

is positive definite and a *Finsler manifold* is a manifold M^m endowed with a function $F : TM^m \rightarrow [0, +\infty)$ such that

- (i) F is smooth on $TM^m \setminus \{0\} = \bigcup \{T_{\bar{x}}M^m \setminus \{0\} : \bar{x} \in M^m\}$;
- (ii) $F|_{T_{\bar{x}}M^m} \rightarrow [0, +\infty)$ is a Minkowski norm for $\forall \bar{x} \in M^m$.

As a special case, we choose $\omega(\bar{x}) = F(\bar{x})$ for $\bar{x} \in M^m$, then (M^m, ω) is a *Finsler manifold*. Particularly, if $\omega(\bar{x}) = g_{\bar{x}}(y, y) = F^2(x, y)$, then (M^m, ω) is a *Riemann manifold*. Therefore, we get a relation for Smarandache geometries with Finsler or Riemann geometry.

Theorem 3.9 *There is an inclusion for Smarandache, pseudo-manifold, Finsler and Riemann geometries as shown in the following:*

$$\begin{aligned} \{\text{Smarandache geometries}\} &\supset \{\text{pseudo-manifold geometries}\} \\ &\supset \{\text{Finsler geometry}\} \\ &\supset \{\text{Riemann geometry}\}. \end{aligned}$$

Other purely mathematical results on the combinatorially differential geometry, particularly the combinatorially Riemannian geometry can be found in recently finished papers [20] – [23] of mine.

§4. The contribution of combinatorial speculation to theoretical physics

The progress of theoretical physics in last twenty years of the 20th century enables human beings to probe the mystic cosmos: *where are we came from? where are we going to?*. Today, these problems still confuse eyes of human beings. Accompanying with research in cosmos, new puzzling problems also arose: *Whether are there finite or infinite cosmoses? Are there just one? What is the dimension of the Universe? We do not even know what the right degree of freedom in the Universe is*, as Witten said([3]).

We are used to the idea that our living space has three dimensions: *length, breadth and height*, with time providing the fourth dimension of spacetime by Einstein. Applying his principle of general relativity, i.e. *all the laws of physics take the same form in any reference system* and equivalence principle, i.e., *there are no difference for physical effects of the inertial force and the gravitation in a field small enough.*, Einstein got the *equation of gravitational field*

$$R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} + \lambda g_{\mu\nu} = -8\pi GT_{\mu\nu}.$$

where $R_{\mu\nu} = R_{\nu\mu} = R_{\mu\nu}^{\alpha}$,

$$R_{\mu\nu}^{\alpha} = \frac{\partial \Gamma_{\mu\nu}^{\alpha}}{\partial x^{\nu}} - \frac{\partial \Gamma_{\mu\nu}^{\alpha}}{\partial x^{\nu}} + \Gamma_{\mu\nu}^{\alpha} \Gamma_{\alpha\nu}^i - \Gamma_{\mu\nu}^{\alpha} \Gamma_{\alpha i}^{\nu},$$

$$\Gamma_{mn}^g = \frac{1}{2}g^{pq} \left(\frac{\partial g_{mp}}{\partial u^n} + \frac{\partial g_{np}}{\partial u^m} - \frac{\partial g_{mn}}{\partial u^p} \right)$$

and $R = g^{\nu\mu} R_{\nu\mu}$.

Combining the Einstein's equation of gravitational field with the *cosmological principle*, i.e., *there are no difference at different points and different orientations at a point of a cosmos*

on the metric 10^4 l.y. , Friedmann got a standard model of cosmos. The metrics of the standard cosmos are

$$ds^2 = -c^2 dt^2 + a^2(t) \left[\frac{dr^2}{1 - Kr^2} + r^2 (d\theta^2 + \sin^2 \theta d\phi^2) \right]$$

and

$$g_{tt} = 1, \quad g_{rr} = -\frac{R^2(t)}{1 - Kr^2}, \quad g_{\phi\phi} = -r^2 R^2(t) \sin^2 \theta.$$

The standard model of cosmos enables the birth of big bang model of the Universe in thirties of the 20th century. The following diagram describes the developing process of the Universe in different periods after the Big Bang.




Fig.4.1

4.1. The M-theory

The M-theory was established by Witten in 1995 for the unity of those five already known string theories and superstring theories, which postulates that all matter and energy can be reduced to *branes* of energy vibrating in an 11 dimensional space, then in a higher dimensional space solve the Einstein's equation of gravitational field under some physical conditions ([1],[3],[26]-[27]). Here, a *brane* is an object or subspace which can have various spatial dimensions. For any integer $p \geq 0$, a p -*brane* has length in p dimensions. For example, a 0-*brane* is just a point or particle; a 1-*brane* is a string and a 2-*brane* is a surface or membrane, \dots .

We mainly discuss line elements in differential forms in Riemann geometry. By a geometrical view, these p -branes in M-theory can be seen as *volume elements in spaces*. Whence, we

can construct a graph model for p -branes in a space and combinatorially research graphs in spaces.

Definition 4.1 For each m -brane \mathbf{B} of a space \mathbf{R}^m , let $(n_1(\mathbf{B}), n_2(\mathbf{B}), \dots, n_p(\mathbf{B}))$ be its unit vibrating normal vector along these p directions and $q: \mathbf{R}^m \rightarrow \mathbf{R}^4$ a continuous mapping. Now construct a graph phase $(\mathcal{G}, \omega, \Lambda)$ by

$$V(\mathcal{G}) = \{p\text{-branes } q(\mathbf{B})\},$$

$$E(\mathcal{G}) = \{(q(\mathbf{B}_1), q(\mathbf{B}_2)) | \text{there is an action between } \mathbf{B}_1 \text{ and } \mathbf{B}_2\},$$

$$\omega(q(\mathbf{B})) = (n_1(\mathbf{B}), n_2(\mathbf{B}), \dots, n_p(\mathbf{B})),$$

and

$$\Lambda(q(\mathbf{B}_1), q(\mathbf{B}_2)) = \text{forces between } \mathbf{B}_1 \text{ and } \mathbf{B}_2.$$

Then we get a graph phase $(\mathcal{G}, \omega, \Lambda)$ in \mathbf{R}^4 . Similarly, if $m = 11$, it is a graph phase for the M-theory.

As an example for applying M-theory to find an accelerating expansion cosmos of 4-dimensional cosmoses from supergravity compactification on hyperbolic spaces is the *Townsend-Wohlfarth type metric* in which the line element is

$$ds^2 = e^{-m\phi(t)}(-S^6 dt^2 + S^2 dx_3^2) + r_c^2 e^{2\phi(t)} ds_{H_m}^2,$$

where

$$\phi(t) = \frac{1}{m-1}(\ln K(t) - 3\lambda_0 t),$$

$$S^2 = K^{\frac{m}{m-1}} e^{-\frac{m+2}{m-1}\lambda_0 t}$$

and

$$K(t) = \frac{\lambda_0 \zeta r_c}{(m-1) \sin[\lambda_0 \zeta |t + t_1|]}$$

with $\zeta = \sqrt{3 + 6/m}$. This solution is obtainable from space-like brane solution and if the proper time ς is defined by $d\varsigma = S^3(t)dt$, then the conditions for expansion and acceleration are $\frac{dS}{d\varsigma} > 0$ and $\frac{d^2 S}{d\varsigma^2} > 0$. For example, the expansion factor is 3.04 if $m = 7$, i.e., a really expanding cosmos.

According to M-theory, the evolution picture of our cosmos started as a perfect 11 dimensional space. However, this 11 dimensional space was unstable. The original 11 dimensional space finally cracked into two pieces, a 4 and a 7 dimensional subspaces. The cosmos made the 7 of the 11 dimensions curled into a tiny ball, allowing the remaining 4 dimensions to inflate at enormous rates, the Universe at the final.

4.2. The combinatorial cosmos

The combinatorial speculation made the following combinatorial cosmos in the reference [17].

Definition 4.2 A combinatorial cosmos is constructed by a triple (Ω, Δ, T) , where

$$\Omega = \bigcup_{i \geq 0} \Omega_i, \quad \Delta = \bigcup_{i \geq 0} O_i$$

and $T = \{t_i; i \geq 0\}$ are respectively called the cosmos, the operation or the time set with the following conditions hold.

(1) (Ω, Δ) is a Smarandache multi-space dependent on T , i.e., the cosmos (Ω_i, O_i) is dependent on time parameter t_i for any integer $i, i \geq 0$.

(2) For any integer $i, i \geq 0$, there is a sub-cosmos sequence

$$(S) : \Omega_i \supset \cdots \supset \Omega_{i1} \supset \Omega_{i0}$$

in the cosmos (Ω_i, O_i) and for two sub-cosmoses (Ω_{ij}, O_i) and (Ω_{il}, O_i) , if $\Omega_{ij} \supset \Omega_{il}$, then there is a homomorphism $\rho_{\Omega_{ij}, \Omega_{il}} : (\Omega_{ij}, O_i) \rightarrow (\Omega_{il}, O_i)$ such that

(i) for $\forall (\Omega_{i1}, O_i), (\Omega_{i2}, O_i), (\Omega_{i3}, O_i) \in (S)$, if $\Omega_{i1} \supset \Omega_{i2} \supset \Omega_{i3}$, then

$$\rho_{\Omega_{i1}, \Omega_{i3}} = \rho_{\Omega_{i1}, \Omega_{i2}} \circ \rho_{\Omega_{i2}, \Omega_{i3}},$$

where \circ denotes the composition operation on homomorphisms.

(ii) for $\forall g, h \in \Omega_i$, if for any integer i , $\rho_{\Omega, \Omega_i}(g) = \rho_{\Omega, \Omega_i}(h)$, then $g = h$.

(iii) for $\forall i$, if there is an $f_i \in \Omega_i$ with

$$\rho_{\Omega_i, \Omega_i \cap \Omega_j}(f_i) = \rho_{\Omega_j, \Omega_i \cap \Omega_j}(f_j)$$

for integers $i, j, \Omega_i \cap \Omega_j \neq \emptyset$, then there exists an $f \in \Omega$ such that $\rho_{\Omega, \Omega_i}(f) = f_i$ for any integer i .

By this definition, there is just one cosmos Ω and the sub-cosmos sequence is

$$\mathbf{R}^4 \supset \mathbf{R}^3 \supset \mathbf{R}^2 \supset \mathbf{R}^1 \supset \mathbf{R}^0 = \{P\} \supset \mathbf{R}_7^- \supset \cdots \supset \mathbf{R}_1^- \supset \mathbf{R}_0^- = \{Q\}.$$

in the string/M-theory. In Fig.4.2, we have shown the idea of the combinatorial cosmos.




Fig.4.2

For 5 or 6 dimensional spaces, it has been established a dynamical theory by this combinatorial speculation([24]-[25]). In this dynamics, we look for a solution in the Einstein's equation of gravitational field in 6-dimensional spacetime with a metric of the form

$$ds^2 = -n^2(t, y, z)dt^2 + a^2(t, y, z)d\sum_k^2 + b^2(t, y, z)dy^2 + d^2(t, y, z)dz^2$$

where $d\sum_k^2$ represents the 3-dimensional spatial sections metric with $k = -1, 0, 1$ respective corresponding to the hyperbolic, flat and elliptic spaces. For 5-dimensional spacetime, deletes the indefinite z in this metric form. Now consider a 4-brane moving in a 6-dimensional *Schwarzschild-ADS spacetime*, the metric can be written as

$$ds^2 = -h(z)dt^2 + \frac{z^2}{l^2}d\sum_k^2 + h^{-1}(z)dz^2,$$

where

$$d\sum_k^2 = \frac{dr^2}{1 - kr^2} + r^2d\Omega_{(2)}^2 + (1 - kr^2)dy^2$$

and

$$h(z) = k + \frac{z^2}{l^2} - \frac{M}{z^3}.$$

Then the equation of a 4-dimensional cosmos moving in a 6-spacetime is

$$2\frac{\ddot{R}}{R} + 3\left(\frac{\dot{R}}{R}\right)^2 = -3\frac{\kappa_{(6)}^4}{64}\rho^2 - \frac{\kappa_{(6)}^4}{8}\rho p - 3\frac{\kappa}{R^2} - \frac{5}{l^2}$$

by applying the *Darmois-Israel conditions* for a moving brane. Similarly, for the case of $a(z) \neq b(z)$, the equations of motion of the brane are

$$\frac{d^2 \dot{R} - d\ddot{R}}{\sqrt{1 + d^2 \dot{R}^2}} - \frac{\sqrt{1 + d^2 \dot{R}^2}}{n} \left(d\dot{n}\dot{R} + \frac{\partial_z n}{d} - (d\partial_z n - n\partial_z d)\dot{R}^2 \right) = -\frac{\kappa_{(6)}^4}{8} (3(p + \rho) + \hat{p}),$$

$$\frac{\partial_z a}{ad} \sqrt{1 + d^2 \dot{R}^2} = -\frac{\kappa_{(6)}^4}{8} (\rho + p - \hat{p}),$$

$$\frac{\partial_z b}{bd} \sqrt{1 + d^2 \dot{R}^2} = -\frac{\kappa_{(6)}^4}{8} (\rho - 3(p - \hat{p})),$$

where the energy-momentum tensor on the brane is

$$\hat{T}_{\mu\nu} = h_{\nu\alpha} T_{\mu}^{\alpha} - \frac{1}{4} T h_{\mu\nu}$$

with $T_{\mu}^{\alpha} = \text{diag}(-\rho, p, p, p, \hat{p})$ and the *Darmois-Israel conditions*

$$[K_{\mu\nu}] = -\kappa_{(6)}^2 \hat{T}_{\mu\nu},$$

where $K_{\mu\nu}$ is the extrinsic curvature tensor.

The combinatorial cosmos also presents new questions to combinatorics, such as:

- (i) to embed a graph into spaces with dimensional ≥ 4 ;
- (ii) to research the phase space of a graph embedded in a space;
- (iii) to establish graph dynamics in a space with dimensional ≥ 4 , \dots , etc..

For example, we have gotten the following result for graphs in spaces in [17].

Theorem 4.1 *A graph G has a nontrivial including multi-embedding on spheres $P_1 \supset P_2 \supset \dots \supset P_s$ if and only if there is a block decomposition $G = \bigsqcup_{i=1}^s G_i$ of G such that for any integer $i, 1 < i < s$,*

- (i) G_i is planar;
- (ii) for $\forall v \in V(G_i)$, $N_G(x) \subseteq \left(\bigcup_{j=i-1}^{i+1} V(G_j) \right)$.

Further research of the combinatorial cosmos will richen the knowledge of combinatorics and cosmology, also get the combinatorialization for cosmology.

References

- [1] I.Antoniadis, Physics with large extra dimensions: String theory under experimental test, *Current Science*, Vol.81, No.12, 25(2001),1609-1613
- [2] S.S.Chern and W.H.Chern, *Lectures in Differential Geometry*(in Chinese), Peking University Press, 2001.
- [3] M.J.Duff, A layman's guide to M-theory, *arXiv*: hep-th/9805177, v3, 2 July(1998).
- [4] B.J.Fei, *Relativity Theory and Non-Euclid Geometries*, Science Publisher Press, Beijing, 2005.

- [5] U.Günther and A.Zhuk, Phenomenology of brane-world cosmological models, *arXiv: gr-qc/0410130*.
- [6] S.Hawking, *A Brief History of Times*, A Bantam Books/November, 1996.
- [7] S.Hawking, *The Universe in Nutshell*, A Bantam Books/November, 2001.
- [8] D.Ida, Brane-world cosmology, *arXiv: gr-qc/9912002*.
- [9] H.Iseri, *Smarandache Manifolds*, American Research Press, Rehoboth, NM,2002.
- [10] H.Iseri, *Partially Paradoxist Smarandache Geometries*, <http://www.gallup.unm.edu/~smarandache/Howard-Iseri-paper.htm>.
- [11] M.Kaku, *Hyperspace: A Scientific Odyssey through Parallel Universe, Time Warps and 10th Dimension*, Oxford Univ. Press.
- [12] P.Kanti, R.Madden and K.A.Olive, A 6-D brane world model, *arXiv: hep-th/0104177*.
- [13] L.Kuciuk and M.Antholy, An Introduction to Smarandache Geometries, *Mathematics Magazine, Aurora, Canada*, Vol.12(2003)
- [14] L.F.Mao, *On Automorphisms groups of Maps, Surfaces and Smarandache geometries, Scientia Magna*, Vol.1(2005), No.2, 55-73.
- [15] L.F.Mao, A new view of combinatorial maps by Smarandache's notion, in *Selected Papers on Mathematical Combinatorics(I)*, World Academic Union, 2006, also in *arXiv: math.GM/0506232*.
- [16] L.F.Mao, *Automorphism Groups of Maps, Surfaces and Smarandache Geometries*, American Research Press, 2005.
- [17] L.F.Mao, *Smarandache multi-space theory*, Hexis, Phoenix, AZ2006.
- [18] L.F.Mao, On algebraic multi-groups, *Scientia Magna*, Vol.2(2006), No.1, 64-70.
- [19] L.F.Mao, On multi-metric spaces, *Scientia Magna*, Vol.2(2006), No.1, 87-94.
- [20] L.F.Mao, Geometrical theory on combinatorial manifolds, *JP J.Geometry and Topology*, Vol.7, No.1(2007),65-114.
- [21] L.F.Mao, An introduction to Smarandache multi-spaces and mathematical combinatorics, *Scientia Magna*, Vol.3, No.1(2007), 54-80.
- [22] L.F.Mao, A combinatorially generalized Stokes' theorem on integrations, *International J.Mathematical Combinatorics*, Vol.1, No.1(2007),67-86.
- [23] L.F.Mao, Curvature equations on combinatorial manifolds with applications to theoretical physics, *International J.Mathematical Combinatorics* (accepted).
- [24] E.Papantonopoulos, Braneworld cosmological models, *arXiv: gr-qc/0410032*.
- [25] E.Papantonopoulos, Cosmology in six dimensions, *arXiv: gr-qc/0601011*.
- [26] J.A.Peacock, *Cosmological Physics*, Cambridge University Press, 2003.
- [27] J.Polchinski, *String Theory*, Vol.1-Vol.2, Cambridge University Press, 2003.
- [28] D.Rabounski, Declaration of academic freedom: Scientific Human Rights, *Progress in Physics*, January (2006), Vol.1, 57-60.
- [29] F.Smarandache, Mixed non-euclidean geometries, *eprint arXiv: math/0010119*, 10/2000.
- [30] F.Smarandache, *A Unifying Field in Logics. Neutrosopy: Neutrosophic Probability, Set, and Logic*, American research Press, Rehoboth, 1999.
- [31] F.Smarandache, Neutrosophy, a new Branch of Philosophy, *Multi-Valued Logic*, Vol.8, No.3(2002)(special issue on Neutrosophy and Neutrosophic Logic), 297-384.

- [32] F.Smarandache, A Unifying Field in Logic: Neutrosophic Field, *Multi-Valued Logic*, Vol.8, No.3(2002)(special issue on Neutrosophy and Neutrosophic Logic), 385-438.
- [33] J.Stillwell, *Classical topology and combinatorial group theory*, Springer-Verlag New York Inc., (1980).