



On Independence Neutrosophic Random Variables

Carlos Granados¹ and José Sanabria²

¹Estudiante de Doctorado en Matemáticas, Universidad de Antioquia, Medellín, Colombia.

Email: carlosgranadosortiz@outlook.es

²Departamento de Matemáticas, Universidad de Sucre, Sincelejo, Colombia.

Email: jesanabri@gmail.com

Abstract. In this article we study independence neutrosophic random variables and conditioned expectation, we prove that conditional variance is equal to neutrosophic conditional variance. These notions are important for neutrosophic probability due to neutrosophic random variables, neutrosophic central limit theorem and neutrosophic laws of large numbers can be studied.

Keywords: Neutrosophic random vector; neutrosophic expectation; independence neutrosophic random variables; Neutrosophic Logic.

1. Introduction

F. Smarandache introduced the notion of neutrosophic probability measure as a function $\mathcal{NP} : Y \rightarrow [0, 1]^3$ where Y is a neutrosophic sample space, and introduced the probability mapping to take the form $\mathcal{NP}(B) = (ch(B), ch(neutB), ch(antiB)) = (\alpha, \beta, \gamma)$ where $0 \leq \alpha, \beta, \gamma \leq 1$ and $0 \leq \alpha + \beta + \gamma \leq 3$ [33]. Moreover, many researchers have investigated many neutrosophic probability distributions like Poisson, exponential, binomial, normal, uniform, Weibull,...etc. (See [32], [2], [18], [26]). Furthermore, researchers have introduced the notion of neutrosophic queueing theory in [35], [36] this is one branch of neutrosophic stochastic modelling. Besides, researchers have also investigated neutrosophic time series prediction and modelling in many cases like neutrosophic moving averages, neutrosophic logarithmic models, neutrosophic linear models and so on. [3], [4], [12].

On the other hand, neutrosophic logic is an extension of intuitionistic fuzzy logic by adding indeterminacy component (I) where $I^2 = I, \dots, I^n = I, 0.I = 0; n \in \mathbb{N}$ and I^{-1} is undefined (see [20], [32]). Neutrosophic logic has a huge brand of applications in many fields including

decision making [29], [19], [25], machine learning [6], [27], intelligent disease diagnosis [30], [11], communication services [8], pattern recognition [28], social network analysis and e-learning systems [21], physics [34], sequences spaces [14] and so on. Neutrosophic logic has solved many decision-making problems efficiently like evaluating green credit rating, personnel selection, etc. [22], [23], [24], [1]. For more notions related to neutrosophic theory, we refer the reader to [9, 10, 14–17].

The study of neutrosophic random variables has become one of the fundamental pillars in neutrosophic theory and probability. Recent results of great importance can be seen in [37] and [13]. Taking into account mentioned above, in this article we study independent neutrosophic random variables and conditioned expectation.

2. Preliminaries

In this section, we show some well-known definitions and properties of neutrosophic logic and neutrosophic probability which are useful for the development of this paper.

Definition 2.1. (see [31]) Let X be a non-empty fixed set. A neutrosophic set A is an object having the form $\{x, (\mu A(x), \delta A(x), \gamma A(x)) : x \in X\}$, where $\mu A(x)$, $\delta A(x)$ and $\gamma A(x)$ represent the degree of membership, the degree of indeterminacy, and the degree of non-membership respectively of each element $x \in X$ to the set A .

Definition 2.2. (see [5]) Let K be a field, the neutrosophic field generated by K and I is denoted by $\langle K \cup I \rangle$ under the operations of K , where I is the neutrosophic element with the property $I^2 = I$.

Definition 2.3. (see [32]) Classical neutrosophic number has the form $a + bI$ where a, b are real or complex numbers and I is the indeterminacy such that $0.I = 0$ and $I^2 = I$ which results that $I^n = I$ for all positive integers n .

Definition 2.4. (see [33]) The neutrosophic probability of event A occurrence is $NP(A) = (ch(A), ch(neutA), ch(antiA)) = (T, I, F)$ where T, I, F are standard or non-standard subsets of the non-standard unitary interval $]^{-0, 1^+}$.

Recently, Bisher and Hatip [37] introduced and studied the notions of neutrosophic random variables by using the concepts presented by [33], these notions were defined as follows:

Definition 2.5. Consider the real valued crisp random variable X which is defined as follows:

$$X : \Omega \rightarrow \mathbb{R}$$

where Ω is the events space. Now, they defined a neutrosophic random variable X_N as follows:

$$X_N : \Omega \rightarrow \mathbb{R}(I)$$

and

$$X_N = X + I$$

where I is indeterminacy.

Theorem 2.6. Consider the neutrosophic random variable $X_N = X + I$ where cumulative distribution function of X is $F_X(x) = P(X \leq x)$. Then, the following statements hold:

- (1) $F_{X_N}(x) = F_X(x - I)$,
- (2) $f_{X_N}(x) = f_X(x - I)$.

Where F_{X_N} and f_{X_N} are cumulative distribution function and probability density function of X_N , respectively.

Theorem 2.7. Consider the neutrosophic random variable $X_N = X + I$, expected value can be found as follows:

$$E(X_N) = E(X) + I.$$

Proposition 2.8 (Properties of expected value of a neutrosophic random variable). Let X_N and Y_N be neutrosophic random variables, then the following properties holds:

- (1) $E(aX_N + b + cI) = aE(X_N) + b + cI$; $a, b, c \in \mathbb{R}$,
- (2) If X_N and Y_N are neutrosophic random variables, then $E(X_N \pm E(Y_N)) = E(X_N) \pm E(Y_N)$,
- (3) $E[(a + bI)X_N] = aE(X_N) + bIE(X_N)$; $a, b \in \mathbb{R}$,
- (4) $|E(X_N)| \leq E|X_N|$.

Theorem 2.9. Consider the neutrosophic random variable $X_N = X + I$, variance of X_N is equal to variance of X , i.e. $V(X_N) = V(X)$.

Now, Granados [13] studied the notions of neutrosophic random vector and joint neutrosophic random variable, these notions were defined as follows:

Definition 2.10. A neutrosophic random vector of two dimension is a vector (X_N, Y_N) in which each coordinate is a neutrosophic random variable. Analogously, we can define a neutrosophic random vector multidimensional as follows $(X_{N_1}, X_{N_2}, \dots, X_{N_n})$ in which $X_{N_1}, X_{N_2}, \dots, X_{N_n}$ are neutrosophic random variables for each $n = 1, 2, \dots$

Definition 2.11. Let (X_N, Y_N) be a neutrosophic random vector, we define probability function of a neutrosophic continuous random vector (X_N, Y_N) . Then, joint probability neutrosophic function of a discrete random vector (X_N, Y_N) $f_N(x, y) : \mathbb{R}^2 \rightarrow [0, \infty)$ in which is non-negative and integrable, and for any $(x, y) \in \mathbb{R}^2$, it is defined as follows

$$P(X_N \leq x, Y_N \leq y) = P(X \leq x - I, Y \leq y - I) = \int_{-\infty}^{y-I} \int_{-\infty}^{x-I} f_{(X_N, Y_N)}(u, v) dv du$$

Similarly, probability function of a neutrosophic discrete random vector (X_N, Y_N) is defined similar by using sum.

Definition 2.12. Let (X_N, Y_N) be a neutrosophic random vector, we define neutrosophic joint distribution function which will be denoted by $F_{(X_N, Y_N)}(x, y) = P(X_N \leq x, Y_N \leq y) = P(X \leq x - I, Y \leq y - I)$.

Definition 2.13. Let $f_{(X_N, Y_N)}(x, y)$ be a joint probability neutrosophic function of a continuous random variable (X_N, Y_N) . We define neutrosophic marginal function of X_N as follows:

$$f_{X_N}(x) = \int_{-\infty}^{+\infty} f_{(X_N, Y_N)}(x, y) dy$$

and we define neutrosophic marginal function of Y_N as follows:

$$f_{Y_N}(y) = \int_{-\infty}^{+\infty} f_{(X_N, Y_N)}(x, y) dx$$

Similarly, joint probability neutrosophic function of a discrete random variable is defined similar by using sum.

Definition 2.14. Expected value of a neutrosophic random vector (X_N, Y_N) in which expected value of X_N and Y_N exist, we define $E(X_N, Y_N) = (E(X_N), E(Y_N))$.

Next, we will show some new notions on neutrosophic random variables which have not been studied so far and are needed.

Let (X_N, Y_N) be neutrosophic vector and $\phi : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a function, then $\phi(X_N, Y_N)$ is a neutrosophic random variable and its expectation is defined as follows:

$$E[\phi(X_N, Y_N)] = \int_{-\infty}^{+\infty} (x - I) dF_{\phi(X_N, Y_N)}(x),$$

as well as one dimensional case, we are required to find out the distribution $\phi(X_N, Y_N)$ by which can be difficult in some cases. Next, we establish an alternative way in which we can calculate expectation of $\phi(X_N, Y_N)$ without known its distribution, but we must know (X_N, Y_N) distribution. Let (X_N, Y_N) be neutrosophic vector and $\phi : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a function such that $\phi(X_N, Y_N)$ has finite expected, then:

$$E[\phi(X_N, Y_N)] = \int_{\mathbb{R}^2} \phi(x - I, y - I) dF_{\phi(X_N, Y_N)}(x, y),$$

This can be proved easily due to this is a Riemann-Stieltjes integral in two dimensional.

In case of X_N and Y_N be independence (see section 3), this increment is

$$\begin{aligned} &F_X(x_i - I)F_Y(y_j - I) - F_X(x_i - I)F_Y(y_{j-1} - I) - F_X(x_{i-1} - I)F_Y(y_j - I) \\ &+ F_X(x_i - I)F_Y(y_j - I) = \Delta F_X(x_i - I)\Delta F_Y(y_j - I), \end{aligned}$$

i.e, bidimensional integral can be separated in two integral and can be written as follows

$$E[\phi(X_N, Y_N)] = \int_{\mathbb{R}^2} \phi(x - I, y - I) dF_{X_N}(x) dF_{Y_N}(y).$$

When (X_N, Y_N) is a discrete vector, we have

$$E[\phi(X_N, Y_N)] = \sum_{x-I, y-I} \phi(x - I, y - I) P(X = x - I, Y = y - I),$$

in which the sum is applied over all possible valued $(x - I, y - I)$ on (X_N, Y_N) .

Theorem 2.15. *Let X_N and Y_N be two neutrosophic random variable with finite expectation, then*

$$E(X_N + Y_N) = E(X_N) + E(Y_N).$$

Proof. Let $\phi(x - I, y - I) = x - I + y - I$, $\phi_1(x - I, y - I) = x - I$ and $\phi_2(x - I, y - I) = y - I$. Then,

$$\begin{aligned} E(X_N + Y_N) &= E(\phi(X_N, Y_N)) \\ &= \int_{\mathbb{R}^2} (x - I + y - I) dF_{X_N, Y_N}(x, y) \\ &= \int_{\mathbb{R}^2} (x - I) dF_{X_N, Y_N}(x, y) + \int_{\mathbb{R}^2} (y - I) dF_{X_N, Y_N}(x, y) \\ &= E(\phi_1(X_N, Y_N)) + E(\phi_2(X_N, Y_N)) \\ &= E(X_N) + E(Y_N) \end{aligned}$$

□

Theorem 2.16. *Let X_N and Y_N be two independence neutrosophic random variable and g and h be two functions such that $g(X_N)$ and $h(Y_N)$ have finite expected, then*

$$E[g(X_N)h(Y_N)] = E[g(X_N)]E[h(Y_N)].$$

Proof.

$$\begin{aligned} E[g(X_N)h(Y_N)] &= \int_{\mathbb{R}^2} g(x - I)h(y - I) dF_{X_N, Y_N}(x, y) \\ &= \int_{\mathbb{R}^2} g(x - I)h(y - I) dF_{X_N}(x) dF_{Y_N}(y) \\ &= \int_{\mathbb{R}} g(x - I) dF_{X_N}(x) \int_{\mathbb{R}} h(y - I) dF_{Y_N}(y) \\ &= E[g(X_N)]E[h(Y_N)] \end{aligned}$$

□

3. Independence neutrosophic random variables

Let X_N and Y_N be two neutrosophic random variables. X_N and Y_N are independence if the events $(X \leq x - I)$ and $(Y \leq y - I)$ are independence for any real-value x and y , i.e., if the following equality is satisfied

$$P[(X \leq x - I) \cap (Y \leq y - I)] = P(X \leq x - I)P(Y \leq y - I). \quad (1)$$

The left side of the equality (1) can be written as $P(X \leq x - I, Y \leq y - I)$ or $F_{X,Y}(x - I, y - I)$, and it is said to be the joint distribution function of X_N and Y_N evaluate in the point (x, y) . Therefore, note that (1) can be expressed as follows

$$F_{X,Y}(x - I, y - I) = F_X(x - I)F_Y(y - I), \text{ for } x, y \in \mathbb{R}. \quad (2)$$

In this way, in order to determine whether two neutrosophic random variables are independent, it is necessary to know both the joint probabilities $P(X \leq x - I, Y \leq y - I)$ as individual probabilities $P(X \leq x - I)$ and $P(Y \leq y - I)$, and verify the identity (2) for each real numbers x and y . Hence, it is enough that there exists a pair $(x - I, y - I)$ for which the equality (2) does not hold to be able to conclude that X and Y are not independent. Granados [13] studied on random vectors and explained how to obtain the individual distributions from the joint distribution of two random variables.

Example 3.1. Let (X_N, Y_N) be a neutrosophic random vector with density function $f(x - I, y - I) = 4(x - I)(y - I)$ for $I \leq x, y \leq 1 - I$.

The marginal density function of X_N is calculated as follows for $I \leq x \leq 1 - I$

$$f_X(x - I) = \int_I^{1-I} 4(x - I)(y - I)dy = 2(x - I).$$

Analogously, we can prove that $f_{Y_N}(y) = 2(y - I)$ for $I \leq x \leq 1 - I$. Therefore, X_N and Y_N are independence, due to $F_{X,Y}(x - I, y - I) = F_X(x - I)F_Y(y - I)$.

Proposition 3.2. Let X_N and Y_N be two independence neutrosophic random variables, and let g and h be two functions of $\mathbb{R} \rightarrow \mathbb{R}$. Then, the neutrosophic random variables $g(X_N)$ and $h(Y_N)$ are independence neutrosophic random variables.

Proof: Let $\mathcal{A} = (-\infty, x - I]$ and $\mathcal{B} = (-\infty, y - I]$, then

$$\begin{aligned} P(g(X_N) \leq x, h(Y_N) \leq y) &= P(g(X_N) \in \mathcal{A}, h(Y_N) \in \mathcal{B}) \\ &= P(X_N \in g^{-1}(\mathcal{A}), Y_N \in h^{-1}(\mathcal{B})) \\ &= P(X_N \in g^{-1}(\mathcal{A}))P(Y_N \in h^{-1}(\mathcal{B})) \\ &= P(g(X_N) \in \mathcal{A})P(h(Y_N) \in \mathcal{B}) \\ &= P(g(X_N) \leq x)P(h(Y_N) \leq y) \end{aligned}$$

Theorem 3.3. Let X_N and Y_N two discrete neutrosophic random variables, then

(1) Using $P(X \leq x - I, Y \leq y - I) = \sum_{u \leq x - I} \sum_{v \leq y - I} P(X = u, Y = v)$, we have

$$\begin{aligned} P(X = x - I, Y = y - I) &= P(X \leq x - I, Y \leq y - I) \\ &\quad - P(X \leq x - I - 1, Y \leq y - I) \\ &\quad - P(X \leq x - I, Y \leq y - I - 1) \\ &\quad + P(X \leq x - I - 1, Y \leq y - I - 1). \end{aligned}$$

(2) Using (1), independence condition $P(X \leq x - I, Y \leq y - I) = P(X \leq x - I)P(Y \leq y - I)$ is equivalent to $P(X = x - I, Y = y - I) = P(X = x - I)P(Y = y - I)$.

Proof:

(1) This result can be obtained by using the following equality

$$\begin{aligned} (X = x - I, Y = y - I) &= (X \leq x - I, Y \leq y - I) - (X \leq x - I - 1, Y \leq y - I) \\ &\quad - (X \leq x - I, Y \leq y - I - 1) \\ &\quad + (X \leq x - I - 1, Y \leq y - I - 1). \end{aligned}$$

(2) Using (1) and by hypothesis of independence, we have

$$\begin{aligned} P(X = x - I, Y = y - I) &= P(X \leq x - I)P(Y \leq y - I) - \\ &\quad P(X \leq x - I - 1)P(Y \leq y - I) \\ &\quad - P(X \leq x - I)P(Y \leq y - I - 1) \\ &\quad + P(X \leq x - I - 1)P(Y \leq y - I - 1) \\ &= [P(X \leq x - I) - P(X \leq x - I - 1)] \\ &\quad \times [P(Y \leq y - I) - P(Y \leq y - I - 1)] \\ &= P(X = x - I)P(Y = y - I). \end{aligned}$$

Theorem 3.4. Let X_N and Y_N be two independence neutrosophic random variables which have finite expectation. Then,

$$E(X_N Y_N) = E(X_N)E(Y_N). \quad (3)$$

Proof: We prove the case when X_N and Y_N are discrete neutrosophic random variables, the case for continuous neutrosophic random variables are proved similarly.

Let

$$\begin{aligned}
 E(X_N Y_N) &= \sum_{x,y} (x - I)(y - I)P(X_N = x, Y_N = y) \\
 &= \sum_x \sum_y (x - I)(y - I)P(X_N = x, Y_N = y) \\
 &= \left(\sum_x (x - I)P(X_N = x)\right)\left(\sum_y (y - I)P(Y_N = y)\right) \\
 &= E(X_N)E(Y_N).
 \end{aligned}$$

Definition 3.5. Let $X_{N_1}, X_{N_2}, \dots, X_{N_n}$ be a collection of neutrosophic random variables with joint function distribution $F_X(x_1 - I, x_2 - I, \dots, x_n - I)$, and consider marginals functions distribution $F_{X_{N_1}}(x_1), F_{X_{N_2}}(x_2), \dots, F_{X_{N_n}}(x_n)$, respectively. Then, we say that $X_{N_1}, X_{N_2}, \dots, X_{N_n}$ are independence if for any real numbers $x_1 - I, x_2 - I, \dots, x_n - I$ the following equality holds

$$F_X(x_1 - I, x_2 - I, \dots, x_n - I) = F_{X_{N_1}}(x_1)F_{X_{N_2}}(x_2)\dots F_{X_{N_n}}(x_n).$$

Analogously, we can define it in terms of neutrosophic density function $f(x_1 - I, x_2 - I, \dots, x_n - I)$ if the following equality holds

$$f(x_1 - I, x_2 - I, \dots, x_n - I) = f_{X_{N_1}}(x_1)f_{X_{N_2}}(x_2)\dots f_{X_{N_n}}(x_n).$$

Example 3.6. Let X_N and Y_N be two continuous neutrosophic random variables with joint density function

$$f_{(X,Y)}(x - I, y - I) = \begin{cases} e^{-x-y+2I} & \text{if } x, y > I, \\ 0 & \text{otherwise.} \end{cases}$$

Neutrosophic marginals probability functions are defined as follows

$$f_X(x - I) = \begin{cases} e^{-x+I} & \text{if } x > I, \\ 0 & \text{otherwise.} \end{cases}$$

and

$$f_Y(y - I) = \begin{cases} e^{-y+I} & \text{if } y > I, \\ 0 & \text{otherwise.} \end{cases}$$

Therefore, $f_{(X,Y)}(x - I, y - I) = f_X(x - I)f_Y(y - I)$ for any real numbers $x - I$ and $y - I$, and hence we conclude X_N and Y_N are independence.

Example 3.7. Let X_N and Y_N be two discrete neutrosophic random variables with joint density function

$$f_{(X,Y)}(x - I, y - I) = \begin{cases} \frac{1}{4} & \text{if } x, y \in \{I, 1 + I\}, \\ 0 & \text{otherwise.} \end{cases}$$

Neutrosophic marginals probability functions are defined as follows

$$f_X(x - I) = \begin{cases} \frac{1}{2} & \text{if } x \in \{I, 1 + I\}, \\ 0 & \text{otherwise.} \end{cases}$$

and

$$f_Y(y - I) = \begin{cases} \frac{1}{2} & \text{if } y \in \{I, 1 + I\}, \\ 0 & \text{otherwise.} \end{cases}$$

Therefore, $f_{(X,Y)}(x - I, y - I) = f_X(x - I)f_Y(y - I)$ for any real numbers $x - I$ and $y - I$, and hence we conclude X_N and Y_N are independence.

Remark 3.8. It can be said that an infinite set of neutrosophic random variables is independence if any finite subset is independence.

This statement can be useful due to for future work the concepts of neutrosophic central limit theorem and neutrosophic laws of large numbers can be studied.

4. Conditional expectation

In this section, we introduce the concept of conditional expectation of a neutrosophic random variable with respect to a σ -algebra, and some of its elemental properties are studied. We will consider that has a base probability space (Ω, \mathcal{F}, P) , and \mathcal{G} is a sub-algebra of \mathcal{F} . We have defined expectation of a neutrosophic random variable as a Riemann-Stieltjes integral as follows

$$E(X_N) = \int_{-\infty}^{+\infty} (x - I) dF_{X_N}(x).$$

however, to make the notation simpler in this section, it is sometimes convenient to adopt the notation of measure theory and denote the expectation of a neutrosophic random variable X_N as follows

$$E(X_N) = \int_{\Omega} X_N dP.$$

We shall recall that if we know distribution of neutrosophic vector (X_N, Y_N) and we take valued $y - I$ such that $f_Y(y - I) = f_{Y_N}(y) \neq 0$, conditional expectation of X_N known $Y = y - I$

is the function

$$y - I \mapsto E(X_N|Y_N = y) = \int_{-\infty}^{+\infty} (x - I) dF_{X_N|Y_N}(x|y), \quad (4)$$

when $f_{Y_N}(y) \neq 0$. (4) is equivalent to write

$$E(X_N|Y_N = y) = \int_{-\infty}^{+\infty} (x - I) f_{X_N|Y_N}(x|y) dx.$$

If we make a change in order of integration, we can see that

$$E(X_N) = \int_{-\infty}^{+\infty} E(X_N|Y_N = y) f_{Y_N}(y) dy,$$

if we apply this expression by using total probability theorem in terms of expectation. In the case when (X_N, Y_N) is a discrete neutrosophic vector, we have

$$\begin{aligned} E(X_N|Y_N = y) &= \sum_x (x - I) f_{X_N|Y_N}(x|y) \\ &= \sum_x (x - I) P(X_N = x|Y_N = y), \end{aligned}$$

considering $f_{Y_N}(y) \neq 0$ and sum is absolutely convergent. Again, applying a change in order of sum, we have

$$E(X_N) = \sum_y E(X_N|Y_N = y) P(Y_N = y).$$

In any cases, we can also see, when Y_N and X_N are independence, we have

$$E(X_N|Y_N = y) = E(X_N).$$

Example 4.1. We will find expectation of $E(X_N|Y_N = y)$ for each $y \in (I, 1 + I)$ when X_N and Y_N have the following joint neutrosophic density function.

$$f_{X|Y}(x - I, y - I) = \begin{cases} 12(x - I)^2 & \text{if } I < x < y < 1 + I, \\ 0 & \text{otherwise.} \end{cases}$$

For $I < y < 1 + I$,

$$\begin{aligned}
 E(X_N|Y_N = y) &= \int_0^{y-I} (x - I) \frac{12(x - I)^2}{4(y - I)^3} dx \\
 &= \frac{3}{(y - I)^3} \int_0^{y-I} (x - I)^3 dx \\
 &= \frac{3}{4} \frac{(y - 2I)^4 - I}{(y - I)^3}
 \end{aligned}$$

Analogously to (4), if Q is an event with positive probability and X_N is a integrable neutrosophic random variable, conditional expectation of X_N known Q is

$$E(X_N|Q) = \int_{-\infty}^{+\infty} (x - I) dF_{X_N|Q}(x),$$

where $F_{X_N|Q}(x) = P(X_N \leq x|A) = P(X \leq x - I|A) = \frac{P(X \leq x - I, Q)}{P(Q)}$. Next, we will show a more generally definition which generalized concepts showed so far.

Definition 4.2. Let X_N be a neutrosophic random variable with finite expectation, and let \mathcal{G} be a sub-algebra of \mathcal{F} . Conditional expectation of X_N known \mathcal{G} , it is a neutrosophic random variable which will be denoted by $E(X_N|\mathcal{G})$ which satisfies the following conditions:

- (1) It is \mathcal{G} -measurable,
- (2) It has finite expectation,
- (3) For any event $G \in \mathcal{G}$, $\int_G E(X_N|\mathcal{G})dP = \int_G X_N dP$.

Remark 4.3. Just as it happens in a random variable, part of the difficulty in understanding this general definition is that an explicit formula is not provided for this neutrosophic random variable but only the properties it satisfies. The objective of this section is to find the meaning of this neutrosophic random variable, interpret its meaning, and explain its relationship with the concept of elementary conditional expectation.

Remark 4.4. When $\mathcal{G} = \sigma(Y_N)$ for any neutrosophic random variable Y_N , conditional expected will be written by $E(X_N|Y_N)$ instead of $E(X_N|\sigma(Y_N))$.

Remark 4.5. Let Q be any event, then conditional expected $E(1_Q|\mathcal{G})$ will be denoted by $P(Q|\mathcal{G})$.

Now, we will show some properties on $E(X_N|Y_N)$ when Y_N is a discrete neutrosophic random variable.

Let X_N and Y_N be two neutrosophic random variables. Now, consider that X_N has finite expected and Y_N is discrete with possible values $y_1 - I, y_2 - I, \dots$ Conditional expectation of

X_N known event $(Y_N = y_j) = (Y = y_j - I)$ is $E(X_N|Y_N = y_j)$, this value depends of the event $(Y_N = y_j)$, and we can consider that we have a function defined over Ω as follows: If ω is such that $Y(\omega) = y_j - I$, then

$$\omega \mapsto E(X_N|Y_N)(\omega) = E(X_N|Y_N = y_j).$$

We can see that $E(X_N|Y_N)$ takes at much different values as Y_N does. Generally, function can be rewritten in terms of indicator function as follows

$$E(X_N|Y_N)(\omega) = \sum_{j=1}^{\infty} E(X_N|Y_N = y_j)1_{(Y_N=y_j)}(\omega).$$

In this way, we can define the function $E(X_N|Y_N) : \Omega \rightarrow \mathbb{R}$ which is denoted by

$$\omega \mapsto E(X_N|Y_N)(\omega) = E(X_N|Y_N = y_j) \text{ if } Y(\omega) = y_j - I$$

is a neutrosophic random variable.

Theorem 4.6. *Let X_N be a integrable neutrosophic random variable, and let Y_N discrete with possible values $y_1 - I, y_2 - I, \dots$ $E(X_N|Y_N) : \Omega \rightarrow \mathbb{R}$ is a neutrosophic random variable which satisfies the following conditions:*

- (1) *It is $\sigma(Y_N)$ -measurable,*
- (2) *It has finite expected,*
- (3) *For any event $G \in \sigma(Y_N)$, $\int_G E(X_N|Y_N)dP = \int_G X_N dP$.*

Proof:

- (1) Through possibles values, the neutrosophic random variables Y_N divides in different part Ω , i.e $(Y_N = y_1), (Y_N = y_2), \dots$ are disjunct events. $\sigma(Y_N) = \sigma\{(Y_N = y_1), (Y_N = y_2), \dots\} \subset \mathcal{F}$. Since $E(X_N|Y_N)$ is constant in each element of partition, implies that $E(X_N|Y_N)$ is $\sigma(Y_N)$ -measurable, and hence it is a neutrosophic random variable.
- (2) Taking the event G as Ω in the third property, we get that X_N and $E(X_N|Y_N)$ have the same finite expectation.
- (3) Since each element of $\sigma(Y_N)$ is union of disjunct elements of $(Y_N = y_j)$, by properties of integral it is enough to show that $\int_G E(X_N|Y_N)dP = \int_G X_N dP$ for these simple

events. Then, we have

$$\begin{aligned} \int_{(Y_N=y_j)} E(X_N|Y_N)(\omega)dP(\omega) &= E(X_N|Y_N = y_j)P(Y_N = y_j) \\ &= \int_{\Omega} X_N(\omega)dP(\omega|Y_N = y_j)P(Y_N = y_j) \\ &= \int_{\Omega} X_N(\omega)dP(\omega, Y_N = y_j) \\ &= \int_{(Y_N=y_j)} X_N(\omega)dP(\omega). \end{aligned}$$

Remark 4.7. We have to see the different between $E(X_N|Y_N = y_j)$ and $E(X_N|Y_N)$. First term is a possible numerical value, second term is a neutrosophic random variable. However, both expressions are called conditional expectation. We will see next a particular case of this neutrosophic random variable. Besides, we will show that the conditional expectation can be seen as a generalization of the basic concept of conditional probability, and it can also be considered as a generalization of the concept of expectation.

Proposition 4.8. *Let X_N be a neutrosophic random variable with finite expectation, then $E(X_N|\{\emptyset, \Omega\}) = E(X_N) = E(X) + I$.*

Proof: This proofs follows since $E(X_N|\mathcal{G})$ is measurable respected to \mathcal{G} , and any measurable function respected to $\{\emptyset, \Omega\}$ is constant. Now, for any event $G \in \{\emptyset, \Omega\}$, $\int_G E(X_N|\{\emptyset, \Omega\})dP = \int_G X_NdP = \int_{\Omega} X_NdP = E(X_N) = E(X) + I$.

Theorem 4.9. *Let X_N and Y_N be two neutrosophic random variables with finite expectation and $b \in \mathbb{R}$. Then, the following statements hold:*

- (1) *If $X \geq I$, then $E(X_N|\mathcal{G}) \geq 0$,*
- (2) *$E(bX_N + Y_N|\mathcal{G}) = bE(X_N|\mathcal{G}) + E(Y_N|\mathcal{G})$,*
- (3) *If $X_N \leq Y_N$, then $E(X_N|\mathcal{G}) \leq E(Y_N|\mathcal{G})$,*
- (4) *$E(E(X_N|\mathcal{G})) = E(X_N)$,*
- (5) *If X_N is \mathcal{G} -measurable, then $E(X_N|\mathcal{G}) = X_N$ a.s. In particular, $E(b|\mathcal{G}) = b$,*
- (6) *If $\mathcal{G}_1 \subset \mathcal{G}_2$, then*

$$E(E(X_N|\mathcal{G}_1)|\mathcal{G}_2) = E(E(X_N|\mathcal{G}_2)|\mathcal{G}_1) = E(X_N|\mathcal{G}_1),$$

- (7) *$|E(X_N|\mathcal{G})| \leq E(|X_N||\mathcal{G})$,*
- (8) *$E|E(X_N|\mathcal{G})| \leq E(|X_N|)$.*

Proof:

- (1) Follows directly from definition and the fact that $X \geq I \equiv X_N \geq 0$.

(2) For all $G \in \mathcal{G}$, we have

$$\begin{aligned} \int_G E(bX_N + Y_N|\mathcal{G})dP &= \int_G (bX_N + Y_N)dP \\ &= b \int_G X_NdP + \int_G Y_NdP. \end{aligned}$$

Now,

$$\begin{aligned} \int_G [bE(X_N|\mathcal{G}) + E(Y_N|\mathcal{G})]dP &= b \int_G E(X_N|\mathcal{G})dP + \int_G E(Y_N|\mathcal{G})dP \\ &= b \int_G X_NdP + \int_G Y_NdP. \end{aligned}$$

(3) Follows directly from definition and the fact that $X_N \leq Y_N$.

(4) Taking $G = \Omega$, we get the equality.

(5) Since X_N is \mathcal{G} -measurable, three condition of the definition hold. Now, $\int_G E(X_N)dP =$

$\int_G E(X_N)dP$. Therefore, $X_N = E(X_N|\mathcal{G})$ a.s.

(6) For all $G \in \mathcal{G}_1 \subset \mathcal{G}_2$, we have

$$\begin{aligned} \int_G E(E(X_N|\mathcal{G}_1)|\mathcal{G}_2)dP &= \int_G E(X_N|\mathcal{G}_1)dP \\ &= \int_G X_NdP. \end{aligned}$$

Analogously,

$$\begin{aligned} \int_G E(E(X_N|\mathcal{G}_2)|\mathcal{G}_1)dP &= \int_G E(X_N|\mathcal{G}_2)dP \\ &= \int_G X_NdP. \end{aligned}$$

(7) Consider that $\int_G |E(X_N|\mathcal{G})|dP = |\int_G E(X_N|\mathcal{G})dP|$. Then,

$$\begin{aligned} |\int_G E(X_N|\mathcal{G})dP| &= |\int_G X_NdP| \\ &\leq \int_G |X_N|dP \\ &= \int_G E(|X_N||\mathcal{G})dP. \end{aligned}$$

(8) Proof follows from parts (4) and (7) of this theorem.

Definition 4.10. Let X_N be a neutrosophic random variable with finite second moment, and \mathcal{G} be a sub-algebra of \mathcal{F} . Conditional variance of X_N known \mathcal{G} which will be denoted by $Var(X_N|\mathcal{G})$, it is defined as a neutrosophic random variable as follows

$$Var(X_N|\mathcal{G}) = E[(X_N - E(X_N|\mathcal{G}))^2|\mathcal{G}].$$

We shall recall that neutrosophic conditional variance is not a number, it is a neutrosophic random variable. Therefore, the only way that we have the neutrosophic variance of a random variable from conditional variance is $Var(X_N) = Var(X) = Var(X_N|\{\emptyset, \Omega\})$.

On the other hand, we have that

$$\begin{aligned} Var(X_N|\mathcal{G}) &= Var(X + I|\mathcal{G}) \\ &= Var(X|\mathcal{G}) \\ &= E[(X - E(X|\mathcal{G}))^2|\mathcal{G}], \end{aligned}$$

this shows that conditional covariance is equal to neutrosophic conditional covariance.

5. Conclusion

In this article we study the notion of neutrosophic random variable taking into account the notions previously studied by [37] and [13]. These results are of great importance because convergence on neutrosophic random variables, neutrosophic central limit theorem and neutrosophic laws of large numbers can be studied. Secondly, this results can be applied in quality control, stochastic modeling, reliability theory, queueing theory, electrical engineering and so on.

Funding

This research received no external funding.

Conflicts of Interest

The author declares no conflict of interest.

References

- [1] M. Abdel-Basset, N. A. Nabeeh, H. A. El-Ghareeb and A. Aboelfetouh, Utilizing Neutrosophic Theory to Solve Transition Difficulties of IoT-Based Enterprises, *Enterprise Information Systems*, 14(9-10)(2019), 1304–1324.
- [2] R. Alhabib, M. M. Ranna, H. Farah and A. Salama, Some Neutrosophic Probability Distributions, *Neutrosophic Sets and Systems*, 22(2018), 30–38.
- [3] R. Alhabib and A. A. Salama, The Neutrosophic Time Series-Study Its Models (Linear-Logarithmic) and test the Coefficients Significance of Its linear model, *Neutrosophic Sets and Systems*, 33(2020), 105–115.
- [4] R. Alhabib and A. A. Salama, Using Moving Averages To Pave The Neutrosophic Time Series, *International Journal of Neutrosophic Science*, 3(1)(2020), 14–20.
- [5] M. Ali, F. Smarandache, M. Shabir and L. Vladareanu, Generalization of Neutrosophic Rings and Neutrosophic Fields, *Neutrosophic Sets and Systems*, 5(2014), 9–14.
- [6] J. Anuradha and V. S, Neutrosophic Fuzzy Hierarchical Clustering for Dengue Analysis in Sri Lanka, *Neutrosophic Sets and Systems*, 31(2020), 179–199.
- [7] M. Bisher and A. Hatip, Neutrosophic Random variables, *Neutrosophic Sets and Systems*, 39(2021), 45–52.

Carlos Granados and José Sanabria, On Independence Neutrosophic Random Variables

- [8] A. Chakraborty, B. Banik, S. P. Mondal and S. Alam, Arithmetic and Geometric Operators of Pentagonal Neutrosophic Number and its Application in Mobile Communication Service Based MCGDM Problem, *Neutrosophic Sets and Systems*, 32(2020), 61–79.
- [9] S. Das, R. Das, C. Granados, A. Mukherjee, Pentapartitioned neutrosophic Q-ideals of Q-algebra, *Neutrosophic Sets and Systems* 41(2021), 53-63.
- [10] S. Das, R. Das, C. Granados, Topology on quadripartitioned neutrosophic sets, *Neutrosophic Sets and Systems* 45(2021), 54-61.
- [11] O. A. Ejaita and P. Asagba ,An Improved Framework for Diagnosing Confusable Diseases Using Neutrosophic Based Neural Network, *Neutrosophic Sets and Systems*, 16(2017), 28–34.
- [12] L. Esther Valencia Cruzaty, M. Reyes Tomal and C. Manuel Castillo Gallo, A Neutrosophic Statistic Method to Predict Tax Time Series in Ecuador, *Neutrosophic Sets and Systems*, 34(2020), 33–39.
- [13] C. Granados, New results on neutrosophic random variables. *Neutrosophic Sets and Systems* (2021).
- [14] C. Granados and A. Dhital, Statistical Convergence of Double Sequences in Neutrosophic Normed Spaces, *Neutrosophic Sets and Systems*, 42(2021), 333–344.
- [15] C. Granados, A. Dhital, New results on Pythagorean neutrosophic open sets in Pythagorean neutrosophic topological spaces, *Neutrosophic Sets and Systems* 43(2021), 12-23.
- [16] C. Granados, Una nueva noció de conjuntos neutrosóficos a través de los conjuntos $*b$ -abiertos en espacios topológicos neutrosóficos, *Eco Matemático* 12(2)(2021), 1-12.
- [17] C. Granados, Un nuevo estudio de los conjuntos supra neutrosophic crisp, *Revista Facultad de Ciencias Básicas* 16(2)(2020), 65-75.
- [18] K. Hamza Alhasan and F. Smarandache, Neutrosophic Weibull distribution and Neutrosophic Family Weibull Distribution, *Neutrosophic Sets and Systems*, 28(2019), 191–199.
- [19] H. Kamaci, Neutrosophic Cubic Hamacher Aggregation Operators and Their Applications in Decision Making, *Neutrosophic Sets and Systems*, 33(2020), 234–255.
- [20] W. B. V. Kandasamy and F. Smarandache, *Neutrosophic Rings*, Hexis, Phoenix, Arizona: Infinite Study, 2006.
- [21] M. M. Lotfy, S. ELhafeez, M. Eisa and A. A. Salama, Review of Recommender Systems Algorithms Utilized in Social Networks based e-Learning Systems & Neutrosophic System, *Neutrosophic Sets and Systems*, 8(2015), 32–41.
- [22] N. A. Nabeeh, M. Abdel-Basset and G. Soliman, A model for evaluating green credit rating and its impact on sustainability performance, *Journal of Cleaner Production*, 280(1)(2021), 124–299.
- [23] N. A. Nabeeh, F. Smarandache, M. Abdel-Basset, H. A. El-Ghareeb and . A. Aboelfetouh, An Integrated Neutrosophic-TOPSIS Approach and its Application to Personnel Selection: A New Trend in Brain Processing and Analysis, *IEEE Access*, 29734–29744, 2017.
- [24] N. A. Nabeeh, M. Abdel-Basset, H. A. El-Ghareeb and A. Aboelfetouh , *Neutrosophic Multi-Criteria Decision Making Approach for IoT-Based Enterprises*, *IEEE Access*, 2019.
- [25] N. Olgun and A. Hatip, The Effect Of The Neutrosophic Logic On The Decision Making, in *Quadruple Neutrosophic Theory And Applications*, Belgium, EU, Pons Editions Brussels, 2020, 238–253.
- [26] S. K. Patro and F. Smarandache, The neutrosophic statistical distribution, more problems, more solutions, *Neutrosophic Sets and Systems*, 12(2016), 73–79.
- [27] R. Sahin, Neutrosophic Hierarchical Clustering Algorithms, *Neutrosophic Sets and Systems*, 2(2014), 19–24.
- [28] M. Sahin, N. Olgun, V. Uluay, A. Kargn and F. Smarandache, A New Similarity Measure Based on Falsity Value between Single Valued Neutrosophic Sets Based on the Centroid Points of Transformed Single Valued Neutrosophic Numbers with Applications to Pattern Recognition, *Neutrosophic Sets and Systems*, 15(2017), 31–48.

- [29] A. Salama, A. Sharaf Al-Din, I. Abu Al-Qasim, R. Alhabib and M. Badran, Introduction to Decision Making for Neutrosophic Environment Study on the Suez Canal Port, *Neutrosophic Sets and Systems*, 35(2020), 22–44.
- [30] G. Shahzadi, M. Akram and A. B. Saeid, An Application of Single-Valued Neutrosophic Sets in Medical Diagnosis, *Neutrosophic Sets and Systems*, 18(2017), 80–88.
- [31] F. Smarandache, Neutrosophic Set a Generalization of the Intuitionistic Fuzzy Sets, *Inter. J. Pure Appl. Math.*, 2005, 287–297.
- [32] F. Smarandache, *Introduction to Neutrosophic Statistics*, USA: Sitech & Education Publishing, 2014.
- [33] F. Smarandache, *Introduction to Neutrosophic Measure, Neutrosophic Integral and Neutrosophic Probability*, Craiova, Romania: Sitech - Education, 2013.
- [34] F. Yuhua, Neutrosophic Examples in Physics, *Neutrosophic Sets and Systems*, 1(2013), 26–33.
- [35] M. B. Zeina, Neutrosophic Event-Based Queueing Model, *International Journal of Neutrosophic Science*, 6(1)(2020), 48–55.
- [36] M. B. Zeina, Erlang Service Queueing Model with Neutrosophic Parameters, *International Journal of Neutrosophic Science*, 6(2)(2020), 106–112.
- [37] M. B. Zeina and A. Hatip, Neutrosophic random variables. *Neutrosophic Sets and Systems*, 39(2021), 44–52.

Received: Aug 15, 2021 / Accepted: Dec 5, 2021