



Similarity Measures for Interval-Valued Neutrosophic Hypersoft Set With Their Application to Solve Decision Making Problem

Rana Muhammad Zulqarnain¹, Aiyared Iampan², Hamiden Abd El-Wahed Khalifa^{3,4},
Imran Siddique^{5*}

¹ College of Mathematics and Computer Science, Zhejiang Normal University, Jinhua, Zhejiang 321004, China. E-mail: ranazulqarnain7777@gmail.com

² Department of Mathematics, School of Science, University of Phayao, Mae Ka, Mueang, Phayao 56000, Thailand. E-mail: aiyared.ia@up.ac.th

³ Department of Mathematics, College of Science and Arts, Al-Badaya, Qassim University E-mail: Ha.Ahmed@qu.edu.sa

⁴ Operations Research Department, Faculty of Graduate Studies for Statistical Research, Cairo University, Giza, Egypt, E-mail: hamiden@cu.edu.eg

⁵ Department of Mathematics, University of Management and Technology Lahore, Pakistan. E-mail: imransiddique@umt.edu.pk

Correspondence: E-mail: imransiddique@umt.edu.pk

Abstract:

A similarity measure is used to tackle many issues that include indistinct and blurred information, excluding is not able to deal with the general fuzziness and obscurity of the problems that have various information. In this paper, we study some basic concepts which are helpful for us to build the structure of the article, such as soft set, neutrosophic soft set, hypersoft set, neutrosophic hypersoft set (NHSS), and interval-valued NHSS, etc. The main objective of the present research is to develop a cosine similarity measure and set-theoretic similarity measure for an IVNHSS with their necessary properties. A decision-making approach has been established by using cosine and set-theoretic similarity measures. Furthermore, we used to develop a technique to solve multi-criteria decision-making problems. Finally, the advantages, effectiveness, flexibility, and comparative analysis of the algorithms are given with prevailing methods.

Keywords: Neutrosophic soft-set; hypersoft set; neutrosophic hypersoft set; interval-valued neutrosophic hypersoft set; similarity measures

1. Introduction

Decision-making is an interesting concern to pick the perfect alternate for any specific persistence. Firstly, it is pretended that details about alternatives are accumulated in crisp numbers, but in real-life situations, collective farm information always conquers wrong and inaccurate information. Fuzzy sets are like sets having an element of membership (Mem) degree. In classical set theory, the Mem degree of the elements in a set is examined in binary form to see that the element is not entirely concomitant. In contrast, the fuzzy set theory enables advanced Mem categorization of the components in the set. The Mem function portrays it, and also the multipurpose unit interval of the Mem function is $[0, 1]$. In some circumstances, decision-makers consider objects' Mem and nonmembership (Nmem) values. Zadeh's FS cannot handle imprecise and vague information in such cases. Atanassov [2] developed the notion of intuitionistic fuzzy sets (IFS) to deal above mentioned difficulties. The IFS accommodates the imprecise and inaccurate information using Mem and Nmem values.

Atanassov IFS was unable to solve those problems in which decision-makers considered the membership degree (MD) and nonmembership degrees (NMD) such as $MD = 0.5$ and $NMD = 0.8$, then $0.5 + 0.8 \not\leq 1$. Yager [3, 4] extended the notion of IFS to Pythagorean fuzzy sets (PFSs) to

overcome above-discussed complications by modifying $MD + NMD \leq 1$ to $MD^2 + NMD^2 \leq 1$. After developing PFSs, Zhang and Xu [5] proposed operational laws for PFSs and established a DM approach to resolving the MCDM problem. Wei and Lu [6] planned some power aggregation operators (AOs) and proposed a DM technique to solve multi-attribute decision-making (MADM) issues under the Pythagorean fuzzy environment. Wang and Li [7] presented power Bonferroni mean operators for PFSs with their basic properties using interaction. Gao et al. [8] presented several aggregation operators by considering the interaction and proposed a DM approach to solving MADM difficulties utilizing the developed operators. Wei [9] developed the interaction operational laws for Pythagorean fuzzy numbers (PFNs) by considering interaction and established interaction aggregation operators by using the developed interaction operations. Zhang [10] developed the accuracy function and presented a DM approach to solving multiple criteria group decision-making (MCGDM) problems using PFNs. Wang et al. [11] extended the PFS and introduced an interactive Hamacher operation with some novel AOs. They also established a DM method to solve MADM problems using their proposed operators. Wang and Li [12] developed some interval-valued PFSs and utilized their operators to resolve multi-attribute group decision-making (MAGDM) issues. Peng and Yuan [13] established novel operators such as Pythagorean fuzzy point operators and developed a DM technique using their proposed operators. Peng and Yang [14] introduced fundamental operations and their necessary possessions under PFSs and planned DM methodology. Garg [15] developed the logarithmic operational laws for PFSs and proposed some AOs. Arora and Garg [16] presented the operational laws for linguistic IFS and developed prioritized AOs. Ma and Xu [17] presented some innovative AOs for PFSs and proposed the score and accuracy functions for PFNs.

Above mentioned theories and their DM methodologies have been used in several fields of life. But, these theories cannot deal with the parametrization of the alternatives. Molodtsov [18] developed soft sets (SS) to overcome the complications, as mentioned earlier. Molodtsov's SS competently deals with imprecise, vague, and unclear objects considering their parametrization. Maji et al. [19] prolonged the notion of SS and introduced some necessary operators with their properties. Maji et al. [20] established a DM technique using their developed operations for SS. They also merged two well-known theories, such as FS and SS, and established the concept of fuzzy soft sets (FSS) [21]. They also proposed an intuitionistic fuzzy soft set (IFSS) [22] and discussed their basic operations. Garg and Arora [23] extended the notion of IFSS and presented a generalized form of IFSS with AOs. They also planned a DM technique to resolve undefined and inaccurate information under IFSS information. Garg and Arora [24] presented the correlation and weighted correlation coefficients for IFSS and developed the TOPSIS approach utilizing established correlation procedures. Zulqarnain et al. [25] introduced some AOs and correlation coefficients for interval-valued IFSS. They also extended the TOPSIS technique using their developed correlation measures to solve the MADM problem. Peng et al. [26] proposed the Pythagorean fuzzy soft sets (PFSSs) and presented fundamental operations of PFSSs with their desirable properties by merging PFS and SS. Athira et al. [27] proposed the entropy measure for PFSSs. They also presented some distance measures for PFSSs and utilized their developed distance measures to solve DM [28] issues. Zulqarnain et al. [29] introduced Einstein operational laws for Pythagorean fuzzy soft numbers (PFSNs) and developed Einstein AOs utilizing defined operational laws for PFSNs. They also planned a DM approach to solve MADM problems with the help of presented operators. Riaz et al. [30] prolonged the idea of PFSSs and developed the m polar PFSSs. They also established the TOPSIS method under the considered hybrid structure and proposed a DM methodology to solve the MCGDM problem. Riaz et al. [31] developed the similarity measures for PFSS with their fundamental properties. Zulqarnain et al. [32, 33] protracted the Einstein ordered AOs for PFSSs and utilized their developed approach to solving the MAGDM problem.

All the above studies only deal the inadequate information because of membership and non-membership values. However, these theories cannot handle the overall incompatible and imprecise data. To address such inconsistent and vague records, the idea of the neutrosophic set (NS) was developed by Smarandache [34]. Maji [35] offered the perception of a neutrosophic soft set (NSS) with necessary operations. The idea of the possibility of NSS was developed by Karaaslan [36] and

introduced a possibility of neutrosophic soft DM method based on And-product. Broumi [37] developed the generalized NSS with some operations and properties and used the projected concept for DM. Deli and Subas [38] developed the single-valued Neutrosophic numbers (SVNNs) to solve MCDM problems. They also established the cut sets for SVNNs. Wang et al. [39] proposed the correlation coefficient (CC) for SVNSSs. Ye [40] introduced the simplified NSs with operational laws and AOs. Also, he presented an MCDM technique utilizing his planned AOs. Masooma et al. [41] progressed a new concept by combining the multipolar fuzzy set and NS, known as the multipolar NS. They also established various characterization and operations with examples. Zulqarnain et al. [42] offered the generalized neutrosophic TOPSIS and used their presented technique for supplier selection.

All the above studies have some limitations. When any attribute from a set contains further sub-attributes, then the above-presented theories fail to solve such problems. To overcome the limitations above, Smarandache [43] protracted the idea of SS to hypersoft sets (HSS) by substituting the one-parameter function f to a multi-parameter (sub-attribute) function. Smarandache claimed that the established HSS competently deals with uncertain objects compared to SS. Several researchers explored the HSS and presented a lot of extensions for HSS [44, 45]. Zulqarnain et al. [46] presented the IFHSS, the generalized version of IFSS. They established the TOPSIS method utilizing the developed correlation coefficient. Zulqarnain et al. [47] proposed the Pythagorean fuzzy hypersoft sets with AOs and correlation coefficients. They also established the TOPSIS technique using their developed correlation coefficient and utilized the presented approach to select appropriate anti-virus face masks. Zulqarnain et al. [48, 49] presented some fundamental operations with their properties for interval-valued NHSS. Also, they proposed the CC and WCC for interval-valued NHSS and established a decision-making approach utilizing their developed CC. Several researchers extended the notion of HSS and introduced different operators for hybrid structures of fuzzy hypersoft sets [50-55]. Broumi et al. [56] extended the mathematical algebra of neutrosophic graphs to the interval-valued neutrosophic graph. Broumi et al. [57] discussed several operations for interval-valued neutrosophic graphs and discussed their desirable properties. Singh [58] introduced a fuzzy three-way context using the properties of the interval-valued neutrosophic set, and its graphical properties deal with partial ignorance. Zulqarnain et al. [59, 60] proposed some fundamental operations for generalized multipolar neutrosophic soft sets and multipolar interval-valued neutrosophic soft sets. However, all existing studies only deal with the scenario by using MD and NMD of sub-attributes of the considered attributes. If any decision-maker considers the MD = 0.7 and NMD = 0.6, then $0.7 + 0.6 \leq 1$ of any sub-attribute of the alternatives. We can observe that the theories mentioned above cannot handle it. Zulqarnain et al. [47] proposed the more generalized notion of PFHSS comparative to IFHSS. The PFHSS accommodates more uncertainty compared to IFHSS by updating the condition $MD + NMD \leq 1$ to $(\mathcal{J}_{\mathcal{F}(\bar{a})}(\delta))^2 + (\mathcal{J}_{\mathcal{F}(\bar{a})}(\delta))^2 \leq 1$. All existing hybrid structures of FS cannot handle the indeterminacy of sub-attributes of considered n-tuple attributes. On the other hand, developed aggregation operators can accommodate the sub-attributes of considered attributes using truthiness, indeterminacy, and falsity objects of sub-attributes with the following condition $0 \leq \sigma_{\mathcal{F}(\bar{a}_k)}(\delta) + \tau_{\mathcal{F}(\bar{a}_k)}(\delta) + \gamma_{\mathcal{F}(\bar{a}_k)}(\delta) \leq 3$. It may be seen that the best selection of the suggested approach is to resemble the verbalized own method, and that ensures the liableness along with the effectiveness of the recommended approach.

The following research is organized: In section 2, we recollected some basic definitions used in the subsequent sequel, such as NS, SS, NSS, HSS, NHSS, and IVNHSS. Section 3 proposes the similarity measures such as cosine and set-theoretic for NHSS with its properties. We established a decision-making technique to solve decision-making complications utilizing our developed similarity measures. In section 4, we use the proposed similarity measures for decision-making. A brief comparative analysis has been conducted between proposed techniques with existing methodologies in section 5. Finally, the conclusion and future directions are presented in section 6.

2. Preliminaries

In this section, we recollect some basic definitions which are helpful to build the structure of the following manuscript such as soft set, hypersoft set, and neutrosophic hypersoft set.

Definition 2.1 [4]

Let \mathcal{U} be the universal set and \mathcal{E} be the set of attributes concerning \mathcal{U} . Let $\mathcal{P}(\mathcal{U})$ be the power set of \mathcal{U} and $\mathcal{A} \subseteq \mathcal{E}$. A pair $(\mathcal{F}, \mathcal{A})$ is called a soft set over \mathcal{U} , and its mapping is given as

$$\mathcal{F}:\mathcal{A} \rightarrow \mathcal{P}(\mathcal{U})$$

It is also defined as:

$$(\mathcal{F}, \mathcal{A}) = \{\mathcal{F}(e) \in \mathcal{P}(\mathcal{U}): e \in \mathcal{E}, \mathcal{F}(e) = \emptyset \text{ if } e \notin \mathcal{A}\}$$

Definition 2.2 [38]

Let \mathcal{U} be a universe of discourse and $\mathcal{P}(\mathcal{U})$ be a power set of \mathcal{U} and $k = \{k_1, k_2, k_3, \dots, k_n\}, (n \geq 1)$ be a set of attributes and set K_i a set of corresponding sub-attributes of k_i respectively with $K_i \cap K_j = \emptyset$ for $n \geq 1$ for each $i, j \in \{1, 2, 3 \dots n\}$ and $i \neq j$. Assume $K_1 \times K_2 \times K_3 \times \dots \times K_n = \ddot{\mathcal{A}} = \{a_{1h} \times a_{2k} \times \dots \times a_{nl}\}$ be a collection of multi-attributes, where $1 \leq h \leq \alpha, 1 \leq k \leq \beta,$ and $1 \leq l \leq \gamma,$ and $\alpha, \beta,$ and $\gamma \in \mathbb{N}$. Then the pair $(\mathcal{F}, K_1 \times K_2 \times K_3 \times \dots \times K_n = \ddot{\mathcal{A}})$ is said to be HSS over \mathcal{U} , and its mapping is defined as

$$\mathcal{F}: K_1 \times K_2 \times K_3 \times \dots \times K_n = \ddot{\mathcal{A}} \rightarrow \mathcal{P}(\mathcal{U}).$$

It is also defined as

$$(\mathcal{F}, \ddot{\mathcal{A}}) = \{\ddot{\mathcal{A}}, \mathcal{F}_{\ddot{\mathcal{A}}}(\ddot{\mathcal{A}}): \ddot{\mathcal{A}} \in \ddot{\mathcal{A}}, \mathcal{F}_{\ddot{\mathcal{A}}}(\ddot{\mathcal{A}}) \in \mathcal{P}(\mathcal{U})\}$$

Definition 2.3 [38]

Let \mathcal{U} be a universe of discourse and $\mathcal{P}(\mathcal{U})$ be a power set of \mathcal{U} and $k = \{k_1, k_2, k_3, \dots, k_n\}, (n \geq 1)$ be a set of attributes and set K_i a set of corresponding sub-attributes of k_i respectively with $K_i \cap K_j = \emptyset$ for $n \geq 1$ for each $i, j \in \{1, 2, 3 \dots n\}$ and $i \neq j$. Assume $K_1 \times K_2 \times K_3 \times \dots \times K_n = \ddot{\mathcal{A}} = \{a_{1h} \times a_{2k} \times \dots \times a_{nl}\}$ be a collection of sub-attributes, where $1 \leq h \leq \alpha, 1 \leq k \leq \beta,$ and $1 \leq l \leq \gamma,$ and $\alpha, \beta,$ and $\gamma \in \mathbb{N}$ and $NS^{\mathcal{U}}$ be a collection of all neutrosophic subsets over \mathcal{U} . Then the pair $(\mathcal{F}, K_1 \times K_2 \times K_3 \times \dots \times K_n = \ddot{\mathcal{A}})$ is said to be NHSS over \mathcal{U} , and its mapping is defined as

$$\mathcal{F}: K_1 \times K_2 \times K_3 \times \dots \times K_n = \ddot{\mathcal{A}} \rightarrow NS^{\mathcal{U}}.$$

It is also defined as

$$(\mathcal{F}, \ddot{\mathcal{A}}) = \{(\ddot{\mathcal{A}}, \mathcal{F}_{\ddot{\mathcal{A}}}(\ddot{\mathcal{A}})): \ddot{\mathcal{A}} \in \ddot{\mathcal{A}}, \mathcal{F}_{\ddot{\mathcal{A}}}(\ddot{\mathcal{A}}) \in NS^{\mathcal{U}}\}, \text{ where } \mathcal{F}_{\ddot{\mathcal{A}}}(\ddot{\mathcal{A}}) = \{(\delta, \sigma_{\mathcal{F}(\ddot{\mathcal{A}})}(\delta), \tau_{\mathcal{F}(\ddot{\mathcal{A}})}(\delta), \gamma_{\mathcal{F}(\ddot{\mathcal{A}})}(\delta)): \delta \in \mathcal{U}\},$$

where $\sigma_{\mathcal{F}(\ddot{\mathcal{A}})}(\delta), \tau_{\mathcal{F}(\ddot{\mathcal{A}})}(\delta),$ and $\gamma_{\mathcal{F}(\ddot{\mathcal{A}})}(\delta)$ represent the truth, indeterminacy, and falsity grades of the attributes such as $\sigma_{\mathcal{F}(\ddot{\mathcal{A}})}(\delta), \tau_{\mathcal{F}(\ddot{\mathcal{A}})}(\delta), \gamma_{\mathcal{F}(\ddot{\mathcal{A}})}(\delta) \in [0, 1],$ and $0 \leq \sigma_{\mathcal{F}(\ddot{\mathcal{A}})}(\delta) + \tau_{\mathcal{F}(\ddot{\mathcal{A}})}(\delta) + \gamma_{\mathcal{F}(\ddot{\mathcal{A}})}(\delta) \leq 3.$

Example 2.4

Consider the universe of discourse $\mathcal{U} = \{\delta_1, \delta_2\}$ and $\mathcal{Q} = \{\ell_1 = \text{Teaching methodology}, \ell_2 = \text{Subjects}, \ell_3 = \text{Classes}\}$ be a collection of attributes with following their corresponding attribute values are given as teaching methodology = $L_1 = \{a_{11} = \text{project base}, a_{12} = \text{class discussion}\},$ Subjects = $L_2 = \{a_{21} = \text{Mathematics}, a_{22} = \text{Computer Science}, a_{23} = \text{Statistics}\},$ and Classes = $L_3 = \{a_{31} = \text{Masters}, a_{32} = \text{Doctorol}\}.$ Let $\ddot{\mathcal{A}} = L_1 \times L_2 \times L_3$ be a set of attributes

$$\begin{aligned} \ddot{\mathcal{A}} &= L_1 \times L_2 \times L_3 = \{a_{11}, a_{12}\} \times \{a_{21}, a_{22}, a_{23}\} \times \{a_{31}, a_{32}\} \\ &= \{(a_{11}, a_{21}, a_{31}), (a_{11}, a_{21}, a_{32}), (a_{11}, a_{22}, a_{31}), (a_{11}, a_{22}, a_{32}), (a_{11}, a_{23}, a_{31}), (a_{11}, a_{23}, a_{32}), \\ &\quad (a_{12}, a_{21}, a_{31}), (a_{12}, a_{21}, a_{32}), (a_{12}, a_{22}, a_{31}), (a_{12}, a_{22}, a_{32}), (a_{12}, a_{23}, a_{31}), (a_{12}, a_{23}, a_{32}),\} \\ \ddot{\mathcal{A}} &= \{\check{\alpha}_1, \check{\alpha}_2, \check{\alpha}_3, \check{\alpha}_4, \check{\alpha}_5, \check{\alpha}_6, \check{\alpha}_7, \check{\alpha}_8, \check{\alpha}_9, \check{\alpha}_{10}, \check{\alpha}_{11}, \check{\alpha}_{12}\} \end{aligned}$$

Then the NHSS over \mathcal{U} is given as follows

$$(\mathcal{F}, \ddot{\mathbb{A}}) = \left\{ \begin{array}{l} (\check{\alpha}_1, (\delta_1, (.6, .3, .8)), (\delta_2, (.9, .3, .5))), (\check{\alpha}_2, (\delta_1, (.5, .2, .7)), (\delta_2, (.7, .1, .5))), (\check{\alpha}_3, (\delta_1, (.5, .2, .8)), (\delta_2, (.4, .3, .4))), \\ (\check{\alpha}_4, (\delta_1, (.2, .5, .6)), (\delta_2, (.5, .1, .6))), (\check{\alpha}_5, (\delta_1, (.8, .4, .3)), (\delta_2, (.2, .3, .5))), (\check{\alpha}_6, (\delta_1, (.9, .6, .4)), (\delta_2, (.7, .6, .8))), \\ (\check{\alpha}_7, (\delta_1, (.6, .5, .3)), (\delta_2, (.4, .2, .8))), (\check{\alpha}_8, (\delta_1, (.8, .2, .5)), (\delta_2, (.6, .8, .4))), (\check{\alpha}_9, (\delta_1, (.7, .4, .9)), (\delta_2, (.7, .3, .5))), \\ (\check{\alpha}_{10}, (\delta_1, (.8, .4, .6)), (\delta_2, (.7, .2, .9))), (\check{\alpha}_{11}, (\delta_1, (.8, .4, .5)), (\delta_2, (.4, .2, .5))), (\check{\alpha}_5, (\delta_1, (.7, .5, .8)), (\delta_2, (.7, .5, .9))) \end{array} \right\}$$

Definition 2.5 [42]

Let \mathcal{U} be a universe of discourse and $\mathcal{P}(\mathcal{U})$ be a power set of \mathcal{U} and $k = \{k_1, k_2, k_3, \dots, k_n\}, (n \geq 1)$ be a set of attributes and set K_i a set of corresponding sub-attributes of k_i respectively with $K_i \cap K_j = \varphi$ for $n \geq 1$ for each $i, j \in \{1, 2, 3 \dots n\}$ and $i \neq j$. Assume $K_1 \times K_2 \times K_3 \times \dots \times K_n = \ddot{\mathbb{A}} = \{a_{1h} \times a_{2k} \times \dots \times a_{nl}\}$ be a collection of sub-attributes, where $1 \leq h \leq \alpha, 1 \leq k \leq \beta,$ and $1 \leq l \leq \gamma,$ and $\alpha, \beta,$ and $\gamma \in \mathbb{N}$ and $IVNS^{\mathcal{U}}$ be a collection of all interval-valued neutrosophic subsets over \mathcal{U} . Then the pair $(\mathcal{F}, K_1 \times K_2 \times K_3 \times \dots \times K_n = \ddot{\mathbb{A}})$ is said to be IVNHSS over \mathcal{U} , and its mapping is defined as

$$\mathcal{F}: K_1 \times K_2 \times K_3 \times \dots \times K_n = \ddot{\mathbb{A}} \rightarrow IVNS^{\mathcal{U}}.$$

It is also defined as

$$(\mathcal{F}, \ddot{\mathbb{A}}) = \{(\check{\alpha}_k, \mathcal{F}_{\ddot{\mathbb{A}}}(\check{\alpha}_k)): \check{\alpha}_k \in \ddot{\mathbb{A}}, \mathcal{F}_{\ddot{\mathbb{A}}}(\check{\alpha}_k) \in NS^{\mathcal{U}}\}, \text{ where } \mathcal{F}_{\ddot{\mathbb{A}}}(\check{\alpha}) = \{ \langle \delta, \sigma_{\mathcal{F}(\check{\alpha}_k)}(\delta), \tau_{\mathcal{F}(\check{\alpha}_k)}(\delta), \gamma_{\mathcal{F}(\check{\alpha}_k)}(\delta) \rangle : \delta \in \mathcal{U} \},$$

where $\sigma_{\mathcal{F}(\check{\alpha}_k)}(\delta), \tau_{\mathcal{F}(\check{\alpha}_k)}(\delta),$ and $\gamma_{\mathcal{F}(\check{\alpha}_k)}(\delta)$ represent the interval truth, indeterminacy, and falsity grades of the attributes such as $\sigma_{\mathcal{F}(\check{\alpha}_k)}(\delta) = [\sigma_{\mathcal{F}(\check{\alpha}_k)}^{\ell}(\delta), \sigma_{\mathcal{F}(\check{\alpha}_k)}^{\bar{\cup}}(\delta)], \tau_{\mathcal{F}(\check{\alpha}_k)}(\delta) = [\tau_{\mathcal{F}(\check{\alpha}_k)}^{\ell}(\delta), \tau_{\mathcal{F}(\check{\alpha}_k)}^{\bar{\cup}}(\delta)], \gamma_{\mathcal{F}(\check{\alpha}_k)}(\delta) = [\gamma_{\mathcal{F}(\check{\alpha}_k)}^{\ell}(\delta), \gamma_{\mathcal{F}(\check{\alpha}_k)}^{\bar{\cup}}(\delta)],$

where $\sigma_{\mathcal{F}(\check{\alpha}_k)}^{\ell}(\delta), \sigma_{\mathcal{F}(\check{\alpha}_k)}^{\bar{\cup}}(\delta), \tau_{\mathcal{F}(\check{\alpha}_k)}^{\ell}(\delta), \tau_{\mathcal{F}(\check{\alpha}_k)}^{\bar{\cup}}(\delta), \gamma_{\mathcal{F}(\check{\alpha}_k)}^{\ell}(\delta), \gamma_{\mathcal{F}(\check{\alpha}_k)}^{\bar{\cup}}(\delta) \subseteq [0, 1],$ and $0 \leq$

$$\sigma_{\mathcal{F}(\check{\alpha}_k)}^{\bar{\cup}}(\delta) + \tau_{\mathcal{F}(\check{\alpha}_k)}^{\bar{\cup}}(\delta) + \gamma_{\mathcal{F}(\check{\alpha}_k)}^{\bar{\cup}}(\delta) \leq 3.$$

Simply an interval-valued neutrosophic hypersoft number (IVNHSSN) can be expressed as $\mathcal{F} = \{ \langle \sigma_{\mathcal{F}(\check{\alpha}_k)}^{\ell}(\delta), \sigma_{\mathcal{F}(\check{\alpha}_k)}^{\bar{\cup}}(\delta), [\tau_{\mathcal{F}(\check{\alpha}_k)}^{\ell}(\delta), \tau_{\mathcal{F}(\check{\alpha}_k)}^{\bar{\cup}}(\delta)], [\gamma_{\mathcal{F}(\check{\alpha}_k)}^{\ell}(\delta), \gamma_{\mathcal{F}(\check{\alpha}_k)}^{\bar{\cup}}(\delta)] \rangle, \text{ where } 0 \leq \sigma_{\mathcal{F}(\check{\alpha}_k)}^{\bar{\cup}}(\delta) + \tau_{\mathcal{F}(\check{\alpha}_k)}^{\bar{\cup}}(\delta) + \gamma_{\mathcal{F}(\check{\alpha}_k)}^{\bar{\cup}}(\delta) \leq 3.$

3. Similarity Measures for Interval-Valued Neutrosophic Hypersoft Set

Many mathematicians developed various methodologies to solve MCDM problems in the past few years, such as aggregation operators for different hybrid structures, CC, similarity measures, and decision-making applications. The following section develops the cosine and set-theoretic similarity measure for IVNHSS.

Definition 3.1

Let $(\mathcal{F}, \ddot{\mathbb{A}}) = \{ \langle \delta_i, [\sigma_{\mathcal{F}(\check{\alpha}_k)}^{\ell}(\delta_i), \sigma_{\mathcal{F}(\check{\alpha}_k)}^{\bar{\cup}}(\delta_i)], [\tau_{\mathcal{F}(\check{\alpha}_k)}^{\ell}(\delta_i), \tau_{\mathcal{F}(\check{\alpha}_k)}^{\bar{\cup}}(\delta_i)], [\gamma_{\mathcal{F}(\check{\alpha}_k)}^{\ell}(\delta_i), \gamma_{\mathcal{F}(\check{\alpha}_k)}^{\bar{\cup}}(\delta_i)] \rangle \mid \delta_i \in \mathcal{U} \}$ and $(\mathcal{G}, \ddot{\mathbb{M}}) = \{ \langle \delta_i, [\sigma_{\mathcal{G}(\check{\alpha}_k)}^{\ell}(\delta_i), \sigma_{\mathcal{G}(\check{\alpha}_k)}^{\bar{\cup}}(\delta_i)], [\tau_{\mathcal{G}(\check{\alpha}_k)}^{\ell}(\delta_i), \tau_{\mathcal{G}(\check{\alpha}_k)}^{\bar{\cup}}(\delta_i)], [\gamma_{\mathcal{G}(\check{\alpha}_k)}^{\ell}(\delta_i), \gamma_{\mathcal{G}(\check{\alpha}_k)}^{\bar{\cup}}(\delta_i)] \rangle \mid \delta_i \in \mathcal{U} \}$ be two IVNHSSs defined over a universe of discourse \mathcal{U} . Then, the then cosine similarity measure of $(\mathcal{F}, \ddot{\mathbb{A}})$ and $(\mathcal{G}, \ddot{\mathbb{M}})$ can be described as follows:

$$S_{IVNHSS}^1((\mathcal{F}, \check{\mathbb{A}}), (\mathcal{G}, \check{\mathbb{M}})) = \frac{1}{mn} \sum_{k=1}^m \sum_{i=1}^n \frac{(\sigma_{\mathcal{F}(\check{a}_k)}^{\ell}(\delta_i) + \sigma_{\mathcal{F}(\check{a}_k)}^{\check{\lambda}}(\delta_i))(\sigma_{\mathcal{G}(\check{a}_k)}^{\ell}(\delta_i) + \sigma_{\mathcal{G}(\check{a}_k)}^{\check{\lambda}}(\delta_i)) + (\tau_{\mathcal{F}(\check{a}_k)}^{\ell}(\delta_i) + \tau_{\mathcal{F}(\check{a}_k)}^{\check{\lambda}}(\delta_i))(\tau_{\mathcal{G}(\check{a}_k)}^{\ell}(\delta_i) + \tau_{\mathcal{G}(\check{a}_k)}^{\check{\lambda}}(\delta_i)) + (\gamma_{\mathcal{F}(\check{a}_k)}^{\ell}(\delta_i) + \gamma_{\mathcal{F}(\check{a}_k)}^{\check{\lambda}}(\delta_i))(\gamma_{\mathcal{G}(\check{a}_k)}^{\ell}(\delta_i) + \gamma_{\mathcal{G}(\check{a}_k)}^{\check{\lambda}}(\delta_i))}{\left(\sqrt{\left((\sigma_{\mathcal{F}(\check{a}_k)}^{\ell}(\delta_i))^2 + (\sigma_{\mathcal{F}(\check{a}_k)}^{\check{\lambda}}(\delta_i))^2 + (\tau_{\mathcal{F}(\check{a}_k)}^{\ell}(\delta_i))^2 + (\tau_{\mathcal{F}(\check{a}_k)}^{\check{\lambda}}(\delta_i))^2 + (\gamma_{\mathcal{F}(\check{a}_k)}^{\ell}(\delta_i))^2 + (\gamma_{\mathcal{F}(\check{a}_k)}^{\check{\lambda}}(\delta_i))^2 \right)} \right)} \right)$$

Proposition 3.2

Let $(\mathcal{F}, \check{\mathbb{A}})$, $(\mathcal{G}, \check{\mathbb{M}})$, and $(\mathcal{H}, \check{\mathbb{C}}) \in \text{NHSS}$, then the following properties hold

1. $0 \leq S_{IVNHSS}^1((\mathcal{F}, \check{\mathbb{A}}), (\mathcal{G}, \check{\mathbb{M}})) \leq 1$
2. $S_{IVNHSS}^1((\mathcal{F}, \check{\mathbb{A}}), (\mathcal{G}, \check{\mathbb{M}})) = S_{IVNHSS}^1((\mathcal{G}, \check{\mathbb{M}}), (\mathcal{F}, \check{\mathbb{A}}))$
3. If $(\mathcal{F}, \check{\mathbb{A}}) \subseteq (\mathcal{G}, \check{\mathbb{M}}) \subseteq (\mathcal{H}, \check{\mathbb{C}})$, then $S_{IVNHSS}^1((\mathcal{F}, \check{\mathbb{A}}), (\mathcal{H}, \check{\mathbb{C}})) \leq S_{IVNHSS}^1((\mathcal{F}, \check{\mathbb{A}}), (\mathcal{G}, \check{\mathbb{M}}))$ and $S_{IVNHSS}^1((\mathcal{F}, \check{\mathbb{A}}), (\mathcal{H}, \check{\mathbb{C}})) \leq S_{IVNHSS}^1((\mathcal{G}, \check{\mathbb{M}}), (\mathcal{H}, \check{\mathbb{C}}))$.

Proof: Using the above definition, the proof of these properties can be done easily.

Definition 3.3

Let $(\mathcal{F}, \check{\mathbb{A}}) = \{ \{ \delta_i, [\sigma_{\mathcal{F}(\check{a}_k)}^{\ell}(\delta_i), \sigma_{\mathcal{F}(\check{a}_k)}^{\check{\lambda}}(\delta_i)], [\tau_{\mathcal{F}(\check{a}_k)}^{\ell}(\delta_i), \tau_{\mathcal{F}(\check{a}_k)}^{\check{\lambda}}(\delta_i)], [\gamma_{\mathcal{F}(\check{a}_k)}^{\ell}(\delta_i), \gamma_{\mathcal{F}(\check{a}_k)}^{\check{\lambda}}(\delta_i)] \} \mid \delta_i \in \mathcal{U} \}$ and $(\mathcal{G}, \check{\mathbb{M}}) = \{ \{ \delta_i, [\sigma_{\mathcal{G}(\check{a}_k)}^{\ell}(\delta_i), \sigma_{\mathcal{G}(\check{a}_k)}^{\check{\lambda}}(\delta_i)], [\tau_{\mathcal{G}(\check{a}_k)}^{\ell}(\delta_i), \tau_{\mathcal{G}(\check{a}_k)}^{\check{\lambda}}(\delta_i)], [\gamma_{\mathcal{G}(\check{a}_k)}^{\ell}(\delta_i), \gamma_{\mathcal{G}(\check{a}_k)}^{\check{\lambda}}(\delta_i)] \} \mid \delta_i \in \mathcal{U} \}$ be two IVNHSSs defined over a universe of discourse \mathcal{U} . Then, the then cosine similarity measure of $(\mathcal{F}, \check{\mathbb{A}})$ and $(\mathcal{G}, \check{\mathbb{M}})$ can be described as follows:

$$S_{IVNHSS}^1((\mathcal{F}, \check{\mathbb{A}}), (\mathcal{G}, \check{\mathbb{M}})) = \frac{1}{mn} \sum_{k=1}^m \sum_{i=1}^n \frac{(\sigma_{\mathcal{F}(\check{a}_k)}^{\ell}(\delta_i) + \sigma_{\mathcal{F}(\check{a}_k)}^{\check{\lambda}}(\delta_i))(\sigma_{\mathcal{G}(\check{a}_k)}^{\ell}(\delta_i) + \sigma_{\mathcal{G}(\check{a}_k)}^{\check{\lambda}}(\delta_i)) + (\tau_{\mathcal{F}(\check{a}_k)}^{\ell}(\delta_i) + \tau_{\mathcal{F}(\check{a}_k)}^{\check{\lambda}}(\delta_i))(\tau_{\mathcal{G}(\check{a}_k)}^{\ell}(\delta_i) + \tau_{\mathcal{G}(\check{a}_k)}^{\check{\lambda}}(\delta_i)) + (\gamma_{\mathcal{F}(\check{a}_k)}^{\ell}(\delta_i) + \gamma_{\mathcal{F}(\check{a}_k)}^{\check{\lambda}}(\delta_i))(\gamma_{\mathcal{G}(\check{a}_k)}^{\ell}(\delta_i) + \gamma_{\mathcal{G}(\check{a}_k)}^{\check{\lambda}}(\delta_i))}{\max \left(\left(\left((\sigma_{\mathcal{F}(\check{a}_k)}^{\ell}(\delta_i))^2 + (\sigma_{\mathcal{F}(\check{a}_k)}^{\check{\lambda}}(\delta_i))^2 + (\tau_{\mathcal{F}(\check{a}_k)}^{\ell}(\delta_i))^2 + (\tau_{\mathcal{F}(\check{a}_k)}^{\check{\lambda}}(\delta_i))^2 + (\gamma_{\mathcal{F}(\check{a}_k)}^{\ell}(\delta_i))^2 + (\gamma_{\mathcal{F}(\check{a}_k)}^{\check{\lambda}}(\delta_i))^2 \right) \right), \left(\left((\sigma_{\mathcal{G}(\check{a}_k)}^{\ell}(\delta_i))^2 + (\sigma_{\mathcal{G}(\check{a}_k)}^{\check{\lambda}}(\delta_i))^2 + (\tau_{\mathcal{G}(\check{a}_k)}^{\ell}(\delta_i))^2 + (\tau_{\mathcal{G}(\check{a}_k)}^{\check{\lambda}}(\delta_i))^2 + (\gamma_{\mathcal{G}(\check{a}_k)}^{\ell}(\delta_i))^2 + (\gamma_{\mathcal{G}(\check{a}_k)}^{\check{\lambda}}(\delta_i))^2 \right) \right) \right)}$$

Proposition 3.4

Let $(\mathcal{F}, \check{\mathbb{A}}) = \{ \{ \delta_i, [\sigma_{\mathcal{F}(\check{a}_k)}^{\ell}(\delta_i), \sigma_{\mathcal{F}(\check{a}_k)}^{\check{\lambda}}(\delta_i)], [\tau_{\mathcal{F}(\check{a}_k)}^{\ell}(\delta_i), \tau_{\mathcal{F}(\check{a}_k)}^{\check{\lambda}}(\delta_i)], [\gamma_{\mathcal{F}(\check{a}_k)}^{\ell}(\delta_i), \gamma_{\mathcal{F}(\check{a}_k)}^{\check{\lambda}}(\delta_i)] \} \mid \delta_i \in \mathcal{U} \}$ and $(\mathcal{G}, \check{\mathbb{M}}) = \{ \{ \delta_i, [\sigma_{\mathcal{G}(\check{a}_k)}^{\ell}(\delta_i), \sigma_{\mathcal{G}(\check{a}_k)}^{\check{\lambda}}(\delta_i)], [\tau_{\mathcal{G}(\check{a}_k)}^{\ell}(\delta_i), \tau_{\mathcal{G}(\check{a}_k)}^{\check{\lambda}}(\delta_i)], [\gamma_{\mathcal{G}(\check{a}_k)}^{\ell}(\delta_i), \gamma_{\mathcal{G}(\check{a}_k)}^{\check{\lambda}}(\delta_i)] \} \mid \delta_i \in \mathcal{U} \}$ be two IVNHSSs. Then, the following properties hold.

1. $0 \leq S_{IVNHSS}^2((\mathcal{F}, \check{\mathbb{A}}), (\mathcal{G}, \check{\mathbb{M}})) \leq 1$
2. $S_{IVNHSS}^2((\mathcal{F}, \check{\mathbb{A}}), (\mathcal{G}, \check{\mathbb{M}})) = S_{IVNHSS}^2((\mathcal{G}, \check{\mathbb{M}}), (\mathcal{F}, \check{\mathbb{A}}))$
3. If $(\mathcal{F}, \check{\mathbb{A}}) \subseteq (\mathcal{G}, \check{\mathbb{M}}) \subseteq (\mathcal{H}, \check{\mathbb{C}})$, then $S_{IVNHSS}^2((\mathcal{F}, \check{\mathbb{A}}), (\mathcal{H}, \check{\mathbb{C}})) \leq S_{IVNHSS}^2((\mathcal{F}, \check{\mathbb{A}}), (\mathcal{G}, \check{\mathbb{M}}))$ and $S_{IVNHSS}^2((\mathcal{F}, \check{\mathbb{A}}), (\mathcal{H}, \check{\mathbb{C}})) \leq S_{IVNHSS}^2((\mathcal{G}, \check{\mathbb{M}}), (\mathcal{H}, \check{\mathbb{C}}))$.

Proof: Using the above definition, the proof of these properties can be done easily.

4. Application of Correlation Coefficient for Decision Making Under IVNHSS Environment

This section utilized the developed approaches based on cosine and set-theoretic similarity measures for decision making.

4.1 Algorithm for Similarity Measures of IVNHSS

- Step 1. Pick out the set containing parameters.
- Step 2. Construct the IVNHSS according to experts.

- Step 3. Compute the cosine similarity measure by using definition 3.1.
- Step 4. Compute the set-theoretic similarity measure for NHSS by utilizing definition 3.3.
- Step 5. An alternative with a maximum value with cosine similarity measure has the maximum rank according to considered numerical illustration.
- Step 6. An alternative with a maximum value with a set-theoretic similarity measure has the maximum rank according to considered numerical illustration.
- Step 7. Analyze the ranking.”




Figure 1: Flow chart of the presented similarity measures

4.2. Problem Formulation and Application of IVNHSS For Decision Making

A construction company calls for the appointment of a civil engineer to supervise the workers. Several engineers apply for the civil engineer post, simply four engineers call for an interview based on experience for undervaluation such as $S = \{S_1, S_2, S_3, S_4\}$ be a set of selected engineers call for the interview. The managing director hires a committee of four experts $U = \{\kappa_1, \kappa_2, \kappa_3, \kappa_4\}$ for the selection of civil engineer. The group of experts chooses the set of attributes for the selection of an appropriate civil engineer such as $\mathcal{Q} = \{\ell_1 = Experience, \ell_2 = Dealing\ skills, \ell_3 = Qualification\}$ be a set of parameters for the selection of MS. Experience = $\ell_1 = \{a_{11} = more\ than\ 20, a_{12} = less\ than\ 20\}$, Dealing skills = $\ell_2 = \{a_{21} = public\ dealing, a_{22} = Staff\ dealing\}$, and Qualification = $\ell_3 = \{a_{31} = Doctoral\ degree\ in\ medical\ education, a_{32} = Masters\ degree\ in\ medical\ education\}$. Let $\mathcal{Q}' = \ell_1 \times \ell_2 \times \ell_3$ be a set of sub-attributes. The experts evaluate the applicants under defined parameters and forward the evaluation performa to the company's managing director. Finally, the director scrutinizes the best applicant based on the expert's evaluation report. Let $\mathcal{Q}' = \ell_1 \times \ell_2 \times \ell_3$ be a set of sub-attributes

$$\mathcal{Q}' = \ell_1 \times \ell_2 \times \ell_3 = \{a_{11}, a_{12}\} \times \{a_{21}, a_{22}\} \times \{a_{31}, a_{32}\}$$

$$= \{(a_{11}, a_{21}, a_{31}), (a_{11}, a_{21}, a_{32}), (a_{11}, a_{22}, a_{31}), (a_{11}, a_{22}, a_{32}), (a_{12}, a_{21}, a_{31}), (a_{12}, a_{21}, a_{32}), (a_{12}, a_{22}, a_{31}), (a_{12}, a_{22}, a_{32})\}, \mathcal{Q}' = \{\check{a}_1, \check{a}_2, \check{a}_3, \check{a}_4, \check{a}_5, \check{a}_6, \check{a}_7, \check{a}_8\}$$

be a set of all multi sub-attributes. Each DM will evaluate the ratings of each alternative in the form of IVNHSNs under the considered multi sub-attributes. The developed method to find the best alternative is as follows.

4.2.1. Application of IVNHSS For Decision Making

Assume $S = \{S_1, S_2, S_3, S_4\}$ be a set of alternatives which are shortlisted for interview and $\mathfrak{L} = \{\ell_1 = \text{Experiance}, \ell_2 = \text{Dealing skills}, \ell_3 = \text{Qualification}\}$ be a set of parameters for the selection of MS. Experience = $\ell_1 = \{a_{11} = \text{more than 20}, a_{12} = \text{less than 20}\}$, Dealing skills = $\ell_2 = \{a_{21} = \text{public dealing}, a_{22} = \text{Staff dealing}\}$, and Qualification = $\ell_3 = \{a_{31} = \text{Doctoral degree in medical education}, a_{32} = \text{Masters degree in medical education}\}$. Let $\mathfrak{L}' = \ell_1 \times \ell_2 \times \ell_3$ be a set of sub-attributes. The health ministry defines a criterion for selecting MS for all alternatives in terms of IVNHSNs given in Table 1.

Table 1. Decision Matrix of Concerning Department

ρ	$\check{\alpha}_1$	$\check{\alpha}_2$	$\check{\alpha}_3$	$\check{\alpha}_4$	$\check{\alpha}_5$	$\check{\alpha}_6$	$\check{\alpha}_7$	$\check{\alpha}_8$
δ_1	([.3,.5],[.2,.4],[.2,.6])	([.2,.3],[.5,.7],[.1,.3])	([.5,.6],[.1,.3],[.4,.6])	([.2,.4],[.3,.5],[.3,.6])	([.2,.3],[.2,.4],[.4,.5])	([.4,.6],[.1,.3],[.2,.4])	([.6,.7],[.2,.3],[.3,.4])	([.4,.5],[.5,.8],[.1,.2])
δ_2	([.5,.6],[.1,.3],[.4,.6])	([.5,.7],[.1,.2],[.4,.6])	([.2,.4],[.3,.4],[.2,.5])	([.6,.8],[.1,.2],[.3,.5])	([.4,.6],[.4,.5],[.3,.5])	([.3,.5],[.4,.5],[.1,.3])	([.1,.2],[.5,.8],[.2,.4])	([.5,.7],[.1,.2],[.5,.6])
δ_3	([.2,.4],[.5,.6],[.4,.6])	([.2,.4],[.3,.4],[.2,.5])	([.4,.6],[.2,.3],[.1,.4])	([.2,.5],[.2,.3],[.1,.6])	([.3,.4],[.2,.5],[.5,.7])	([.3,.5],[.4,.5],[.1,.3])	([.2,.4],[.7,.8],[.1,.2])	([.1,.2],[.7,.8],[.2,.3])
δ_4	([.2,.3],[.5,.7],[.1,.3])	([.3,.4],[.2,.5],[.5,.7])	([.2,.4],[.3,.5],[.3,.6])	([.5,.7],[.1,.2],[.4,.6])	([.4,.6],[.1,.3],[.2,.4])	([.1,.2],[.5,.8],[.2,.4])	([.2,.4],[.3,.4],[.2,.5])	([.5,.6],[.1,.3],[.4,.6])

Table 2. Decision Matrix for Alternative $\aleph^{(1)}$

$\aleph^{(1)}$	$\check{\alpha}_1$	$\check{\alpha}_2$	$\check{\alpha}_3$	$\check{\alpha}_4$	$\check{\alpha}_5$	$\check{\alpha}_6$	$\check{\alpha}_7$	$\check{\alpha}_8$
δ_1	([.2,.4],[.4,.5],[.3,.4])	([.3,.4],[.4,.5],[.2,.5])	([.3,.6],[.2,.3],[.1,.2])	([.2,.4],[.4,.6],[.1,.2])	([.1,.3],[.6,.7],[.2,.3])	([.4,.5],[.2,.5],[.2,.3])	([.6,.7],[.1,.2],[.2,.3])	([.4,.6],[.2,.3],[.4,.5])
δ_2	([.3,.4],[.2,.5],[.5,.7])	([.4,.7],[.1,.2],[.1,.2])	([.4,.5],[.2,.5],[.1,.2])	([.5,.7],[.1,.2],[.2,.4])	([.6,.8],[.1,.2],[.1,.5])	([.2,.4],[.7,.8],[.1,.2])	([.2,.4],[.3,.5],[.3,.6])	([.3,.4],[.4,.5],[.2,.4])
δ_3	([.5,.6],[.2,.3],[.4,.5])	([.5,.7],[.1,.2],[.2,.4])	([.7,.8],[.1,.2],[.2,.4])	([.1,.3],[.1,.5],[.2,.5])	([.1,.4],[.2,.4],[.1,.2])	([.2,.5],[.2,.4],[.3,.5])	([.3,.5],[.2,.4],[.4,.6])	([.5,.7],[.1,.2],[.5,.6])
δ_4	([.3,.5],[.3,.4],[.6,.7])	([.2,.4],[.3,.4],[.2,.5])	([.2,.4],[.7,.8],[.1,.2])	([.4,.7],[.1,.2],[.1,.2])	([.5,.6],[.2,.3],[.4,.5])	([.2,.4],[.3,.5],[.3,.6])	([.4,.6],[.2,.3],[.4,.5])	([.1,.3],[.1,.5],[.2,.5])

Table 3. Decision Matrix for Alternative $\aleph^{(2)}$

$\aleph^{(2)}$	$\check{\alpha}_1$	$\check{\alpha}_2$	$\check{\alpha}_3$	$\check{\alpha}_4$	$\check{\alpha}_5$	$\check{\alpha}_6$	$\check{\alpha}_7$	$\check{\alpha}_8$
δ_1	([.2,.4],[.4,.6],[.4,.5])	([.2,.3],[.4,.6],[.3,.5])	([.1,.2],[.6,.8],[.2,.5])	([.4,.5],[.2,.5],[.1,.2])	([.2,.3],[.4,.6],[.3,.5])	([.1,.2],[.6,.8],[.2,.5])	([.7,.8],[.1,.2],[.2,.3])	([.1,.3],[.6,.7],[.2,.5])
δ_2	([.4,.5],[.2,.5],[.1,.2])	([.5,.7],[.1,.2],[.2,.4])	([.1,.3],[.6,.7],[.2,.6])	([.1,.4],[.2,.5],[.4,.6])	([.1,.4],[.2,.4],[.1,.2])	([.1,.2],[.2,.5],[.4,.6])	([.1,.4],[.2,.5],[.4,.6])	([.1,.4],[.2,.5],[.4,.6])
δ_3	([.3,.4],[.2,.6],[.4,.6])	([.2,.4],[.3,.4],[.2,.5])	([.4,.5],[.2,.5],[.1,.2])	([.1,.2],[.2,.5],[.4,.6])	([.3,.5],[.3,.5],[.6,.7])	([.3,.5],[.3,.5],[.6,.7])	([.1,.2],[.2,.5],[.4,.6])	([.5,.7],[.1,.2],[.2,.4])
δ_4	([.2,.4],[.4,.5],[.6,.8])	([.3,.5],[.3,.5],[.6,.7])	([.1,.2],[.2,.5],[.4,.6])	([.1,.4],[.2,.4],[.1,.2])	([.4,.5],[.2,.5],[.1,.2])	([.1,.2],[.2,.5],[.4,.6])	([.4,.5],[.2,.5],[.1,.2])	([.1,.2],[.2,.5],[.4,.6])

Table 4. Decision Matrix for Alternative $\aleph^{(3)}$

$\aleph^{(3)}$	$\check{\alpha}_1$	$\check{\alpha}_2$	$\check{\alpha}_3$	$\check{\alpha}_4$	$\check{\alpha}_5$	$\check{\alpha}_6$	$\check{\alpha}_7$	$\check{\alpha}_8$
δ_1	([.6,.7],[.1,.2],[.3,.5])	([.6,.8],[.1,.2],[.2,.3])	([.6,.7],[.3,.5],[.1,.2])	([.7,.8],[.1,.2],[.2,.5])	([.6,.7],[.1,.2],[.1,.2])	([.5,.8],[.1,.2],[.2,.4])	([.1,.3],[.6,.7],[.2,.5])	([.7,.8],[.1,.2],[.2,.3])
δ_2	([.5,.7],[.3,.4],[.2,.3])	([.5,.7],[.2,.5],[.2,.3])	([.5,.6],[.3,.4],[.1,.2])	([.7,.8],[.3,.5],[.1,.3])	([.1,.2],[.2,.5],[.4,.6])	([.1,.4],[.2,.5],[.4,.6])	([.4,.6],[.2,.3],[.1,.2])	([.4,.6],[.2,.3],[.1,.2])
δ_3	([.2,.4],[.3,.4],[.2,.5])	([.4,.7],[.2,.3],[.3,.7])	([.4,.6],[.2,.3],[.1,.2])	([.3,.5],[.3,.5],[.6,.7])	([.6,.8],[.1,.2],[.1,.2])	([.7,.8],[.1,.2],[.2,.4])	([.1,.2],[.2,.5],[.4,.6])	([.6,.8],[.1,.2],[.1,.3])
δ_4	([.6,.8],[.3,.4],[.1,.2])	([.5,.7],[.1,.2],[.4,.5])	([.1,.2],[.2,.5],[.4,.6])	([.5,.6],[.3,.4],[.1,.2])	([.2,.4],[.3,.4],[.2,.5])	([.1,.3],[.6,.7],[.2,.5])	([.7,.8],[.1,.2],[.2,.5])	([.4,.6],[.2,.3],[.1,.2])

Table 5. Decision Matrix for Alternative $\aleph^{(4)}$

$\aleph^{(4)}$	$\check{\alpha}_1$	$\check{\alpha}_2$	$\check{\alpha}_3$	$\check{\alpha}_4$	$\check{\alpha}_5$	$\check{\alpha}_6$	$\check{\alpha}_7$	$\check{\alpha}_8$
δ_1	([.3,.5],[.2,.4],[.1,.2])	([.3,.6],[.1,.2],[.4,.7])	([.4,.7],[.3,.4],[.2,.3])	([.7,.8],[.2,.4],[.3,.5])	([.5,.7],[.3,.4],[.2,.4])	([.4,.6],[.2,.5],[.3,.4])	([.2,.3],[.5,.7],[.2,.4])	([.5,.7],[.2,.4],[.3,.5])
δ_2	([.4,.5],[.5,.7],[.2,.4])	([.4,.7],[.3,.5],[.2,.4])	([.5,.8],[.3,.4],[.2,.3])	([.2,.4],[.2,.3],[.4,.5])	([.3,.5],[.2,.3],[.3,.5])	([.2,.4],[.2,.3],[.3,.6])	([.5,.8],[.3,.6],[.2,.3])	([.4,.6],[.2,.3],[.1,.2])
δ_3	([.2,.4],[.3,.4],[.2,.5])	([.4,.6],[.2,.3],[.3,.5])	([.3,.5],[.3,.5],[.1,.2])	([.3,.5],[.4,.6],[.6,.7])	([.5,.7],[.1,.2],[.4,.5])	([.4,.6],[.3,.5],[.1,.2])	([.6,.7],[.1,.2],[.3,.5])	([.2,.5],[.2,.3],[.4,.6])
δ_4	([.1,.2],[.2,.5],[.4,.6])	([.5,.7],[.2,.4],[.1,.3])	([.3,.5],[.2,.5],[.1,.3])	([.4,.6],[.2,.5],[.3,.4])	([.5,.8],[.3,.4],[.2,.3])	([.4,.6],[.2,.3],[.1,.2])	([.4,.7],[.3,.5],[.2,.4])	([.2,.4],[.3,.4],[.2,.5])

Step 3. Compute the cosine similarity measure by using definition 3.1.

By using Tables 1-5, compute the cosine similarity measure between $\mathcal{S}_{IVNHSS}^1(S, S_1)$, $\mathcal{S}_{IVNHSS}^1(S, S_2)$, $\mathcal{S}_{IVNHSS}^1(S, S_3)$, and $\mathcal{S}_{IVNHSS}^1(S, S_4)$ by using equation 3.1, such as

$$\mathcal{S}_{IVNHSS}^1(S, S_1) = 0.07007.$$

Similarly, we can find the cosine similarity measure between $\mathcal{S}_{IVNHSS}^1(S, S_2)$, $\mathcal{S}_{IVNHSS}^1(S, S_3)$, and $\mathcal{S}_{IVNHSS}^1(S, S_4)$ given as

$\mathcal{S}_{IVNHSS}^1(S, S_2) = 0.06771$, $\mathcal{S}_{IVNHSS}^1(S, S_3) = 0.06943$, and $\mathcal{S}_{IVNHSS}^1(S, S_4) = 0.06874$. This shows that $\mathcal{S}_{IVNHSS}^1(S, S_1) > \mathcal{S}_{IVNHSS}^1(S, S_3) > \mathcal{S}_{IVNHSS}^1(S, S_4) > \mathcal{S}_{IVNHSS}^1(S, S_2)$. It can be seen from this ranking alternative S_1 is most relevant and similar to S . Therefore S_1 is the best alternative for the civil engineer, the ranking of other alternatives given as $S_1 > S_3 > S_4 > S_2$.

Now we compute the set-theoretic similarity measure by using Definition 3.3 between $\mathcal{S}_{IVNHSS}^2(S, S_1)$, $\mathcal{S}_{IVNHSS}^2(S, S_2)$, $\mathcal{S}_{IVNHSS}^2(S, S_3)$, and $\mathcal{S}_{IVNHSS}^2(S, S_4)$ given as From Tables 1-5, we can find the set-theoretic similarity measure for each alternative by using definition 3.3 given as $\mathcal{S}_{IVNHSS}^2(S, S_1) = 0.06986$, $\mathcal{S}_{IVNHSS}^2(S, S_2) = 0.06379$, $\mathcal{S}_{IVNHSS}^2(S, S_3) = 0.06157$, and $\mathcal{S}_{IVNHSS}^2(S, S_4) = 0.06176$. $\mathcal{S}_{IVNHSS}^2(S, S_1) > \mathcal{S}_{IVNHSS}^2(S, S_2) > \mathcal{S}_{IVNHSS}^2(S, S_4) > \mathcal{S}_{IVNHSS}^2(S, S_3)$. It can be seen from this ranking alternative S_1 is most relevant and similar to S . Therefore S_1 is the best alternative for the civil engineer, the ranking of other alternatives given as $S_1 > S_2 > S_4 > S_3$.

5. Discussion and Comparative Analysis

In the subsequent section, we will talk over the usefulness, easiness, manageability, and assistance of the planned method. We also performed a short evaluation of the undermentioned: the planned technique and some prevailing methodologies.

5.1. Superiority of the Proposed Approach

Through this study and comparison, it could be determined that the consequences acquired by the suggested approach have been more common than either available method. Overall, the DM procedure associated with the prevailing DM methods accommodates extra information to address hesitation. Also, FS’s various hybrid structures are becoming a particular feature of NHSS, along with some appropriate circumstances added. The general info associated with the object could be stated precisely and analytically, see Table 6. Therefore, it is a suitable technique to syndicate inaccurate and ambiguous information in the DM process. Hence, the suggested approach is practical, modest, and ahead of fuzzy sets’ distinctive hybrid structures.

Table 6. Comparison between NHSS and some existing techniques

	Set	Truthiness	Indeterminacy	Falsity	Parametrization	Attributes	Sub-attributes
Zadeh [1]	FS	✓	×	×	×	✓	×
Atanassov [2]	IFS	✓	×	✓	×	✓	×
Smarandache [34]	NS	✓	✓	✓	×	✓	×
Maji et al. [21]	FSS	✓	×	×	✓	✓	×
Maji et al. [22]	IFSS	✓	×	✓	✓	✓	×
Peng et al. [26]	PFSS	✓	×	✓	✓	✓	×
Maji [35]	NSS	✓	✓	✓	✓	✓	×
Zulqarnain et al. [46]	IFHSS	✓	×	✓	✓	✓	✓
Zulqarnain et al. [47]	PFHSS	✓	×	✓	✓	✓	✓
Proposed approach	NHSS	✓	✓	✓	✓	✓	✓

It turns out that this is a contemporary issue. Why do we have to embody novel algorithms based on the proposed novel structure? Many indications compared with other existing methods; the

recommended method may be an exception. We remember the following fact: the mixed structure limits IFS, picture fuzzy sets, FS, fuzzy hesitation sets, NS, and other fuzzy sets and cannot provide complete information about the situation. But our m-polar model GmPNSS can deal with truthiness, indeterminacy, and falsity, so it is most suitable for MCDM. Due to the exaggerated multipolar neutrosophy, these three degrees are independent and provide a lot of information about alternative norms. Other similarity measures of available hybrid structures are converted into exceptional cases of GmPNSS. A comparative analysis of some already existing techniques is listed above in Table 6. Therefore, this model has more versatility and can efficiently resolve complications than intuitionistic, neutrosophic, hesitant, image, and ambiguity substitution. The similarity measure established for GmPNSS becomes better than the existing similarity measure for MCDM.

5.1 Comparative Analysis

In the following section, we recommend another algorithmic rule under NHSS by utilizing the progressed cosine similarity measure and set-theoretic similarity measure. Subsequently, we use the suggested algorithm to a realistic problem, namely the appropriate civil engineer in a company. The overall outcomes prove that the algorithmic rule is valuable and practical. It can be observed that S_1 is the most acceptable alternative for the civil engineer position. The recommended approach may be compared to other available methods. From the research findings, it has been concluded that the outcomes acquired by the planned approach exceed the consequences of the prevailing ideas. Therefore, compared to existing techniques, the established similarity measures handled the uncertain and ambiguous information competently. However, under current DM strategies, the core advantage of the planned method is that it can accommodate extra info in data comparative to existing techniques. It is also a beneficial tool to solve inaccurate and imprecise information in DM procedures. The benefit of the planned approach and related measures over current methods is evading conclusions grounded on adverse reasons.

5.2. Discussion

Zadeh's [1] FS handled the inaccurate and imprecise information using MD of sub-attributes of considered attributes for each alternative. But the FS has no evidence around the NMD of the considered parameters. Atanassov's [2] IFS accommodates the unclear and undefined objects using MD and NMD. However, IFS cannot handle the circumstances when $MD + NMD \geq 1$, conversely, is presented notion competently deals with such difficulties. Meanwhile, these theories have no information about the indeterminacy of the attributes. To overcome such problems, Smarandache [34] proposed the idea of NS. Maji et al. [21] presented the notion of FSS to deal with the parametrization of the objects, which contains uncertainty by considering the MD of the attributes. But, the presented FSS provides no information about the NMD of the thing. To overcome the presented drawback, Maji et al. [22] offered the concept of IFSS. The proposed notion handles the uncertain object more accurately by using the MD and NMD of the attributes with their parametrization, and the sum of MD and NMD does not exceed 1. To handle this scenario, Peng et al. [26] proposed the notion of PFSS by modifying the condition $MD + NMD \leq 1$ to $MD^2 + NMD^2 \leq 1$ with their parametrization. The PFSS is unable to deal with the indeterminacy of the attributes. Maji [35] introduced the concept of NSS, in which decision-makers competently solve the DM problems comparative to the above-studied theories using truthiness, falsity, and indeterminacy of the object. But all the studies mentioned above have no information about the sub-attributes of the considered attributes. So the theories discussed earlier cannot handle the scenario when characteristics have their corresponding sub-attributes. Utilizing the MD and NMD, Zulqarnain et al. [46] extended the IFSS to IFHSS and proposed the CC and WCC for IFHSS in which $MD + NMD \leq 1$ for each sub-attribute. But IFHSS cannot provide any information on the NMem values of the sub-attribute of the considered attribute. Zulqarnain et al. [47] proposed the more generalized notion of

PFHSS comparative to IFHSS. The PFHSS accommodates more uncertainty compared to IFHSS by updating the condition $MD + NMD \leq 1$ to $(\sigma_{\mathcal{F}(\tilde{a})}(\delta))^2 + (\tau_{\mathcal{F}(\tilde{a})}(\delta))^2 \leq 1$. All existing hybrid structures of FS cannot handle the indeterminacy of sub-attributes of considered n-tuple attributes. On the other hand, developed aggregation operators can accommodate the sub-attributes of considered attributes using truthness, indeterminacy, and falsity objects of sub-attributes with the following condition $0 \leq \sigma_{\mathcal{F}(\tilde{a})}(\delta), \tau_{\mathcal{F}(\tilde{a})}(\delta), \gamma_{\mathcal{F}(\tilde{a})}(\delta) \leq 3$. It may be seen that the best selection of the suggested approach is to resemble the verbalized own method, and that ensures the liableness along with the effectiveness of the recommended approach.

6. Conclusion

The interval-valued neutrosophic hypersoft set is a novel concept that is an extension of the interval-valued neutrosophic soft set. This paper studies some basic concepts such as soft set, NSS, HSS, IFHSS, PFHSS, and NHSS. We developed the idea of cosine similarity measure and set-theoretic similarity measure for IVNHSS and described their desirable properties. Furthermore, a decision-making approach has been developed for IVNHSS based on the proposed technique. To verify the effectiveness of our developed techniques, we presented an illustration to solve MCDM problems. We introduced a comprehensive comparative analysis of proposed techniques with existing methods. In the future, the concept of IVNHSS will be extended to m polar interval-valued NHSS. It will solve real-life problems such as medical diagnoses, decision-making, etc. Future research will concentrate on presenting numerous other operators under the mPIVNHSS environment to solve decision-making issues. Many other structures such as topological, algebraic, ordered structures, etc., can be developed and discussed under-considered environment.

References

1. L.A. Zadeh, Fuzzy sets, *Information and Control* **8** (1965), 338–353.
2. K. Atanassov, Intuitionistic fuzzy sets, *Fuzzy Sets, and Systems* **20** (1986), 87–96.
3. R.R. Yager, Pythagorean fuzzy subsets. *Proceedings Joint IFSA World Congress and NAFIPS Annual Meeting*. Edmonton, Canada, (2013), 57–61.
4. R.R. Yager, Pythagorean membership grades in multicriteria decision making, *IEEE Transactions on Fuzzy Systems* **22** (4) (2014), 958–965.
5. X. Zhang and Z. Xu, Extension of TOPSIS to multiple criteria decision making with pythagorean fuzzy sets, *International Journal of Intelligent Systems* **29** (12) (2014), 1061–1078.
6. G. Wei and M. Lu, Pythagorean fuzzy power aggregation operators in multiple attribute decision making, *International Journal of Intelligent Systems* **33** (1) (2018), 169–186.
7. L. Wang and N. Li, Pythagorean fuzzy interaction power bonferroni mean aggregation operators in multiple attribute decision making, *International Journal of Intelligent Systems*, **35** (1) (2020), 150–183.

8. H. Gao, M. Lu, G. Wei, Some Novel Pythagorean Fuzzy Interaction Aggregation Operators in Multiple Attribute Decision Making, *Fundamenta Informaticae* **159** (2018), 385-428.
9. G. Wei, Pythagorean fuzzy interaction aggregation operators and their application to multiple attribute decision making, *Journal of Intelligent & Fuzzy Systems* **33** (2017), 2119-2132.
10. X.L. Zhang, A novel approach based on similarity measure for pythagorean fuzzy multiple criteria group decision making, *International Journal of Intelligent Systems* **31** (6) (2016), 593-611.
11. L. Wang, H. Garg and N. Li, Pythagorean fuzzy interactive hamacher power aggregation operators for assessment of express service quality with entropy weight, *Soft Computing* (2020), 1-21.
12. L. Wang and N. Li, Continuous interval-valued pythagorean fuzzy aggregation operators for multiple attribute group decision making, *Journal of Intelligent and Fuzzy Systems* **36** (6) (2019), 6245-6263.
13. X. Peng and H. Yuan, Fundamental properties of pythagorean fuzzy aggregation operators, *Fundamenta Informaticae* **147** (2016), 415-446.
14. X. Peng and Y. Yang, Some results for pythagorean fuzzy sets, *International Journal of Intelligent Systems* **30** (11) (2015), 1133-1160.
15. H. Garg, New logarithmic operational laws and their aggregation operators for pythagorean fuzzy set and their applications, *International Journal of Intelligent Systems* **34** (1) (2019), 82-106.
16. R. Arora and H. Garg, Group decision-making method based on prioritized linguistic intuitionistic fuzzy aggregation operators and its fundamental properties, *Computational and Applied Mathematics* **38** (2) (2019), 1-36.
17. Z. Ma and Z. Xu, Symmetric pythagorean fuzzy weighted geometric/averaging operators and their application in multi criteria decision-making problems, *International Journal of Intelligent Systems* **31** (12) (2016), 1198-1219.
18. D. Molodtsov, Soft set theory first results, *Computers & Mathematics with Applications* **37** (1999), 19-31.
19. P.K. Maji, R. Biswas and A.R. Roy, Soft set theory, *Computers and Mathematics with Applications* **45** (4-5) (2003), 555-562.
20. P.K. Maji, A.R. Roy and R. Biswas, An application of soft sets in a decision making problem, *Computers and Mathematics with Applications* **44** (2002), 1077-1083.
21. P.K. Maji, R. Biswas and A.R. Roy, Fuzzy soft sets, *Journal of Fuzzy Mathematics* **9** (2001), 589-602.
22. P.K. Maji, R. Biswas and A.R. Roy, Intuitionistic fuzzy soft sets, *Journal of Fuzzy Mathematics* **9** (2001), 677-692.

23. H. Garg and R. Arora, Generalized and group-based generalized intuitionistic fuzzy soft sets with applications in decision-making, *Applied Intelligence* **48** (2018), 343–356.
24. H. Garg and R. Arora, TOPSIS method based on correlation coefficient for solving decision-making problems with intuitionistic fuzzy soft set information, *AIMS Mathematics* **5** (4) (2020), 2944–2966.
25. R. M. Zulqarnain, X. L. Xin, M. Saqlain, W. A. Khan, TOPSIS Method Based on the Correlation Coefficient of Interval-Valued Intuitionistic Fuzzy Soft Sets and Aggregation Operators with Their Application in Decision-Making, *Journal of Mathematics*, 2021, 1-16. <https://doi.org/10.1155/2021/6656858>.
26. X. Peng, Y. Yang and J. Song, Pythagoren fuzzy soft set and its application, *Computer Engineering* **41** (7) (2015), 224–229.
27. T.M. Athira, S.J. John and H. Garg, A novel entropy measure of pythagorean fuzzy soft sets, *AIMS Mathematics* **5** (2) (2020), 1050–1061.
28. T.M. Athira, S.J. John and H. Garg, Entropy and distance measures of pythagorean fuzzy soft sets and their applications, *Journal of Intelligent & Fuzzy Systems* **37** (3) (2019), 4071-4084.
29. Zulqarnain, R. M., Siddique, I., Jarad, F., Hamed, Y. S., Abualnaja, K. M., & Iampan, A. (2022). Einstein Aggregation Operators for Pythagorean Fuzzy Soft Sets with Their Application in Multiattribute Group Decision-Making. *Journal of Function Spaces*, 2022.
30. M. Riaz, K. Naeem and D. Afzal, Pythagorean m-polar fuzzy soft sets with TOPSIS method for MCGDM, *Punjab University Journal of Mathematics* **52** (3) (2020), 21-46.
31. M. Riaz, N. Khalid and D. Afzal, A similarity measure under pythagorean fuzzy soft environment with applications, *Computational and Applied Mathematics* **39** (4) (2020), 1-17.
32. Zulqarnain, R. M., Siddique, I., Ahmad, S., Iampan, A., Jovanov, G., Vranješ, Đ., & Vasiljević, J. (2021). Pythagorean fuzzy soft Einstein ordered weighted average operator in sustainable supplier selection problem. *Mathematical Problems in Engineering*, 2021.
33. Zulqarnain, R. M., Siddique, I., & El-Morsy, S. (2022). Einstein-ordered weighted geometric operator for Pythagorean fuzzy soft set with its application to solve MAGDM problem. *Mathematical Problems in Engineering*, 2022.
34. F. Smarandache, Neutrosophy: neutrosophic probability, set, and logic: analytic synthesis & synthetic analysis, Rehoboth Am. Res. Press 1998.
35. Maji, P.K. Neutrosophic soft set. *Annals of Fuzzy Mathematics and Informatics*. **2013**, 5(1), 157–168.
36. Karaaslan, F. Possibility neutrosophic soft sets and PNS-decision making method. *Applied Soft Computing Journal*. **2016**, 54, 403–414.

37. Broumi, S. Generalized Neutrosophic Soft Set. *International Journal of Computer Science, Engineering and Information Technology*. **2013**, 3(2), 17–30.
38. Deli, I.; Şubaş, Y. A ranking method of single valued neutrosophic numbers and its applications to multi-attribute decision making problems. *Int. J. Mach. Learn. & Cyber*. **2017**, 8, 1309–1322.
39. Wang, H.; Smarandache, F.; Zhang, Y. Single valued neutrosophic sets. *Int. J. Gen. Syst*, **2013**, 42, 386–394.
40. Ye, J. A multicriteria decision-making method using aggregation operators for simplified neutrosophic sets. *Journal of Intelligent and Fuzzy Systems*. **2014**, 26, 2459–2466.
41. Hashmi, M.R.; Riaz, M.; Smarandache, F. m-Polar Neutrosophic Topology with Applications to Multi criteria Decision-Making in Medical Diagnosis and Clustering Analysis. *Int. J. Fuzzy Syst*, **2019**, Dec (2019).
42. R. M. Zulqarnain, X. L. Xin, M. Saeed, F. Smarandache, N. Ahmad, Generalized Neutrosophic TOPSIS to Solve Multi-Criteria Decision-Making Problems, *Neutrosophic Sets and Systems*, 38, 276-292, (2020).
43. F. Smarandache, *Extension of Soft Set to Hypersoft Set, and then to Plithogenic Hypersoft Set*, *Neutrosophic Sets and Systems*, **22** (2018), 168-170.
44. S. Rana, M. Qayyum, M. Saeed, F. Smarandache, *Plithogenic Fuzzy Whole Hypersoft Set: Construction of Operators and their Application in Frequency Matrix Multi Attribute Decision Making Technique*, *Neutrosophic Sets and Systems*, **28** (2019), 34-50.
45. R. M. Zulqarnain, X. L. Xin, M. Saqlain, F. Smarandache, *Generalized Aggregate Operators on Neutrosophic Hypersoft Set*, *Neutrosophic Sets and Systems*, **36** (2020), 271-281.
46. R. M. Zulqarnain, X. L. Xin, M. Saeed, Extension of TOPSIS method under intuitionistic fuzzy hypersoft environment based on correlation coefficient and aggregation operators to solve decision making problem. *AIMS Mathematics* **2020**, 6 (3), 2732-2755.
47. R. M. Zulqarnain, I. Saddique, F. Jarad, R. Ali, T. Abdeljawad, Development of TOPSIS Technique Under Pythagorean Fuzzy Hypersoft Environment Based on Correlation Coefficient and Its Application Towards the Selection of Antivirus Mask in COVID-19 Pandemic. *Complexity, Complexity, Volume 2021, Article ID 6634991, 27 pages, <https://doi.org/10.1155/2021/6634991>*.
48. R. M. Zulqarnain, X. L. Xin, B. Ali, S. Broumi, S. Abdal, M. I. Ahamad, Decision-Making Approach Based on Correlation Coefficient with its Properties Under Interval-Valued Neutrosophic hypersoft set environment. *Neutrosophic Sets and Systems*, 40,12-28, (2021).

49. R. M. Zulqarnain, X. L. Xin, M. Saqlain, M. Saeed, F. Smarandache, M. I. Ahamad, Some Fundamental Operations on Interval Valued Neutrosophic Hypersoft Set with Their Properties. *Neutrosophic Sets and Systems*, 40,134-148, (2021).
50. R. M. Zulqarnain, I. Siddique, S. Ahmad, S. Ayaz, B. Said, An Algorithm Based on Correlation Coefficient Under Neutrosophic hyper soft set environment with its Application for Decision-Making, *Neutrosophic Sets and Systems*, 46, 128-140.
51. R. M. Zulqarnain, I. Siddique, R. Ali, F. Jarad, A. Iampan, Multi Criteria Decision Making Approach For Pythagorean Fuzzy Hypersoft Sets Interaction Aggregation Operators, *Mathematical Problems in Engineering*, 2021.
52. Zulqarnain, R. M., Siddique, I., Jarad, F., & Iampan, A. (2022). Aggregation Operators for Interval-Valued Intuitionistic Fuzzy Hypersoft Set with Their Application in Material Selection. *Mathematical Problems in Engineering*, 2022.
53. I. Siddique, R. M. Zulqarnain, R. Ali, F. Jarad, A. Iampan, Multicriteria Decision-Making Approach for Aggregation Operators of Pythagorean Fuzzy Hypersoft Sets, *Computational Intelligence and Neuroscience*, 2021.
54. A. Samad, R. M. Zulqarnain, E. Sermutlu, R. Ali, I. Siddique, F. Jarad , T. Abdeljawad, Selection of an Effective Hand Sanitizer to Reduce COVID-19 Effects and Extension of TOPSIS Technique Based on Correlation Coefficient under Neutrosophic Hypersoft Set, *Complexity*, 2021, Article ID 5531830, 1-22. <https://doi.org/10.1155/2021/5531830>.
55. R. M. Zulqarnain, Imran Siddique, Rifaqat Ali, Fahd Jarad, Abdul Samad, and Thabet Abdeljawad. "Neutrosophic Hypersoft Matrices with Application to Solve Multiattributive Decision-Making Problems." *Complexity* 2021 (2021).
56. S. Broumi, A. Bakali, M. Talea, Singh P. K., F. Smarandache, Energy and Spectrum Analysis of Interval-valued Neutrosophic graph Using MATLAB, *Neutrosophic Set and Systems*, vol. 24, Mar 2019, pp. 46-60.
57. S. Broumi, M. Talea, A. Bakali, F. Smarandache, Singh PK, Properties of Interval Valued Neutrosophic Graphs, In *Fuzzy Multi-criteria Decision Making Using Neutrosophic Sets, Studies in Fuzziness and Soft Computing Vol 369*, https://doi.org/10.1007/978-3-030-00045-5_8.
58. P. K. Singh , Interval-valued neutrosophic graph representation of concept lattice and its (α, β, γ) -decomposition, *Arabian Journal for Science and Engineering*, 2018, Vol. 43, Issue 2, pp. 723-740.
59. R. M. Zulqarnain, H. Garg, I. Siddique, A. Alsubie, N. Hamadneh, I. Khan, Algorithms for a generalized multi-polar neutrosophic soft set with information measures to solve medical

diagnoses and decision-making problems, *Journal of Mathematics*, Volume 2021, Article ID 6654657, 1-30, <https://doi.org/10.1155/2021/6654657>.

60. R. M. Zulqarnain, I. Siddique, A. Iampan, E. Bonyah, Algorithms For Multi-Polar Interval-Valued Neutrosophic Soft Set With Information Measures to Solve Multi-Criteria Decision-Making Problem, *Computational Intelligence and Neuroscience*, 2021.

Received: Apr 13, 2021. Accepted: Nov.10, 2022