

Received: 10.02.2016

Accepted: 03.03.2016

Year: 2016, Number: 11, Pages: 84-102

Original Article\*\*

## ON BIPOLAR SINGLE VALUED NEUTROSOPHIC GRAPHS

**Said Broumi**<sup>1,\*</sup> <broumisaid78@gmail.com>

**Mohamed Talea**<sup>1</sup> <taleamohamed@yahoo.fr>

**Assia Bakali**<sup>2</sup> <assiabakali@yahoo.fr>

**Florentin Smarandache**<sup>3</sup> <fsmarandache@gmail.com>

<sup>1</sup>Laboratory of Information processing, Faculty of Science Ben M'Sik, University Hassan II, B.P 7955, Sidi Othman, Casablanca, Morocco.

<sup>2</sup>Ecole Royale Navale-Boulevard Sour Jdid, B.P 16303 Casablanca-Morocco,

<sup>3</sup>Department of Mathematics, University of New Mexico, 705 Gurley Avenue, Gallup, NM 87301, USA

**Abstract** - In this article, we combine the concept of bipolar neutrosophic set and graph theory. We introduce the notions of bipolar single valued neutrosophic graphs, strong bipolar single valued neutrosophic graphs, complete bipolar single valued neutrosophic graphs, regular bipolar single valued neutrosophic graphs and investigate some of their related properties.

**Keywords** - Bipolar neutrosophic sets, bipolar single valued neutrosophic graph, strong bipolar single valued neutrosophic graph, complete bipolar single valued neutrosophic graph.

### 1. Introduction

Zadeh [32] coined the term 'degree of membership' and defined the concept of fuzzy set in order to deal with uncertainty. Atanassov [29, 31] incorporated the degree of non-membership in the concept of fuzzy set as an independent component and defined the concept of intuitionistic fuzzy set. Smarandache [12, 13] grounded the term 'degree of indeterminacy as an independent component and defined the concept of neutrosophic set from the philosophical point of view to deal with incomplete, indeterminate and inconsistent information in real world. The concept of neutrosophic sets is a generalization of the theory of fuzzy sets, intuitionistic fuzzy sets. Each element of a neutrosophic sets has three membership degrees including a truth membership degree, an indeterminacy membership degree, and a falsity membership degree which are within the real standard or nonstandard unit interval ]-0, 1+[. Therefore, if their range is restrained within the real standard unit interval [0, 1], the neutrosophic set is easily applied to engineering problems. For this purpose, Wang et al. [17] introduced the concept of a single valued neutrosophic set (SVNS) as a subclass of the neutrosophic set. Recently, Deli et al. [23] defined the concept of bipolar neutrosophic as an extension of the fuzzy sets, bipolar fuzzy sets, intuitionistic fuzzy sets

\*\* Edited by Mumtaz Ali and Naim Çağman (Editor-in- Chief).

\*Corresponding Author.

and neutrosophic sets studied some of their related properties including the score, certainty and accuracy functions to compare the bipolar neutrosophic sets. The neutrosophic sets theory of and their extensions have been applied in various part [1, 2, 3, 16, 18, 19, 20, 21, 25, 26, 27, 41, 42, 50, 51, 53].

A graph is a convenient way of representing information involving relationship between objects. The objects are represented by vertices and the relations by edges. When there is vagueness in the description of the objects or in its relationships or in both, it is natural that we need to design a fuzzy graph Model. The extension of fuzzy graph theory [4, 6, 11] have been developed by several researchers including intuitionistic fuzzy graphs [5, 35, 44] considered the vertex sets and edge sets as intuitionistic fuzzy sets. Interval valued fuzzy graphs [32, 34] considered the vertex sets and edge sets as interval valued fuzzy sets. Interval valued intuitionistic fuzzy graphs [8, 52] considered the vertex sets and edge sets as interval valued intuitionistic fuzzy sets. Bipolar fuzzy graphs [6, 7, 40] considered the vertex sets and edge sets as bipolar fuzzy sets. M-polar fuzzy graphs [39] considered the vertex sets and edge sets as m-polar fuzzy sets. Bipolar intuitionistic fuzzy graphs [9] considered the vertex sets and edge sets as bipolar intuitionistic fuzzy sets. But, when the relations between nodes (or vertices) in problems are indeterminate, the fuzzy graphs and their extensions are failed. For this purpose, Samarandache [10, 11] have defined four main categories of neutrosophic graphs, two based on literal indeterminacy (I), which called them; I-edge neutrosophic graph and I-vertex neutrosophic graph, these concepts are studied deeply and has gained popularity among the researchers due to its applications via real world problems [7, 14, 15, 54, 55, 56]. The two others graphs are based on (t, i, f) components and called them; The (t, i, f)-Edge neutrosophic graph and the (t, i, f)-vertex neutrosophic graph, these concepts are not developed at all. Later on, Broumi et al. [46] introduced a third neutrosophic graph model. This model allows the attachment of truth-membership (t), indeterminacy-membership (i) and falsity-membership degrees (f) both to vertices and edges, and investigated some of their properties. The third neutrosophic graph model is called single valued neutrosophic graph (SVNG for short). The single valued neutrosophic graph is the generalization of fuzzy graph and intuitionistic fuzzy graph. Also the same authors [45] introduced neighborhood degree of a vertex and closed neighborhood degree of vertex in single valued neutrosophic graph as a generalization of neighborhood degree of a vertex and closed neighborhood degree of vertex in fuzzy graph and intuitionistic fuzzy graph. Also, Broumi et al. [47] introduced the concept of interval valued neutrosophic graph as a generalization fuzzy graph, intuitionistic fuzzy graph, interval valued fuzzy graph, interval valued intuitionistic fuzzy graph and single valued neutrosophic graph and have discussed some of their properties with proof and examples. In addition Broumi et al [48] have introduced some operations such as cartesian product, composition, union and join on interval valued neutrosophic graphs and investigate some their properties. On the other hand, Broumi et al [49] have discussed a sub class of interval valued neutrosophic graph called strong interval valued neutrosophic graph, and have introduced some operations such as, cartesian product, composition and join of two strong interval valued neutrosophic graph with proofs. In the literature the study of bipolar single valued neutrosophic graphs (BSVN-graph) is still blank, we shall focus on the study of bipolar single valued neutrosophic graphs in this paper. In the present paper, bipolar neutrosophic sets are employed to study graphs and give rise to a new class of graphs called bipolar single valued neutrosophic graphs. We introduce the notions of bipolar single valued neutrosophic graphs, strong bipolar single valued neutrosophic graphs, complete bipolar single valued neutrosophic graphs, regular bipolar single valued neutrosophic graphs and investigate some of their related properties. This paper is organized as follows;

In section 2, we give all the basic definitions related bipolar fuzzy set, neutrosophic sets, bipolar neutrosophic set, fuzzy graph, intuitionistic fuzzy graph, bipolar fuzzy graph,  $N$ -graph and single valued neutrosophic graph which will be employed in later sections. In section 3, we introduce certain notions including bipolar single valued neutrosophic graphs, strong bipolar single valued neutrosophic graphs, complete bipolar single valued neutrosophic graphs, the complement of strong bipolar single valued neutrosophic graphs, regular bipolar single valued neutrosophic graphs and illustrate these notions by several examples, also we described degree of a vertex, order, size of bipolar single valued neutrosophic graphs. In section 4, we give the conclusion.

## 2. Preliminaries

In this section, we mainly recall some notions related to bipolar fuzzy set, neutrosophic sets, bipolar neutrosophic set, fuzzy graph, intuitionistic fuzzy graph, bipolar fuzzy graph,  $N$ -graph and single valued neutrosophic graph relevant to the present work. The readers are referred to [9, 12, 17, 35, 36, 38, 43, 46, 57] for further details and background.

**Definition 2.1** [12]. Let  $U$  be an universe of discourse; then the neutrosophic set  $A$  is an object having the form  $A = \{ \langle x: T_A(x), I_A(x), F_A(x) \rangle, x \in U \}$ , where the functions  $T, I, F: U \rightarrow ]0, 1^+]$  define respectively the degree of membership, the degree of indeterminacy, and the degree of non-membership of the element  $x \in U$  to the set  $A$  with the condition:

$$]0 \leq T_A(x) + I_A(x) + F_A(x) \leq 3^+. \quad (1)$$

The functions  $T_A(x), I_A(x)$  and  $F_A(x)$  are real standard or nonstandard subsets of  $]0, 1^+]$ .

Since it is difficult to apply NSs to practical problems, Wang et al. [16] introduced the concept of a SVNS, which is an instance of a NS and can be used in real scientific and engineering applications.

**Definition 2.2** [17]. Let  $X$  be a space of points (objects) with generic elements in  $X$  denoted by  $x$ . A single valued neutrosophic set  $A$  (SVNS  $A$ ) is characterized by truth-membership function  $T_A(x)$ , an indeterminacy-membership function  $I_A(x)$ , and a falsity-membership function  $F_A(x)$ . For each point  $x$  in  $X$   $T_A(x), I_A(x), F_A(x) \in [0, 1]$ . A SVNS  $A$  can be written as

$$A = \{ \langle x: T_A(x), I_A(x), F_A(x) \rangle, x \in X \} \quad (2)$$

**Definition 2.3** [9]. A bipolar neutrosophic set  $A$  in  $X$  is defined as an object of the form

$$A = \{ \langle x, T^P(x), I^P(x), F^P(x), T^N(x), I^N(x), F^N(x) \rangle: x \in X \},$$

where  $T^P, I^P, F^P: X \rightarrow [1, 0]$  and  $T^N, I^N, F^N: X \rightarrow [-1, 0]$ . The Positive membership degree  $T^P(x), I^P(x), F^P(x)$  denotes the truth membership, indeterminate membership and false membership of an element  $\in X$  corresponding to a bipolar neutrosophic set  $A$  and the negative membership degree  $T^N(x), I^N(x), F^N(x)$  denotes the truth membership, indeterminate membership and false membership of an element  $\in X$  to some implicit counter-property corresponding to a bipolar neutrosophic set  $A$ .

**Example 2.4** Let  $X = \{x_1, x_2, x_3\}$

$$A = \left\{ \begin{array}{l} \langle x_1, 0.5, 0.3, 0.1, -0.6, -0.4, -0.05 \rangle \\ \langle x_2, 0.3, 0.2, 0.7, -0.02, -0.3, -0.02 \rangle \\ \langle x_3, 0.8, 0.05, 0.4, -0.6, -0.6, -0.03 \rangle \end{array} \right\}$$

is a bipolar neutrosophic subset of  $X$ .

**Definition 2.5 [9].** Let  $A_1 = \{ \langle x, T_1^P(x), I_1^P(x), F_1^P(x), T_1^N(x), I_1^N(x), F_1^N(x) \rangle \}$  and  $A_2 = \{ \langle x, T_2^P(x), I_2^P(x), F_2^P(x), T_2^N(x), I_2^N(x), F_2^N(x) \rangle \}$  be two bipolar neutrosophic sets . Then  $A_1 \subseteq A_2$  if and only if

$$T_1^P(x) \leq T_2^P(x), I_1^P(x) \leq I_2^P(x), F_1^P(x) \geq F_2^P(x)$$

and

$$T_1^N(x) \geq T_2^N(x), I_1^N(x) \geq I_2^N(x), F_1^N(x) \leq F_2^N(x) \text{ for all } x \in X.$$

**Definition 2.6 [9].** Let  $A_1 = \{ \langle x, T_1^P(x), I_1^P(x), F_1^P(x), T_1^N(x), I_1^N(x), F_1^N(x) \rangle \}$  and  $A_2 = \{ \langle x, T_2^P(x), I_2^P(x), F_2^P(x), T_2^N(x), I_2^N(x), F_2^N(x) \rangle \}$  be two bipolar neutrosophic sets. Then  $A_1 = A_2$  if and only if

$$T_1^P(x) = T_2^P(x), I_1^P(x) = I_2^P(x), F_1^P(x) = F_2^P(x)$$

and

$$T_1^N(x) = T_2^N(x), I_1^N(x) = I_2^N(x), F_1^N(x) = F_2^N(x) \text{ for all } x \in X$$

**Definition 2.7 [9].** Let  $A_1 = \{ \langle x, T_1^P(x), I_1^P(x), F_1^P(x), T_1^N(x), I_1^N(x), F_1^N(x) \rangle \}$  and  $A_2 = \{ \langle x, T_2^P(x), I_2^P(x), F_2^P(x), T_2^N(x), I_2^N(x), F_2^N(x) \rangle \}$  be two bipolar neutrosophic sets . Then their union is defined as:

$$(A_1 \cup A_2)(x) = \left( \begin{array}{l} \max(T_1^P(x), T_2^P(x)), \frac{I_1^P(x) + I_2^P(x)}{2}, \min(F_1^P(x), F_2^P(x)) \\ \min(T_1^N(x), T_2^N(x)), \frac{I_1^N(x) + I_2^N(x)}{2}, \max(F_1^N(x), F_2^N(x)) \end{array} \right)$$

for all  $x \in X$ .

**Definition 2.8 [9].** Let  $A_1 = \{ \langle x, T_1^P(x), I_1^P(x), F_1^P(x), T_1^N(x), I_1^N(x), F_1^N(x) \rangle \}$  and  $A_2 = \{ \langle x, T_2^P(x), I_2^P(x), F_2^P(x), T_2^N(x), I_2^N(x), F_2^N(x) \rangle \}$  be two bipolar neutrosophic sets . Then their intersection is defined as:

$$(A_1 \cap A_2)(x) = \left( \begin{array}{l} \min(T_1^P(x), T_2^P(x)), \frac{I_1^P(x) + I_2^P(x)}{2}, \max(F_1^P(x), F_2^P(x)) \\ \max(T_1^N(x), T_2^N(x)), \frac{I_1^N(x) + I_2^N(x)}{2}, \min(F_1^N(x), F_2^N(x)) \end{array} \right)$$

for all  $x \in X$ .

**Definition 2.9 [9].** Let  $A_1 = \{ \langle x, T_1^P(x), I_1^P(x), F_1^P(x), T_1^N(x), I_1^N(x), F_1^N(x) \rangle : x \in X \}$  be a bipolar neutrosophic set in  $X$ . Then the complement of  $A$  is denoted by  $A^c$  and is defined by

$$T_{A^c}^P(x) = \{1^P\} - T_A^P(x), I_{A^c}^P(x) = \{1^P\} - I_A^P(x), F_{A^c}^P(x) = \{1^P\} - F_A^P(x)$$

and

$$T_{A^c}^N(x) = \{1^N\} - T_A^N(x), I_{A^c}^N(x) = \{1^N\} - I_A^N(x), F_{A^c}^N(x) = \{1^N\} - F_A^N(x)$$

**Definition 2.10 [43].** A fuzzy graph is a pair of functions  $G = (\sigma, \mu)$  where  $\sigma$  is a fuzzy subset of a non empty set  $V$  and  $\mu$  is a symmetric fuzzy relation on  $\sigma$ . i.e  $\sigma : V \rightarrow [0,1]$  and  $\mu: V \times V \rightarrow [0,1]$  such that  $\mu(uv) \leq \sigma(u) \wedge \sigma(v)$  for all  $u, v \in V$  where  $uv$  denotes the edge between  $u$  and  $v$  and  $\sigma(u) \wedge \sigma(v)$  denotes the minimum of  $\sigma(u)$  and  $\sigma(v)$ .  $\sigma$  is called the fuzzy vertex set of  $V$  and  $\mu$  is called the fuzzy edge set of  $E$ .

**Definition 2.11[38]:** By a  $N$ -graph  $G$  of a graph  $G^*$ , we mean a pair  $G = (\mu_1, \mu_2)$  where  $\mu_1$  is an  $N$ -function in  $V$  and  $\mu_2$  is an  $N$ -relation on  $E$  such that  $\mu_2(u,v) \geq \max (\mu_1(u), \mu_1(v))$  all  $u, v \in V$ .

**Definition 2.12[35] :** An Intuitionistic fuzzy graph is of the form  $G = (V, E)$  where

i.  $V = \{v_1, v_2, \dots, v_n\}$  such that  $\mu_1: V \rightarrow [0,1]$  and  $\gamma_1: V \rightarrow [0,1]$  denote the degree of membership and non-membership of the element  $v_i \in V$ , respectively, and

$$0 \leq \mu_1(v_i) + \gamma_1(v_i) \leq 1$$

for every  $v_i \in V, (i = 1, 2, \dots, n)$ ,

ii.  $E \subseteq V \times V$  where  $\mu_2: V \times V \rightarrow [0,1]$  and  $\gamma_2: V \times V \rightarrow [0,1]$  are such that

$$\mu_2(v_i, v_j) \leq \min [\mu_1(v_i), \mu_1(v_j)] \text{ and } \gamma_2(v_i, v_j) \geq \max [\gamma_1(v_i), \gamma_1(v_j)]$$

and  $0 \leq \mu_2(v_i, v_j) + \gamma_2(v_i, v_j) \leq 1$  for every  $(v_i, v_j) \in E, (i, j = 1, 2, \dots, n)$

**Definition 2.13 [57].** Let  $X$  be a non-empty set. A bipolar fuzzy set  $A$  in  $X$  is an object having the form  $A = \{(x, \mu_A^P(x), \mu_A^N(x)) \mid x \in X\}$ , where  $\mu_A^P(x): X \rightarrow [0, 1]$  and  $\mu_A^N(x): X \rightarrow [-1, 0]$  are mappings.

**Definition 2.14 [ 57]** Let  $X$  be a non-empty set. Then we call a mapping

$$A = (\mu_A^P, \mu_A^N): X \times X \rightarrow [-1, 0] \times [0, 1]$$

a bipolar fuzzy relation on  $X$  such that  $\mu_A^P(x, y) \in [0, 1]$  and  $\mu_A^N(x, y) \in [-1, 0]$ .

**Definition 2.15 [36].** Let  $A = (\mu_A^P, \mu_A^N)$  and  $B = (\mu_B^P, \mu_B^N)$  be bipolar fuzzy sets on a set  $X$ . If  $A = (\mu_A^P, \mu_A^N)$  is a bipolar fuzzy relation on a set  $X$ , then  $A = (\mu_A^P, \mu_A^N)$  is called a bipolar fuzzy relation on

$$B = (\mu_B^P, \mu_B^N) \text{ if } \mu_B^P(x, y) \leq \min(\mu_A^P(x), \mu_A^P(y))$$

and

$$\mu_B^N(x, y) \geq \max(\mu_A^N(x), \mu_A^N(y)) \text{ for all } x, y \in X.$$

A bipolar fuzzy relation  $A$  on  $X$  is called symmetric if  $\mu_A^P(x, y) = \mu_A^P(y, x)$  and  $\mu_A^N(x, y) = \mu_A^N(y, x)$  for all  $x, y \in X$ .

**Definition 2.16 [36].** A bipolar fuzzy graph of a graph  $G^* = (V, E)$  is a pair  $G = (A, B)$ , where  $A = (\mu_A^P, \mu_A^N)$  is a bipolar fuzzy set in  $V$  and  $B = (\mu_B^P, \mu_B^N)$  is a bipolar fuzzy set on  $E \subseteq V \times V$  such that  $\mu_B^P(xy) \leq \min\{\mu_A^P(x), \mu_A^P(y)\}$  for all  $xy \in E, \mu_B^N(xy) \geq$

$\min\{\mu_A^N(x), \mu_A^N(y)\}$  for all  $xy \in E$  and  $\mu_B^P(xy) = \mu_B^N(xy) = 0$  for all  $xy \in \tilde{V}^2 - E$ . Here  $A$  is called bipolar fuzzy vertex set of  $V$ ,  $B$  the bipolar fuzzy edge set of  $E$ .

**Definition 2.17 [46]** A single valued neutrosophic graph (SVNG) of a graph  $G^* = (V, E)$  is a pair  $G = (A, B)$ , where

1.  $V = \{v_1, v_2, \dots, v_n\}$  such that  $T_A: V \rightarrow [0, 1]$ ,  $I_A: V \rightarrow [0, 1]$  and  $F_A: V \rightarrow [0, 1]$  denote the degree of truth-membership, degree of indeterminacy-membership and falsity-membership of the element  $v_i \in V$ , respectively, and

$$0 \leq T_A(v_i) + I_A(v_i) + F_A(v_i) \leq 3$$

for every  $v_i \in V$  ( $i=1, 2, \dots, n$ )

2.  $E \subseteq V \times V$  where  $T_B: V \times V \rightarrow [0, 1]$ ,  $I_B: V \times V \rightarrow [0, 1]$  and  $F_B: V \times V \rightarrow [0, 1]$  are such that

$$T_B(v_i, v_j) \leq \min [T_A(v_i), T_A(v_j)], I_B(v_i, v_j) \geq \max [I_A(v_i), I_A(v_j)]$$

and

$$F_B(v_i, v_j) \geq \max [F_A(v_i), F_A(v_j)]$$

and

$$0 \leq T_B(v_i, v_j) + I_B(v_i, v_j) + F_B(v_i, v_j) \leq 3$$

for every  $(v_i, v_j) \in E$  ( $i, j = 1, 2, \dots, n$ )

**Definition 2.18 [46]:** Let  $G=(V, E)$  be a single valued neutrosophic graph. Then the degree of a vertex  $v$  is defined by  $d(v) = (d_T(v), d_I(v), d_F(v))$  where

$$d_T(v) = \sum_{u \neq v} T_B(u, v), d_I(v) = \sum_{u \neq v} I_B(u, v) \text{ and } d_F(v) = \sum_{u \neq v} F_B(u, v)$$

### 3. Bipolar Single Valued Neutrosophic Graph

**Definition 3.1.** Let  $X$  be a non-empty set. Then we call a mapping  $A = (x, T^P(x), I^P(x), F^P(x), T^N(x), I^N(x), F^N(x)): X \times X \rightarrow [-1, 0] \times [0, 1]$  a bipolar single valued neutrosophic relation on  $X$  such that  $T_A^P(x, y) \in [0, 1]$ ,  $I_A^P(x, y) \in [0, 1]$ ,  $F_A^P(x, y) \in [0, 1]$ , and  $T_A^N(x, y) \in [-1, 0]$ ,  $I_A^N(x, y) \in [-1, 0]$ ,  $F_A^N(x, y) \in [-1, 0]$ .

**Definition 3.2.** Let  $A = (T_A^P, I_A^P, F_A^P, T_A^N, I_A^N, F_A^N)$  and  $B = (T_B^P, I_B^P, F_B^P, T_B^N, I_B^N, F_B^N)$  be bipolar single valued neutrosophic graph on a set  $X$ . If  $B = (T_B^P, I_B^P, F_B^P, T_B^N, I_B^N, F_B^N)$  is a bipolar single valued neutrosophic relation on  $A = (T_A^P, I_A^P, F_A^P, T_A^N, I_A^N, F_A^N)$  then

$$T_B^P(x, y) \leq \min(T_A^P(x), T_A^P(y)), T_B^N(x, y) \geq \max(T_A^N(x), T_A^N(y))$$

$$I_B^P(x, y) \geq \max(I_A^P(x), I_A^P(y)), I_B^N(x, y) \leq \min(I_A^N(x), I_A^N(y))$$

$$F_B^P(x, y) \geq \max(F_A^P(x), F_A^P(y)), F_B^N(x, y) \leq \min(F_A^N(x), F_A^N(y)) \text{ for all } x, y \in X.$$

A bipolar single valued neutrosophic relation  $B$  on  $X$  is called symmetric if

$$T_B^P(x, y) = T_B^P(y, x), I_B^P(x, y) = I_B^P(y, x), F_B^P(x, y) = F_B^P(y, x)$$

and

$$T_B^N(x, y) = T_B^N(y, x), I_B^N(x, y) = I_B^N(y, x), F_B^N(x, y) = F_B^N(y, x), \text{ for all } x, y \in X.$$

**Definition 3.3.** A bipolar single valued neutrosophic graph of a graph  $G^* = (V, E)$  is a pair  $G = (A, B)$ , where  $A = (T_A^P, I_A^P, F_A^P, T_A^N, I_A^N, F_A^N)$  is a bipolar single valued neutrosophic set in  $V$  and  $B = (T_B^P, I_B^P, F_B^P, T_B^N, I_B^N, F_B^N)$  is a bipolar single valued neutrosophic set in  $\tilde{V}^2$  such that

$$\begin{aligned} T_B^P(v_i, v_j) &\leq \min(T_A^P(v_i), T_A^P(v_j)) \\ I_B^P(v_i, v_j) &\geq \max(I_A^P(v_i), I_A^P(v_j)) \\ F_B^P(v_i, v_j) &\geq \max(F_A^P(v_i), F_A^P(v_j)) \end{aligned}$$

and

$$\begin{aligned} T_B^N(v_i, v_j) &\geq \max(T_A^N(v_i), T_A^N(v_j)) \\ I_B^N(v_i, v_j) &\leq \min(I_A^N(v_i), I_A^N(v_j)) \\ F_B^N(v_i, v_j) &\leq \min(F_A^N(v_i), F_A^N(v_j)) \end{aligned}$$

for all  $v_i v_j \in \tilde{V}^2$ .

**Notation :** An edge of BSVNG is denoted by  $e_{ij} \in E$  or  $v_i v_j \in E$


Here the sextuple  $(v_i, T_A^P(v_i), I_A^P(v_i), F_A^P(v_i), T_A^N(v_i), I_A^N(v_i), F_A^N(v_i))$  denotes the positive degree of truth-membership, the positive degree of indeterminacy-membership, the positive degree of falsity-membership, the negative degree of truth-membership, the negative degree of indeterminacy-membership, the negative degree of falsity- membership of the vertex  $v_i$ .

The sextuple  $(e_{ij}, T_B^P, I_B^P, F_B^P, T_B^N, I_B^N, F_B^N)$  denotes the positive degree of truth-membership, the positive degree of indeterminacy-membership, the positive degree of falsity-membership, the negative degree of truth-membership, the negative degree of indeterminacy-membership, the negative degree of falsity- membership of the edge relation  $e_{ij} = (v_i, v_j)$  on  $V \times V$ .

**Note 1.** (i) When  $T_A^P = I_A^P = F_A^P = 0$  and  $T_A^N = I_A^N = F_A^N = 0$  for some  $i$  and  $j$ , then there is no edge between  $v_i$  and  $v_j$ .

Otherwise there exists an edge between  $v_i$  and  $v_j$ .

(ii) If one of the inequalities is not satisfied in (1) and (2), then  $G$  is not an BSVNG




**Figure 1:** Bipolar single valued neutrosophic graph.

**Proposition 3.5:** A bipolar single valued neutrosophic graph is the generalization of fuzzy graph

**Proof:** Suppose  $G = (A, B)$  be a bipolar single valued neutrosophic graph. Then by setting the positive indeterminacy-membership, positive falsity-membership and negative truth-membership, negative indeterminacy-membership, negative falsity-membership values of vertex set and edge set equals to zero reduces the bipolar single valued neutrosophic graph to fuzzy graph.

**Example 3.6:**




**Figure 2: Fuzzy graph**

**Proposition 3.7:** A bipolar single valued neutrosophic graph is the generalization of intuitionistic fuzzy graph

**Proof:** Suppose  $G = (A, B)$  be a bipolar single valued neutrosophic graph. Then by setting the positive indeterminacy-membership, negative truth-membership, negative indeterminacy-membership, negative falsity-membership values of vertex set and edge set equals to zero reduces the bipolar single valued neutrosophic graph to intuitionistic fuzzy graph.

**Example 3.8**




**Figur 3: Intuitionistic fuzzy graph**

**Proposition 3.9:** A bipolar single valued neutrosophic graph is the generalization of single valued neutrosophic graph

**Proof:** Suppose  $G = (A, B)$  be a bipolar single valued neutrosophic graph. Then by setting the negative truth-membership, negative indeterminacy-membership, negative falsity-membership values of vertex set and edge set equals to zero reduces the bipolar single valued neutrosophic graph to single valued neutrosophic graph.



**Example 3.10**




**Figure 4:** Single valued neutrosophic graph

**Proposition 3.11:** A bipolar single valued neutrosophic graph is the generalization of bipolar intuitionistic fuzz graph

**Proof:** Suppose  $G = (A, B)$  be a bipolar single valued neutrosophic graph. Then by setting the positive indeterminacy-membership, negative indeterminacy-membership values of vertex set and edge set equals to zero reduces the bipolar single valued neutrosophic graph to bipolar intuitionistic fuzzy graph

**Example 3.12**




**Figure 5:** Bipolar intuitionistic fuzzy graph.

**Proposition 3.13:** A bipolar single valued neutrosophic graph is the generalization of  $N$ -graph

**Proof:** Suppose  $G = (A, B)$  be a bipolar single valued neutrosophic graph. Then by setting the positive degree membership such truth-membership, indeterminacy- membership, falsity-membership and negative indeterminacy-membership, negative falsity-membership values of vertex set and edge set equals to zero reduces the single valued neutrosophic graph to  $N$ -graph.

**Example 3.14:**




**Figure 6:** N- graph

**Definition 3.15.** A bipolar single valued neutrosophic graph that has neither self loops nor parallel edge is called simple bipolar single valued neutrosophic graph.

**Definition 3.16.** A bipolar single valued neutrosophic graph is said to be connected if every pair of vertices has at least one bipolar single valued neutrosophic graph between them, otherwise it is disconnected.

**Definition 3.17.** When a vertex  $v_i$  is end vertex of some edges  $(v_i, v_j)$  of any BSVN-graph  $G=(A, B)$ . Then  $v_i$  and  $(v_i, v_j)$  are said to be **incident** to each other.



**Figure 7 :** Incident BSVN-graph

In this graph  $v_2v_3, v_3v_4$  and  $v_3v_5$  are incident on  $v_3$ .

**Definition 3.18** Let  $G=(V, E)$  be a bipolar single valued neutrosophic graph. Then the degree of any vertex  $v$  is sum of positive degree of truth-membership, positive sum of degree of indeterminacy-membership, positive sum of degree of falsity-membership, negative degree of truth-membership, negative sum of degree of indeterminacy-membership, and negative sum of degree of falsity-membership of all those edges which are incident on vertex  $v$  denoted by  $d(v)=(d_T^P(v), d_I^P(v), d_F^P(v), d_T^N(v), d_I^N(v), d_F^N(v))$  where

$$d_T^P(v) = \sum_{u \neq v} T_B^P(u, v) \text{ denotes the positive T- degree of a vertex } v,$$

$$d_I^P(v) = \sum_{u \neq v} I_B^P(u, v) \text{ denotes the positive I- degree of a vertex } v,$$

$$d_F^P(v) = \sum_{u \neq v} F_B^P(u, v) \text{ denotes the positive F- degree of a vertex } v,$$

$$d_T^N(v) = \sum_{u \neq v} T_B^N(u, v) \text{ denotes the negative T- degree of a vertex } v,$$

$$d_I^N(v) = \sum_{u \neq v} I_B^N(u, v) \text{ denotes the negative I- degree of a vertex } v,$$

$d_F^N(v) = \sum_{u \neq v} F_B^N(u, v)$  denotes the negative F- degree of a vertex  $v$

**Definition 3.19:** The minimum degree of  $G$  is

$$\delta(G) = (\delta_T^P(G), \delta_I^P(G), \delta_F^P(G), \delta_T^N(G), \delta_I^N(G), \delta_F^N(G))$$

where

- $\delta_T^P(G) = \wedge \{d_T^P(v) \mid v \in V\}$  denotes the minimum positive T- degree,
- $\delta_I^P(G) = \wedge \{d_I^P(v) \mid v \in V\}$  denotes the minimum positive I- degree,
- $\delta_F^P(G) = \wedge \{d_F^P(v) \mid v \in V\}$  denotes the minimum positive F- degree,
- $\delta_T^N(G) = \wedge \{d_T^N(v) \mid v \in V\}$  denotes the minimum negative T- degree,
- $\delta_I^N(G) = \wedge \{d_I^N(v) \mid v \in V\}$  denotes the minimum negative I- degree,
- $\delta_F^N(G) = \wedge \{d_F^N(v) \mid v \in V\}$  denotes the minimum negative F- degree


**Definition 3.20:** The maximum degree of  $G$  is

$$\Delta(G) = (\Delta_T^P(G), \Delta_I^P(G), \Delta_F^P(G), \Delta_T^N(G), \Delta_I^N(G), \Delta_F^N(G))$$

where

- $\Delta_T^P(G) = \vee \{d_T^P(v) \mid v \in V\}$  denotes the maximum positive T- degree,
- $\Delta_I^P(G) = \vee \{d_I^P(v) \mid v \in V\}$  denotes the maximum positive I- degree,
- $\Delta_F^P(G) = \vee \{d_F^P(v) \mid v \in V\}$  denotes the maximum positive F- degree,
- $\Delta_T^N(G) = \vee \{d_T^N(v) \mid v \in V\}$  denotes the maximum negative T- degree,
- $\Delta_I^N(G) = \vee \{d_I^N(v) \mid v \in V\}$  denotes the maximum negative I- degree,
- $\Delta_F^N(G) = \vee \{d_F^N(v) \mid v \in V\}$  denotes the maximum negative F- degree

**Example 3.21.** Let us consider a bipolar single valued neutrosophic graph  $G = (A, B)$  of  $G^* = (V, E)$ , such that  $V = \{v_1, v_2, v_3, v_4\}$ ,  $E = \{(v_1, v_2), (v_2, v_3), (v_3, v_4), (v_4, v_1)\}$



**Figure 8:** Degree of a bipolar single valued neutrosophic graph  $G$ .

In this example, the degree of  $v_1$  is  $(0.3, 0.6, 1.1, -0.4, -0.6, -0.6)$ . the degree of  $v_2$  is  $(0.2, 0.6, 1.2, -0.3, -0.9, -0.8)$ . the degree of  $v_3$  is  $(0.2, 0.8, 1.2, -0.2, -1.2, -1.2)$ . the degree of  $v_4$  is  $(0.3, 0.8, 1.1, -0.3, -0.9, -1)$

Order and size of a bipolar single valued neutrosophic graph is an important term in bipolar single valued neutrosophic graph theory. They are defined below.


**Definition 3.22:** Let  $G = (V, E)$  be a BSVNG. The order of  $G$ , denoted  $O(G)$  is defined as  $O(G) = (O_T^P(G), O_I^P(G), O_F^P(G), O_T^N(G), O_I^N(G), O_F^N(G))$ , where

$$\begin{aligned}
 O_T^P(G) &= \sum_{v \in V} T_1^P(v) \text{ denotes the positive T- order of a vertex } v, \\
 O_I^P(G) &= \sum_{v \in V} I_1^P(v) \text{ denotes the positive I- order of a vertex } v, \\
 O_F^P(G) &= \sum_{v \in V} F_1^P(v) \text{ denotes the positive F- order of a vertex } v, \\
 O_T^N(G) &= \sum_{v \in V} T_1^N(v) \text{ denotes the negative T- order of a vertex } v, \\
 O_I^N(G) &= \sum_{v \in V} I_1^N(v) \text{ denotes the negative I- order of a vertex } v, \\
 O_F^N(G) &= \sum_{v \in V} F_1^N(v) \text{ denotes the negative F- order of a vertex } v.
 \end{aligned}$$

**Definition 3.23:** Let  $G=(V, E)$  be a BSVNG. The size of  $G$ , denoted  $S(G)$  is defined as  $S(G)=(S_T^P(G), S_I^P(G), S_F^P(G), S_T^N(G), S_I^N(G), S_F^N(G))$ , where

$$\begin{aligned}
 S_T^P(G) &= \sum_{u \neq v} T_2^P(u, v) \text{ denotes the positive T- size of a vertex } v, \\
 S_I^P(G) &= \sum_{u \neq v} I_2^P(u, v) \text{ denotes the positive I- size of a vertex } v, \\
 S_F^P(G) &= \sum_{u \neq v} F_2^P(u, v) \text{ denotes the positive F- size of a vertex } v, \\
 S_T^N(G) &= \sum_{u \neq v} T_2^N(u, v) \text{ denotes the negative T- size of a vertex } v, \\
 S_I^N(G) &= \sum_{u \neq v} I_2^N(u, v) \text{ denotes the negative I- size of a vertex } v, \\
 S_F^N(G) &= \sum_{u \neq v} F_2^N(u, v) \text{ denotes the negative F- size of a vertex } v.
 \end{aligned}$$

**Definition 3.24** A bipolar single valued neutrosophic graph  $G=(V, E)$  is called constant if degree of each vertex is  $k=(k_1, k_2, k_3, k_4, k_5, k_6)$ . That is,  $d(v)=(k_1, k_2, k_3, k_4, k_5, k_6)$  for all  $v \in V$ .



**Figure 9:** Constant bipolar single valued neutrosophic graph  $G$ .

In this example, the degree of  $v_1, v_2, v_3, v_4$  is  $(0.2, 0.6, 1.2, -0.4, -0.6, -1.4)$ .

$$\begin{aligned}
 O(G) &= (0.8, 1, 1.8, -1.5, -1.1, -1.8) \\
 S(G) &= (0.4, 1.2, 2.4, -0.7, -1.2, -2.8)
 \end{aligned}$$

**Remark 3.25.**  $G$  is a  $(k_i, k_j, k_l, k_m, k_n, k_o)$ -constant BSVNG iff  $\delta = \Delta = k$ , where  $k = k_i + k_j + k_l + k_m + k_n + k_o$ .

**Definition 3.26.** A bipolar single valued neutrosophic graph  $G=(A, B)$  is called strong bipolar single valued neutrosophic graph if


$$\begin{aligned}
 T_B^P(u, v) &= \min(T_A^P(u), T_A^P(v)), \\
 I_B^P(u, v) &= \max(I_A^P(u), I_A^P(v)), \\
 F_B^P(u, v) &= \max(F_A^P(u), F_A^P(v)), \\
 T_B^N(u, v) &= \max(T_A^N(u), T_A^N(v)),
 \end{aligned}$$

$$I_B^N(u, v) = \min(I_A^N(u), I_A^N(v)),$$

$$F_B^N(u, v) = \min(F_A^N(u), F_A^N(v))$$

for all  $(u, v) \in E$

**Example 3.27.** Consider a strong BSVN-graph  $G$  such that  $V = \{v_1, v_2, v_3, v_4\}$  and  $E = \{(v_1, v_2), (v_2, v_3), (v_3, v_4), (v_4, v_1)\}$



**Figure 10:** Strong bipolar single valued neutrosophic graph  $G$ .

**Definition 3.28.** A bipolar single valued neutrosophic graph  $G = (A, B)$  is called complete if

$$T_B^P(u, v) = \min(T_A^P(u), T_A^P(v)),$$

$$I_B^P(u, v) = \max(I_A^P(u), I_A^P(v)),$$

$$F_B^P(u, v) = \max(F_A^P(u), F_A^P(v)),$$


$$T_B^N(u, v) = \max(T_A^N(u), T_A^N(v)),$$

$$I_B^N(u, v) = \min(I_A^N(u), I_A^N(v)),$$

$$F_B^N(u, v) = \min(F_A^N(u), F_A^N(v))$$

for all  $u, v \in V$ .

**Example 3.29.** Consider a complete BSVN-graph  $G$  such that  $V = \{v_1, v_2, v_3, v_4\}$  and  $E = \{(v_1, v_2), (v_2, v_3), (v_3, v_4), (v_4, v_1), (v_1, v_3), (v_2, v_4)\}$



**Figure 11:** Complete bipolar single valued neutrosophic graph  $G$ .

$$d(v_1) = (0.5, 0.8, 1.4, -0.9, -1, -1.5)$$

$$d(v_2) = (0.4, 0.9, 1.5, -1.2, -1, -1.6)$$

$$d(v_3) = (0.4, 0.9, 1.5, -0.7, -1.3, -1.7)$$

$$d(\mathbf{v}_4) = (0.5, 0.8, 1.4, -0.6, -1.1, -1.6)$$

**Definition 3.30.** The complement of a bipolar single valued neutrosophic graph  $G = (A, B)$  of a graph  $G^* = (V, E)$  is a bipolar single valued neutrosophic graph  $\bar{G} = (\bar{A}, \bar{B})$  of  $\bar{G}^* = (V, V \times V)$ , where  $\bar{A} = A = (T_A^P, I_A^P, F_A^P, T_A^N, I_A^N, F_A^N)$  and  $\bar{B} = (\bar{T}_B^P, \bar{I}_B^P, \bar{F}_B^P, \bar{T}_B^N, \bar{I}_B^N, \bar{F}_B^N)$  is defined by

$$\begin{aligned} \bar{T}_B^P(u,v) &= \min(T_A^P(u), T_A^P(v)) - T_B^P(u, v) \text{ for all } u, v \in V, uv \in \tilde{V}^2 \\ \bar{I}_B^P(u,v) &= \max(I_A^P(u), I_A^P(v)) - I_B^P(u, v) \text{ for all } u, v \in V, uv \in \tilde{V}^2 \\ \bar{F}_B^P(u,v) &= \max(F_A^P(u), F_A^P(v)) - F_B^P(u, v) \text{ for all } u, v \in V, uv \in \tilde{V}^2 \\ \bar{T}_B^N(u,v) &= \max(T_A^N(u), T_A^N(v)) - T_B^N(u, v) \text{ for all } u, v \in V, uv \in \tilde{V}^2 \\ \bar{I}_B^N(u,v) &= \min(I_A^N(u), I_A^N(v)) - I_B^N(u, v) \text{ for all } u, v \in V, uv \in \tilde{V}^2 \\ \bar{F}_B^N(u,v) &= \min(F_A^N(u), F_A^N(v)) - F_B^N(u, v) \text{ for all } u, v \in V, uv \in \tilde{V}^2 \end{aligned}$$

**Proposition 3.31:** The complement of complete BSVN-graph is a BSVN-graph with no edge. Or if  $G$  is a complete then in  $\bar{G}$  the edge is empty.

**Proof.** Let  $G = (V, E)$  be a complete BSVN-graph.  $T_B^P(u, v) = \min(T_A^P(u), T_A^P(v))$ ,  
So  $T_B^P(u, v) = \min(T_A^P(u), T_A^P(v))$ ,  $T_B^N(u, v) = \max(T_A^N(u), T_A^N(v))$ ,

$$\begin{aligned} I_B^P(u, v) &= \max(T_A^P(u), T_A^P(v)), & I_B^N(u, v) &= \min(I_A^N(u), I_A^N(v)), \\ F_B^P(u, v) &= \max(T_A^P(u), T_A^P(v)), & F_B^N(u, v) &= \min(F_A^N(u), F_A^N(v)) \end{aligned}$$

for all  $u, v \in V$ . Hence in  $\bar{G}$ ,

$$\begin{aligned} \bar{T}_B^P &= \min(T_A^P(u), T_A^P(v)) - T_B^P(u, v) \text{ for all } u, v \in V \\ &= \min(T_A^P(u), T_A^P(v)) - \min(T_A^P(u), T_A^P(v)) \text{ for all } u, v \in V \\ &= 0 \text{ for all } u, v \in V \end{aligned}$$

and

$$\begin{aligned} \bar{I}_B^P &= \max(I_A^P(u), I_A^P(v)) - I_B^P(u, v) \text{ for all } u, v \in V \\ &= \max(I_A^P(u), I_A^P(v)) - \max(I_A^P(u), I_A^P(v)) \text{ for all } u, v \in V \\ &= 0 \text{ for all } u, v \in V \end{aligned}$$

Also

$$\begin{aligned} \bar{F}_B^P &= \max(F_A^P(u), F_A^P(v)) - F_B^P(u, v) \text{ for all } u, v \in V \\ &= \max(F_A^P(u), F_A^P(v)) - \max(F_A^P(u), F_A^P(v)) \text{ for all } u, v \in V \\ &= 0 \text{ for all } u, v \in V \end{aligned}$$

Similarly

$$\begin{aligned} \bar{T}_B^N &= \max(T_A^N(u), T_A^N(v)) - T_B^N(u, v) \text{ for all } u, v \in V \\ &= \max(T_A^N(u), T_A^N(v)) - \max(T_A^N(u), T_A^N(v)) \text{ for all } u, v \in V \\ &= 0 \text{ for all } u, v \in V \end{aligned}$$

and

$$\begin{aligned} \bar{I}_B^P &= \min(I_A^N(u), I_A^N(v)) - I_B^N(u, v) \text{ for all } u, v \in V \\ &= \min(I_A^N(u), I_A^N(v)) - \min(I_A^N(u), I_A^N(v)) \text{ for all } u, v \in V \\ &= 0 \text{ for all } u, v \in V \end{aligned}$$

Also

$$\begin{aligned} \bar{F}_B^N &= \min(F_A^N(u), F_A^N(v)) - F_B^N(u, v) \text{ for all } u, v \in V \\ &= \min(F_A^N(u), F_A^N(v)) - \min(F_A^N(u), F_A^N(v)) \text{ for all } u, v \in V \\ &= 0 \text{ for all } u, v \in V \end{aligned}$$

$(\bar{T}_B^P, \bar{I}_B^P, \bar{F}_B^P, \bar{T}_B^N, \bar{I}_B^N, \bar{F}_B^N) = (0, 0, 0, 0, 0)$ . Hence the edge set of  $\bar{G}$  is empty if  $G$  is a complete BSVNG.


**Definition 3.32:** A regular BSVN-graph is a BSVN-graph where each vertex has the same number of open neighbors degree.  $d_N(v) = (d_{NT}^P(v), d_{NI}^P(v), d_{NF}^P(v), d_{NT}^N(v), d_{NI}^N(v), d_{NF}^N(v))$ .

The following example shows that there is no relationship between regular BSVN-graph and a constant BSVN-graph

**Example 3.33.** Consider a graph  $G^*$  such that  $V = \{v_1, v_2, v_3, v_4\}$ ,  $E = \{v_1v_2, v_2v_3, v_3v_4, v_4v_1\}$ . Let  $A$  be a single valued neutrosophic subset of  $V$  and let  $B$  a single valued neutrosophic subset of  $E$  denoted by

	$v_1$	$v_2$	$v_3$	$v_4$
$T_A^P$	0.2	0.2	0.2	0.2
$I_A^P$	0.2	0.2	0.2	0.2
$F_A^P$	0.4	0.4	0.4	0.4
$T_A^N$	-0.4	-0.4	-0.4	-0.4
$I_A^N$	-0.1	-0.4	-0.1	-0.1
$F_A^N$	-0.4	-0.4	-0.4	-0.4

	$v_1v_2$	$v_2v_3$	$v_3v_4$	$v_4v_1$
$T_B^P$	0.1	0.1	0.1	0.2
$I_B^P$	0.3	0.3	0.5	0.3
$F_B^P$	0.6	0.6	0.6	0.5
$T_B^N$	-0.2	-0.1	-0.1	-0.2
$I_B^N$	-0.3	-0.6	-0.6	-0.3
$F_B^N$	-0.5	-0.7	-0.7	-0.5



**Figure 12:** Regular bipolar single valued neutrosophic graph  $G$ .

By routing calculations show that G is regular BSVN-graph since each open neighbors degree is same , that is (0.4, 0.4, 0.8 , -0.8, -0.2,-0.8). But it is not constant BSVN-graph since degree of each vertex is not same.

**Definition 3.34 :**Let  $G=(V, E)$  be a bipolar single valued neutrosophic graph. Then the totally degree of a vertex  $v \in V$  is defined by

$$td(v)=(td_T^P(v), td_I^P(v), td_F^P(v), td_T^N(v), td_I^N(v), td_F^N(v))$$

where

$$td_T^P(v) = \sum_{u \neq v} T_B^P(u, v) + T_A^P(v)$$

denotes the totally positive T- degree of a vertex v,

$$td_I^P(v) = \sum_{u \neq v} I_B^P(u, v) + I_A^P(v)$$

denotes the totally positive I- degree of a vertex v,

$$td_F^P(v) = \sum_{u \neq v} F_B^P(u, v) + F_A^P(v)$$

denotes the totally positive F- degree of a vertex v,

$$td_T^N(v) = \sum_{u \neq v} T_B^N(u, v) + T_A^N(v)$$

denotes the totally negative T- degree of a vertex v,

$$td_I^N(v) = \sum_{u \neq v} I_B^N(u, v) + I_A^N(v)$$


denotes the totally negative I- degree of a vertex v,

$$td_F^N(v) = \sum_{u \neq v} F_B^N(u, v) + F_A^N(v)$$

denotes the totally negative F- degree of a vertex v

If each vertex of G has totally same degree  $\mathbf{m}=(m_1, m_2, m_3, m_4, m_5, m_6)$ , then G is called a **m**-totally constant BSVN-Graph.

**Example 3.35.** Let us consider a bipolar single valued neutrosophic graph  $G= (A, B)$  of  $G^* = (V, E)$ , such that  $V = \{v_1, v_2, v_3, v_4\}$ ,  $E = \{(v_1, v_2),(v_2, v_3), (v_3, v_4), (v_4, v_1)\}$




**Figure 13:** Totally degree of a bipolar single valued neutrosophic graph G.

In this example, the totally degree of  $v_1$  is (0.5, 0.8, 1.4, -0.8, -0.7, -1.4). The totally degree of  $v_2$  is (0.3, 0.9, 1.7, -0.9, -1.1, -1.5). The totally degree of  $v_3$  is (0.4, 1.1, 1.7, -0.5, -1.7, -2). The totally degree of  $v_4$  is (0.6, 1, 1.5, -0.5, -1.1, -1.7).

**Definition 3.36:** A totally regular BSVN-graph is a BSVN-graph where each vertex has the same number of closed neighbors degree, it is noted  $d[v]$ .

**Example 3.37.** Let us consider a BSVN-graph  $G= (A, B)$  of  $G^* = (V, E)$ , such that  $V = \{v_1, v_2, v_3, v_4\}$  and  $E = \{(v_1, v_2),(v_2, v_3), (v_3, v_4), (v_4, v_1)\}$





**Figure 14:** Degree of a bipolar single valued neutrosophic graph G.

By routing calculations we show that G is regular BSVN-graph since the degree of  $v_1, v_2, v_3,$  and  $v_4$  is  $(0.2, 0.6, 1.2, -0.4, -0.6, -1)$ . It is neither totally regular BSVN-graph not constant BSVN-graph.

## 4. Conclusion

In this paper, we have introduced the concept of bipolar single valued neutrosophic graphs and described degree of a vertex, order, size of bipolar single valued neutrosophic graphs, also we have introduced the notion of complement of a bipolar single valued neutrosophic graph, strong bipolar single valued neutrosophic graph, complete bipolar single valued neutrosophic graph, regular bipolar single valued neutrosophic graph. Further, we are going to study some types of single valued neutrosophic graphs such irregular and totally irregular single valued neutrosophic graphs and bipolar single valued neutrosophic graphs.

## References

1. A. Q. Ansari, R.Biswas & S. Aggarwal, (2012). Neutrosophic classifier: An extension of fuzzy classifier. Elsevier- AppliedSoft Computing, 13, p.563-573 (2013) <http://dx.doi.org/10.1016/j.asoc.2012.08.002>.
2. A. Q. Ansari, R.Biswas & S. Aggarwal. (Poster Presentation) Neutrosophication of Fuzzy Models, IEEE Workshop On Computational Intelligence: Theories, Applications and Future Directions (hostedby IIT Kanpur), 14th July'13.
3. A. Q. Ansari, R.Biswas & S. Aggarwal Extension to fuzzy logic representation: Moving towards neutrosophic logic - A new laboratory rat, Fuzzy Systems (FUZZ), 2013 IEEE International Conference, p.1 –8, (2013) DOI:10.1109/FUZZ-IEEE.2013.6622412.
4. A. Nagoor Gani and M.Basheer Ahamed, Order and Size in Fuzzy Graphs, Bulletin of Pure and Applied Sciences, Vol 22E (No.1), p.145-148 (2003).
5. A. Nagoor Gani. A and S. Shajitha Begum, Degree, Order and Size in Intuitionistic Fuzzy Graphs, International Journal of Algorithms, Computing and Mathematics,(3)3 (2010).
6. A. Nagoor Gani and S.R Latha, On Irregular Fuzzy Graphs, Applied Mathematical Sciences, Vol.6, no.11,517-523, (2012).
7. A. V. Devadoss, A. Rajkumar & N. J. P .Praveena, A Study on Miracles through Holy Bible using Neutrosophic Cognitive Maps (NCMS).International Journal of Computer Applications, 69(3) (2013).
8. A. Mohamed Ismayil and A. Mohamed Ali, On Strong Interval-Valued Intuitionistic Fuzzy Graph, International Journal of Fuzzy Mathematics and Systems. Volume 4, Number 2 pp. 161-

- 168 (2014).
9. D. Ezhilmaran & K. Sankar, Morphism of bipolar intuitionistic fuzzy graphs, *Journal of Discrete Mathematical Sciences and Cryptography*, 18:5, 605-621(2015), DOI:10.1080/09720529.2015.1013673.
  10. F. Smarandache. Refined Literal Indeterminacy and the Multiplication Law of Sub-Indeterminacies, *Neutrosophic Sets and Systems*, Vol. 9, 58-(2015),
  11. F. Smarandache, Types of Neutrosophic Graphs and neutrosophic Algebraic Structures together with their Applications in Technology, seminar, Universitatea Transilvania din Brasov, Facultatea de Design de Produs si Mediu, Brasov, Romania 06 June 2015.
  12. F. Smarandache, Neutrosophic set - a generalization of the intuitionistic fuzzy set, *Granular Computing*, 2006 IEEE International Conference, p. 38 – 42 (2006), DOI: 10.1109/GRC.2006.1635754.
  13. F. Smarandache, A geometric interpretation of the neutrosophic set — A generalization of the intuitionistic fuzzy set *Granular Computing (GrC)*, 2011 IEEE International Conference, p.602 – 606 (2011), DOI 10.1109/GRC.2011.6122665.
  14. F. Smarandache, *Symbolic Neutrosophic Theory*, Europanova asbl, Brussels, 195p.
  15. G. Garg, K. Bhutani, M. Kumar and S. Aggarwal, Hybrid model for medical diagnosis using Neutrosophic Cognitive Maps with Genetic Algorithms, *FUZZ-IEEE*, 6page 2015( IEEE International conference on fuzzy systems).
  16. H. Wang, Y. Zhang, R. Sunderraman, Truth-value based interval neutrosophic sets, *Granular Computing*, 2005 IEEE International Conference, vol. 1, p. 274 - 277 (2005), DOI: 10.1109/GRC.2005.1547284.
  17. H. Wang, F. Smarandache, Y. Zhang, and R. Sunderraman, Single valued Neutrosophic Sets, *Multispace and Multistructure* 4, p. 410-413 (2010).
  18. H. Wang, F. Smarandache, Zhang, Y.-Q. and R.Sunderraman, "Interval Neutrosophic Sets and Logic: Theory and Applications in Computing", *Hexis*, Phoenix, AZ, (2005).
  19. H. Y Zhang, J. Wang, X. Chen, An outranking approach for multi-criteria decision-making problems with interval-valued neutrosophic sets, *Neural Computing and Applications*, pp 1-13 (2015).
  20. H.Y. Zhang , P. Ji, J. Q.Wang & X. HChen, An Improved Weighted Correlation Coefficient Based on Integrated Weight for Interval Neutrosophic Sets and its Application in Multi-criteria Decision-making Problems, *International Journal of Computational Intelligence Systems* ,V8, Issue 6 (2015), DOI:10.1080/18756891.2015.1099917.
  21. H. D. Cheng and Y. Guo, A new neutrosophic approach to image thresholding, *New Mathematics and Natural Computation*, 4(3) (2008) 291–308.
  22. H.J. Zimmermann, *Fuzzy Set Theory and its Applications*, Kluwer-Nijhoff, Boston, 1985.
  23. I. Deli, M. Ali, F. Smarandache, Bipolar neutrosophic sets and their application based on multi-criteria decision making problems, *Advanced Mechatronic Systems (ICAMechS)*, 2015 International Conference, p.249 - 254 (2015), DOI: 10.1109/ICAMechS.2015.7287068.
  24. I. Turksen, Interval valued fuzzy sets based on normal forms, *Fuzzy Sets and Systems*, vol. 20, p. 191-210 (1986).
  25. J. Ye, vector similarity measures of simplified neutrosophic sets and their application in multicriteria decision making, *International Journal of Fuzzy Systems*, Vol. 16, No. 2, p.204-211 (2014).
  26. J. Ye, Single-Valued Neutrosophic Minimum Spanning Tree and Its Clustering Method, *Journal of Intelligent Systems* 23(3): 311–324,(2014).
  27. J. Ye, Trapezoidal neutrosophic set and its application to multiple attribute decision making, *Neural Computing and Applications*, (2014) DOI: 10.1007/s00521-014-1787-6.
  28. J. Chen, S. Li, S. Ma, and X. Wang, m-Polar Fuzzy Sets: An Extension of Bipolar Fuzzy Sets, *The Scientific World Journal*, (2014) <http://dx.doi.org/10.1155/2014/416530>.
  29. K. Atanassov, "Intuitionistic fuzzy sets," *Fuzzy Sets and Systems*, vol. 20, p. 87-96 (1986).
  30. K. Atanassov and G. Gargov, "Interval valued intuitionistic fuzzy sets," *Fuzzy Sets and Systems*, vol.31, pp. 343-349 (1989).
  31. K. Atanassov. *Intuitionistic fuzzy sets: theory and applications*. Physica, New York, 1999.

32. L. Zadeh Fuzzy sets, Inform and Control, 8, 338-353(1965)
33. M. Akram and W. A. Dudek, Interval-valued fuzzy graphs, Computers & Mathematics with Applications, vol. 61, no. 2, pp. 289–299 (2011).
34. M. Akram, Interval-valued fuzzy line graphs, Neural Computing and Applications, vol. 21, pp. 145–150 (2012).
35. M. Akram and B. Davvaz, Strong intuitionistic fuzzy graphs, Filomat, vol. 26, no. 1, pp. 177–196 (2012).
36. M. Akram, Bipolar fuzzy graphs, Information Sciences, vol. 181, no. 24, pp. 5548–5564 (2011).
37. M. Akram, Bipolar fuzzy graphs with applications, Knowledge Based Systems, vol. 39, pp. 1–8 (2013).
38. M. Akram, K. H.Dar, On N- graphs, Southeast Asian Bulletin of Mathematics, (2012)
39. Akram, M and A. Adeel, m- polar fuzzy graphs and m-polar fuzzy line graphs, Journal of Discrete Mathematical Sciences and Cryptography, 2015.
40. M. Akram, W. A. Dudek, Regular bipolar fuzzy graphs, Neural Computing and Applications, Vol 21, (2012)197-205.
41. M. Ali, and F. Smarandache, Complex Neutrosophic Set, Neural Computing and Applications, Vol. 25, (2016), 1-18. DOI: 10.1007/s00521-015-2154-y.
42. M. Ali, I. Deli, F. Smarandache, The Theory of Neutrosophic Cubic Sets and Their Applications in Pattern Recognition, Journal of Intelligent and Fuzzy Systems, (In press), 1-7, DOI:10.3233/IFS-151906.
43. P. Bhattacharya, Some remarks on fuzzy graphs, Pattern Recognition Letters 6: 297-302 (1987).
44. R. Parvathi and M. G. Karunambigai, Intuitionistic Fuzzy Graphs, Computational Intelligence, Theory and applications, International Conference in Germany, p.18 -20 (2006).
45. S.Broumi, M. Talea, F. Smarandache, Single Valued Neutrosophic Graphs: Degree, Order and Size, (2016) submitted
46. S.Broumi, M. Talea. A. Bakkali and F.Samarandache, Single Valued Neutrosophic Graph, Journal of New theory, N 10 (2016) 86-101.
47. S.Broumi, M. Talea, A.Bakali, F. Smarandache, Interval Valued Neutrosophic Graphs, critical review (2016) in presse
48. S.Broumi, M. Talea, A.Bakali, F. Smarandache, Operations on Interval Valued Neutrosophic Graphs, (2016) submitted.
49. S.Broumi, M. Talea, A.Bakali, F. Smarandache, On Strong Interval Valued Neutrosophic Graphs, (2016) submitted.
50. S. Broumi, F. Smarandache, New distance and similarity measures of interval neutrosophic sets, Information Fusion (FUSION), 2014 IEEE 17th International Conference, p. 1 – 7 (2014).
51. S. Aggarwal, R. Biswas, A. Q. Ansari, Neutrosophic modeling and control , Computer and Communication Technology (ICCCT), 2010 International Conference, p. 718 – 723 (2010) DOI:10.1109/ICCCT.2010.5640435.
52. S. N. Mishra and A. Pal, Product of Interval Valued Intuitionistic fuzzy graph, Annals of Pure and Applied Mathematics Vol. 5, No.1, 37-46 (2013).
53. Y. Hai-Long, G. She, Yanhonge , L. Xiuwu, On single valued neutrosophic relations, Journal of Intelligent & Fuzzy Systems, vol. Preprint, no. Preprint, p. 1-12 (2015).
54. W. B. Vasantha Kandasamy and F. Smarandache, Fuzzy Cognitive Maps and Neutrosophic Cognitive Maps, 2013.
55. W. B. Vasantha Kandasamy, K. Ilanthenral and Florentin Smarandache, Neutrosophic Graphs: A New Dimension to Graph Theory, Kindle Edition, 2015.
56. W.B. Vasantha Kandasamy and F. Smarandache “Analysis of social aspects of migrant laborers living with HIV/AIDS using Fuzzy Theory and Neutrosophic Cognitive Maps”, Xiquan, Phoenix (2004).
57. W.R. Zhang, "Bipolar fuzzy sets and relations: a computational framework for cognitive modeling and multiagent decision analysis," in Fuzzy Information Processing Society Biannual Conference, 1994. Industrial Fuzzy Control and Intelligent Systems Conference, and the NASA Joint Technology Workshop on Neural Networks and Fuzzy Logic(1994) pp.305-309, doi: 10.1109/IJCF.1994.375115