Neutrosophic Diagram and Classes of Neutrosophic Paradoxes or to the Outer-Limits of Science

Florentin Smarandache

Department of Mathematics, University of New Mexico, Gallup, NM 87301, USA. E-mail: smarand@unm.edu

These paradoxes are called "neutrosophic" since they are based on indeterminacy (or neutrality, i.e. neither true nor false), which is the third component in neutrosophic logic. We generalize the Venn diagram to a Neutrosophic Diagram, which deals with vague, inexact, ambiguous, illdefined ideas, statements, notions, entities with unclear borders. We define the neutrosophic truth table and introduce two neutrosophic operators (neuterization and antonymization operators) give many classes of neutrosophic paradoxes.

1 Introduction to the neutrosophics

Let $\langle A \rangle$ be an idea, or proposition, statement, attribute, theory, event, concept, entity, and $\langle non A \rangle$ what is not $\langle A \rangle$.

Let *<anti A>* be the opposite of *<A>*. We have introduced a new notation [1998], *<neut A>*, which is neither *<A>* nor *<anti A>* but in between. *<neut A>* is related with *<A>* and *<anti A>*.

Let's see an example for vague (not exact) concepts: if <*A*> is "*tall*" (an attribute), then <*anti A*> is "*short*", and <*neut A*> is "*medium*", while <*non A*> is "*not tall*" (which can be "*medium or short*"). Similarly for other <*A*>, <*neut A*>, <*anti A*> such as: <*good*>, <*so so*>, <*bad*>, or <*perfect*>, <*average*>, <*imperfect*>, or <*high*>, <*medium*>, <*small*>, or respectively <*possible*>, <*sometimes possible* and other times impossible>, <*impossible*>, etc.

Now, let's take an exact concept / statement: if $\langle A \rangle$ is the statement "1+1 = 2 *in base* 10", then $\langle anti A \rangle$ is "1+1 \neq 2 *in base* 10", while $\langle neut A \rangle$ is undefined (doesn't exist) since it is not possible to have a statement in between "1 + 1 = 2 *in base* 10" and "1 + 1 \neq 2 *in base* 10" because in base 10 we have 1+1 is either equal to 2 or 1+1 is different from 2. $\langle non A \rangle$ coincides with $\langle anti A \rangle$ in this case, $\langle non A \rangle$ is "1 + 1 \neq 2 *in base* 10".

Neutrosophy is a theory the author developed since 1995 as a generalization of dialectics. This theory considers every notion or idea $\langle A \rangle$ together with its opposite or negation $\langle anti A \rangle$, and the spectrum of "neutralities" in between them and related to them, noted by $\langle neut A \rangle$.

The Neutrosophy is a new branch of philosophy which studies the origin, nature, and scope of neutralities, as well as their interactions with different ideational spectra.

Its Fundamental Thesis:

Any idea $\langle A \rangle$ is T% true, I% indeterminate (i.e. neither true nor false, but neutral, unknown), and F% false.

Its Fundamental Theory:

Every idea $\langle A \rangle$ tends to be neutralized, diminished, balanced by $\langle non A \rangle$ ideas (not only by $\langle anti A \rangle$ as Hegel asserted) — as a state of equilibrium. In between $\langle A \rangle$ and $\langle anti A \rangle$ there may be a continuous spectrum of particular $\langle neut A \rangle$ ideas, which can balance $\langle A \rangle$ and $\langle anti A \rangle$.

To neuter an idea one must discover all its three sides: of sense (truth), of nonsense (falsity), and of undecidability (indeterminacy) — then reverse/combine them. Afterwards, the idea will be classified as neutrality.

There exists a *Principle of Attraction* not only between the opposites $\langle A \rangle$ and $\langle anti A \rangle$ (as in dialectics), but also between them and their neutralities $\langle neut A \rangle$ related to them, since $\langle neut A \rangle$ contributes to the *Completeness of Knowledge*.

Hence, neutrosophy is based not only on analysis of oppositional propositions as dialectic does, but on analysis of these contradictions together with the neutralities related to them.

Neutrosophy was extended to Neutrosophic Logic, Neutrosophic Set, Neutrosophic Probability and Neutrosophic Statistics, which are used in technical applications.

In the *Neutrosophic Logic* (which is a generalization of fuzzy logic, especially of intuitionistic fuzzy logic) every logical variable x is described by an ordered triple x = (T, I, F), where T is the degree of truth, F is the degree of falsehood, and I the degree of indeterminacy (or neutrality, i.e. neither true nor false, but vague, unknown, imprecise), with T, I, F standard or non-standard subsets of the non-standard unit interval]⁻⁰, 1⁺[. In addition, these values may vary over time, space, hidden parameters, etc.

Neutrosophic Probability (as a generalization of the classical probability and imprecise probability) studies the chance that a particular event $\langle A \rangle$ will occur, where that chance is represented by three coordinates (variables): T% chance the event will occur, I% indeterminate (unknown) chance, and F% chance the event will not occur.

Neutrosophic Statistics is the analysis of neutrosophic probabilistic events.

Neutrosophic Set (as a generalization of the fuzzy set, and especially of intuitionistic fuzzy set) is a set such that an element belongs to the set with a neutrosophic probability,

i.e. T degree of appurtenance (membership) to the set, I degree of indeterminacy (unknown if it is appurtenance or non-appurtenance to the set), and F degree of non-appurtenance (non-membership) to the set.

There exist, for each particular idea: PRO parameters, CONTRA parameters, and NEUTER parameters which influence the above values.

Indeterminacy results from any hazard which may occur, from unknown parameters, or from new arising conditions. This resulted from practice.

2 Applications of neutrosophics

Neutrosophic logic/set/probability/statistics are useful in artificial intelligence, neural networks, evolutionary programming, neutrosophic dynamic systems, and quantum mechanics.

3 Examples of neutrosophy used in Arabic philosophy (F. Smarandache and S. Osman)

• While Avicenna promotes the idea that the world is contingent if it is necessitated by its causes, Averroes rejects it, and both of them are right from their point of view.

Hence <*A*> and <*anti A*> have common parts.

• Islamic dialectical theology (*kalam*) promoting creationism was connected by Avicenna in an extraordinary way with the opposite Aristotelian-Neoplatonic tradition.

Actually a lot of work by Avicenna falls into the frame of neutrosophy.

- Averroes's religious judges (*qadis*) can be connected with atheists' believes.
- al-Farabi's metaphysics and general theory of emanation vs. al-Ghazali's Sufi writings and mystical treatises [we may think about a coherence of al-Ghazali's "Incoherence of the Incoherence" book].
- al-Kindi's combination of Koranic doctrines with Greek philosophy.
- Islamic Neoplatonism + Western Neoplatonism.
- Ibn-Khaldun's statements in his theory on the cyclic sequence of civilizations, says that: Luxury leads to the raising of civilization (because the

people seek for comforts of life) but also Luxury leads to the decay of civilization (because its correlation with the corruption of ethics).

- On the other hand, there's the method of absent-bypresent syllogism in jurisprudence, in which we find the same principles and laws of neutrosophy.
- In fact, we can also function a lot of Arabic aphorisms, maxims, Koranic miracles (Ayat Al- Qur'an) and Sunna of the prophet, to support the theory of neutrosophy.

Take the colloquial proverb that "The continuance of state is impossible" too, or "Everything, if it's increased over its extreme, it will turn over to its opposite"!

4 The Venn diagram

In a Venn diagram we have with respect to a universal set U the following:



Fig. 1: Venn diagram

Therefore, there are no common parts amongst $\langle A \rangle$, $\langle neut A \rangle$, and $\langle anti A \rangle$, and all three of them are (completely) contained by the universal set *U*. Also, all borders of these sets $\langle A \rangle$, $\langle neut A \rangle$, $\langle anti A \rangle$, and *U* are clear, exact. All these four sets are well–defined.

While $\langle neut A \rangle$ means neutralities related to $\langle A \rangle$ and $\langle anti A \rangle$, what is outside of $\langle A \rangle U \langle neut A \rangle U \langle anti A \rangle$ but inside of U are other neutralities, not related to $\langle A \rangle$ or to $\langle anti A \rangle$.

Given $\langle A \rangle$, there are two types of neutralities: those related to $\langle A \rangle$ (and implicitly related to $\langle anti A \rangle$), and those not related to $\langle A \rangle$ (and implicitly not related to $\langle anti A \rangle$)

5 The neutrosophic diagram, as extension of the Venn diagram

Yet, for ambiguous, vague, not-well-known (or even unknown) imprecise ideas / notions / statements / entities with unclear frontiers amongst them the below relationships may occur because between an approximate idea noted by $\langle A \rangle$ and its opposite *<anti A>* and their neutralities *<neut A>* there are not clear delimitations, not clear borders to distinguish amongst what is $\langle A \rangle$ and what is not $\langle A \rangle$. There are buffer zones in between <*A*> and <*anti A*> and <*neut A*>, and an element x from a buffer zone between $\langle A \rangle$ and <anti A> may or may not belong to both <A> and <anti A> simultaneously. And similarly for an element y in a buffer zone between $\langle A \rangle$ and $\langle neut A \rangle$, or an element z in the buffer zone between *<neut A>* and *<anti A>*. We may have a buffer zone where the confusion of appurtenance to $\langle A \rangle$, or to <neut A>, or to <anti A> is so high, that we can consider that an element w belongs to all of them simultaneously (or to none of them simultaneously).

We say that all four sets $\langle A \rangle$, $\langle neut A \rangle$, $\langle anti A \rangle$, and the neutrosophic universal set U are illdefined, inexact, unknown (especially if we deal with predictions; for example

Florentin Smarandache. Neutrosophic Diagram and Classes of Neutrosophic Paradoxes or to the Outer-Limits of Science

if $\langle A \rangle$ is a statement with some degree of chance of occurring, with another degree of change of not occurring, plus an unknown part). In the general case, none of the sets $\langle A \rangle$, $\langle neut A \rangle$, $\langle anti A \rangle$, $\langle non A \rangle$ are completely included in U, and neither U is completely known; for example, if U is the neutrosophic universal set of some specific given events, what about an unexpected event that might belong to U? That's why an approximate U (with vague borders) leaves room for expecting the unexpected.

The *Neutrosophic Diagram* in the general case is the following (Fig. 2): the borders of <*A*>, *<anti A*>, and *<neut A*> are dotted since they are unclear.



Fig. 2: Neutrosophic Diagram

Similarly, the border of the neutrosophic universal set U is dotted, meaning also unclear, so U may not completely contain $\langle A \rangle$, nor $\langle neut A \rangle$ or $\langle anti A \rangle$, but U "approximately" contains each of them. Therefore, there are elements in $\langle A \rangle$ that may not belong to U, and the same thing for $\langle neut A \rangle$ and $\langle anti A \rangle$. Or elements, in the most ambiguous case, there may be elements in $\langle A \rangle$ and in $\langle neut A \rangle$ and in $\langle anti A \rangle$ which are not contained in the universal set U.

Even the neutrosophic universal set is ambiguous, vague, and with unclear borders.

Of course, the intersections amongst $\langle A \rangle$, $\langle neut A \rangle$, $\langle anti A \rangle$, and U may be smaller or bigger or even empty depending on each particular case.

See below an example of a particular neutrosophic diagram (Fig. 3), when some intersections are contained by the neutrosophic universal set:



Fig. 3: Example of a particular neutrosophic diagram

A neutrosophic diagram is different from a Venn diagram since the borders in a neutrosophic diagram are vague. When all borders are exact and all intersections among $\langle A \rangle$, $\langle neut A \rangle$, and $\langle anti A \rangle$ are empty, and all $\langle A \rangle$, $\langle neut A \rangle$, and $\langle anti A \rangle$ are included in the neutrosophic universal set U, then the neutrosophic diagram becomes a Venn diagram.

The neutrosophic diagram, which complies with the neutrosophic logic and neutrosophic set, is an extension of the Venn diagram.

6 Classes of neutrosophic paradoxes

The below classes of neutrosophic paradoxes are not simply word puzzles. They may look absurd or unreal from the classical logic and classical set theory perspective. If $\langle A \rangle$ is a precise / exact idea, with well-defined borders that delimit it from others, then of course the below relationships do not occur.

But let $\langle A \rangle$ be a vague, imprecise, ambiguous, not-wellknown, not-clear-boundary entity, $\langle non A \rangle$ means what is not $\langle A \rangle$, and $\langle anti A \rangle$ means the opposite of $\langle A \rangle$. $\langle neut A \rangle$ means the neutralities related to $\langle A \rangle$ and $\langle anti A \rangle$, neutralities which are in between them.

When $\langle A \rangle$, $\langle neut A \rangle$, $\langle anti A \rangle$, $\langle non A \rangle$, U are uncertain, imprecise, they may be selfcontradictory. Also, there are cases when the distinction between a set and its elements is not clear.

Although these neutrosophic paradoxes are based on "pathological sets" (those whose properties are considered atypically counterintuitive), they are not referring to the theory of Meinongian objects (*Gegenstandstheorie*) such as round squares, unicorns, etc. Neutrosophic paradoxes are not reported to objects, but to vague, imprecise, unclear ideas or predictions or approximate notions or attributes from our everyday life.

7 Neutrosophic operators

Let's introduce for the first time two new Neutrosophic Operators:

An operator that "neuterizes" an idea. To neuterize [neuter+ize, transitive verb; from the Latin word neuter = neutral, in neither side], n(.), means to map an entity to its neutral part. [We use the Segoe Print for "n(.)".] "To neuterize" is different from "to neutralize" [from the French word neutraliser] which means to declare a territory neutral in war, or to make ineffective an enemy, or to destroy an enemy.

 $n(\langle A \rangle) = \langle neut A \rangle$. By definition $n(\langle neut A \rangle) = \langle neut A \rangle$.

For example, if $\langle A \rangle$ is "tall", then n(tall) = medium, also n(short) = medium, n(medium) = medium. But if $\langle A \rangle$ is "1 + 1 = 2 *in base* 10" then $n(\langle 1 + 1 = 2 in base 10 \rangle)$ is undefined (does not exist), and similarly $n(\langle 1 + 1 \neq 2 in base 10 \rangle)$ is undefined. And an operator that "antonymizes" an idea. To antonymize [antonym+ize, transitive verb; from the Greek work antōnymia = instead of, opposite], α(.), means to map an entity to its opposite. [We use the Segoe Print for α(.)] α(<A>) = <anti A>.

For example, if $\langle A \rangle$ is "*tall*", then $\alpha(tall) = short$, also $\alpha(short) = tall$, and $\alpha(medium) = tall or short$. But if $\langle A \rangle$ is "1 + 1 = 2 *in base*10" then $\alpha(\langle 1 + 1 = 2 in base10\rangle) = \langle 1+1 \neq 2 in base$ 10> and reciprocally $\alpha(\langle 1+1 \neq 2 in base$ 10>) = $\langle 1+1 = 2 in base$ 10>.

The classical operator for *negation / complement* in logics respectively in set theory, "*to negate*" (\neg), which is equivalent in neutrosophy with the operator "*to nonize*" (i.e. to *non+ize*) or *nonization* (i.e. *non+ization*), means to map an idea to its neutral or to its opposite (a union of the previous two neutrosophic operators: *neuterization and antonymization*):

 $\neg <A> = < non A> = < neut A> \cup < anti A> = n(<A>) \cup a(<A>).$

Neutrosophic Paradoxes result from the following *neutrosophic logic / set connectives* following all apparently impossibilities or semi-impossibilities of neutrosophically connecting $\langle A \rangle$, $\langle anti A \rangle$, $\langle neut A \rangle$, $\langle non A \rangle$, and the neutrosophic universal set U.

8 Neutrosophic truth tables

For <*A*> = "*tall*":

< <i>A</i> >	𝔅(<a>)	n(<a>)	$\neg $
tall	short	medium	short or medium
medium	short or tall	medium	short or tall
short	tall	medium	tall or medium

To remark that $n(< medium>) \triangleq medium$. If <A> = tall, then < neut A> = medium, and < neut(neut A)> = < neut A>, or n(< n(<A>)>) = n(<A>).

For $\langle A \rangle =$ "1 + 1 = 2 *in base* 10" we have $\langle anti A \rangle = \langle non A \rangle =$ " $1 + 1 \neq 2$ *in base* 10", while $\langle neut A \rangle$ is undefined (N/A) — whence the neutrosophic truth table becomes:

< <i>A</i> >	<i>a</i> (< <i>A</i> >)	n (<a>)	$\neg $
True	False	N/A	False
False	True	N/A	True

In the case when a statement is given by its neutrosophic logic components $\langle A \rangle = (T, I, F)$, i.e. $\langle A \rangle$ is T% true, I% indeterminate, and F% false, then the neutrosophic truth table depends on the defined neutrosophic operators for each application.

9 Neutrosophic operators and classes of neutrosophic paradoxes

a) Complement/Negation

 $\neg <A > \neq <non A >$ and reciprocally $\neg <non A > \neq <A >$.

 $\neg(\neg < A >) \neq < A >$

 $\neg(\neg < anti A >) \neq < anti A >$ $\neg(\neg < non A >) \neq < non A >$

 $\neg(\neg < neut A >) \neq < neut A >$ $\neg(\neg U) \neq U$, where U is the neutrosophic universal

set. $\neg(\neg < \emptyset >) \neq < \emptyset >$, where $< \emptyset >$ is the neutrosophic empty set.

b) Neuterization

 $n(<A>) \neq <neut A>$ $n(<anti A>) \neq <neut A>$ $n(<non A>) \neq <neut A>$ $n(n(<A>)) \neq <A>$

c) Antonymization

 $a(<A>) \neq <anti A>$ $a(<anti A>) \neq <A>$ $a(<non A>) \neq <A>$ $a(a(<A>)) \neq <A>$

d) Intersection/Conjunction

 $\langle A \rangle \cap \langle non A \rangle \neq \emptyset$ (neutrosophic empty set) [symbolically $(\exists x)(x \in A \land x \in \neg A)$],

or even more $\langle A \rangle \cap \langle anti A \rangle \neq \emptyset$ [symbolically $(\exists x)$ $(x \in A \land x \in a(A))$],

similarly $\langle A \rangle \cap \langle neut | A \rangle \neq \emptyset$ and $\langle anti | A \rangle \cap \langle neut | A \rangle \neq \emptyset$, $\neq \emptyset$,

up to $\langle A \rangle \cap \langle neut A \rangle \cap \langle anti A \rangle \neq \emptyset$.

The symbolic notations will be in a similar way.

This is *Neutrosophic Transdisciplinarity*, which means to find common features to uncommon entities.

For examples:

There are things which are good and bad in the same time.

There are things which are good and bad and medium in the same time (because from one point of view they may be god, from other point of view they may be bad, and from a third point of view they may be medium).

e) Union / Weak Disjunction

 $<A> \cup <neut A> \cup <anti A> \neq U.$ $<anti A> \cup <neut A> \neq <non A>.$ Etc.

f) Inclusion/Conditional

 $\langle A \rangle \subset \langle anti A \rangle$ $(\forall x)(x \in A \rightarrow x \in a(A))$ All is $\langle anti A \rangle$, the $\langle A \rangle$ too. All *good* things are also *bad*. All is *imperfect*, the *perfect* too.

 $\langle anti A \rangle \subset \langle A \rangle$ $(\forall x)(x \in a(A) \rightarrow x \in A)$ All is $\langle A \rangle$, the $\langle anti A \rangle$ too. All bad things have something good in them [this is rather a fuzzy paradox]. All is *perfect* things are *imperfect* in some degree.

 $< non A > \subset <A >$ $(\forall x)(x \in \neg A \rightarrow x \in A)$ All is $\langle A \rangle$, the $\langle non A \rangle$ too. All bad things have something good and something *medium* in them [this is a neutrosophic paradox, since it is based on good, bad, and medium]. All is perfect things have some imperfectness and mediocrity in them at some degree. $<A> \subset <neut A>$ $(\forall x)(x \in A \rightarrow x \in \mathbf{n}(A))$ All is $\langle neutA \rangle$, the $\langle A \rangle$ too. <*non* $A > \subset <$ *neut*A > [partial neutrosophic paradox of inclusion] $(\forall x)(x \in \neg A \rightarrow x \in \mathbf{n}(A))$ All is < neutA >, the < non A > too. $< non A > \subset < antiA > [partial neutrosophic paradox of$ inclusion] $(\forall x)(x \in \neg A \to x \in \pmb{a}(A))$ All is *<antiA>*, the *<non A>* too. $\langle antiA \rangle \subset \langle neut A \rangle$ $(\forall \mathbf{x})(\mathbf{x} \in \boldsymbol{a}(A) \rightarrow \mathbf{x} \in \boldsymbol{n}(A))$ All is *<neut A>*, the *<anti A>* too. $<\!\!A\!\!> \cup <\!\!anti A\!\!> \subset <\!\!neut A\!\!>$ $(\forall \mathbf{x})((x \in \mathbf{A} \lor \mathbf{x} \in \boldsymbol{a}(\mathbf{A})) \to \mathbf{x} \in \boldsymbol{n}(\mathbf{A}))$ All is *<neutA>*, the *<A>* and *<antiA>* too.

Paradoxes of some Neutrosophic Arguments $\langle A \rangle \Rightarrow \langle B \rangle$ $\langle B \rangle \Rightarrow \langle anti A \rangle$ $\therefore \langle A \rangle \Rightarrow \langle anti A \rangle$ Example: too much work produces sickness; sickness

produces less work (absences from work, low efficiency); therefore, too much work implies less work (this is a *Law of Self-Equilibrium*).

 $<\!\!A \!\!> \Rightarrow <\!\!B \!\!> \\<\!\!B \!\!> \Rightarrow <\!\!non A \!\!> \\\therefore <\!\!A \!\!> \Rightarrow <\!\!non A \!\!> \\<\!\!A \!\!> \Rightarrow <\!\!R \!\!>$

 $<\!\!B\!\!> \Rightarrow <\!\!neut A\!\!>$ $\therefore <\!\!A\!\!> \Rightarrow <\!\!neut A\!\!>$

g) Equality/Biconditional

Unequal Equalities $\langle A \rangle \neq \langle A \rangle$ which symbolically becomes $(\exists x)(x \in \neg A \leftrightarrow x \notin \neg A)$ or even stronger inequality $(\forall x)(x \in \neg A \leftrightarrow x \notin \neg A)$. Nothing is $\langle A \rangle$, nor even $\langle A \rangle$. $\langle anti A \rangle \neq \langle anti A \rangle$ which symbolically becomes $(\exists x)(x \in A \leftrightarrow x \notin A)$

or even stronger inequality $(\forall x)(x \in A \leftrightarrow x \notin A)$.

<neut $A > \neq$ <neut A >which symbolically becomes $(\exists x)(x \in vA \leftrightarrow x \notin vA)$ or even stronger inequality $(\forall x)(x \in vA \leftrightarrow x \notin vA)$. $< non A > \neq < non A >$

which symbolically becomes $(\exists x)(x \in \neg A \leftrightarrow x \notin \neg A)$ or even stronger inequality $(\forall x)(x \in \neg A \leftrightarrow x \notin \neg A)$.

Equal Inequalities $\langle A \rangle = \langle anti A \rangle$ $(\forall x)(x \in A \leftrightarrow x \in a(A))$ All is $\langle A \rangle$, the $\langle anti A \rangle$ too; and reciprocally, all is $\langle anti A \rangle$, the $\langle A \rangle$ too. Or, both combined implications give: All is $\langle A \rangle$ is equivalent to all is $\langle anti A \rangle$.

And so on: <*A*> = <*neut A*> <*anti A*> = <*neut A*> <*non A*> = <*A*>

Dilations and Absorptions <anti A> = <non A>, which means that <anti A> is dilated to its neutrosophic superset <non A>, or <non A> is absorbed to its neutrosophic subset <anti A>. Similarly for: <neut A> = <non A>

<A> = U<neut A> = U<anti A> = U

 $\langle non A \rangle = U$

 h) Combinations of the previous single neutrosophic operator equalities and/or inequalities, resulting in more neutrosophic operators involved in the same expression.

For examples:

<*neut* $A > \cap (<A > \cup <$ *anti* $A >) \neq \emptyset$ [two neutrosophic operators].

 $<A>\cup$ < anti $A> \neq \neg$ < neut A> and reciprocally \neg $(<A>\cup$ < anti $A>) \neq$ < neut A>.

 $\langle A \rangle \cup \langle neut A \rangle \neq \neg \langle anti A \rangle$ and reciprocally.

 \neg (*<A*> \cup *<neut A*> \cup *<anti A*>) $\neq \emptyset$ and reciprocally. Etc.

- i) We can also take into consideration other logical connectors, such as strong disjunction (we previously used the *weak disjunction*), *Shaffer's connector*, *Peirce's connector*, and extend them to the neutrosophic form.
- j) We may substitute *<A>* by some entities, attributes, statements, ideas and get nice neutrosophic paradoxes, but not all substitutions will work properly.

10 Some particular paradoxes

Quantum Semi-Paradox

Let's go back to 1931 Schrödinger's paper. Saul Youssef writes (flipping a *quantum coin*) in arXiv.org at quant-ph/ 9509004:

It is interesting. Actually we can say the same for any probability different from 1: If at the beginning, the probability of a quantum event, P(quantum event) = p, with 0 , and if later the event occurs, we get to <math>P(quantum event) = 1; but if the event does not occur, then we get P(quantum event) = 0, so still a kind of contradiction.

Torture's paradox

An innocent person P, who is tortured, would say to the torturer T whatever the torturer wants to hear, even if P doesn't know anything.

So, T would receive incorrect information that will work against him/her. Thus, the torture returns against the torturer.

Paradoxist psychological behavior

Instead of being afraid of something, say $\langle A \rangle$, try to be afraid of its opposite $\langle anti A \rangle$, and thus– because of your fear – you'll end up with the $\langle anti \langle anti \rangle$, which is $\langle A \rangle$.

Paradoxically, *negative publicity* attracts better than positive one (enemies of those who do negative publicity against you will sympathize with you and become your friends).

Paradoxistically [word coming etymologically from *para-doxism*, *paradoxist*], to be in opposition is more poetical and interesting than being opportunistic.

At a sportive, literary, or scientific competition, or in a war, to be on the side of the weaker is more challenging but on the edge of chaos and, as in Complex Adoptive System, more potential to higher creation.

Law of Self-Equilibrium

(Already cited above at the Neutrosophic Inclusion/Conditional Paradoxes) $\langle A \rangle \rightarrow \langle B \rangle$ and $\langle B \rangle \rightarrow \langle anti A \rangle$, therefore $\langle A \rangle \rightarrow \langle anti A \rangle$!

Example: too much work produces sickness; sickness produces less work (absences from work, low efficiency); therefore, too much work implies less work.

Submitted on April 25, 2010 / Accepted on April 30, 2010

References

- Weisstein E.W. Smarandache paradox. CRC Concise Enciclopedia of Mathematics, CRC Press, Boca Raton, FL, 1998, 1661.
- Begay A. The Smarandache Semantic Paradox. *Humanistic* Mathematics Network Journal, Harvey Mudd College, Claremont (CA), 1998, no. 17, 48.
- Greenstein C.H. Dictionary of logical terms and symbols. Van Nostrand Reinhold Co., 1978.
- Devaraj Ramasamy. Florentin Smarandache set up the paradoxist literary movement. In *Parnassus of World Poets 1994*, Madras, India, September 1994.

- 5. Dale J. Logic: the semantics of existence and nonexistence. Berlin, de Gruyter, 1996.
- Le C.T. The Smarandache class of paradoxes. *Bulletin of the Transylvania University of Brasov*, New Series, Series B, 1994, v.1(36), 7–8.
- 7. Le C.T. The Smarandache class of paradoxes. *Bulletin of Pure and Applied Sciences E*, 1995, v.14(2), 109–110.
- Le C.T. The most paradoxist mathematician of the world: Florentin Smarandache. *Bulletin of Pure and Applied Sciences E*, 1996, v.15(1), 81–100.
- 9. Le C.T. The Smarandache class of paradoxes. *Journal of Indian Academy of Mathematics*, 1996, v.18, no.1, 53–55.
- Le C.T. The Smarandache class of paradoxes (mathematical poem). In: Bunner H.C. An anthology in memoriam. Bristol Banner Books, Bristol (IN), 1996, 94.
- 11. Le C.T. Clasa de paradoxuri Smarandache. *Tempus*, 1994, anul III, no.2(5), 4.
- 12. Mitroiescu I. The Smarandache class of paradoxes applied in computer sciences. *Abstracts of Papers Presented to the American Mathematical Society*, 1995, v.16, no.3, issue 101, 651.
- 13. Mitroiescu I. The Smarandache class of paradoxes. *The Mathematical Gazette*, 1995, v.79, no.484, 125.
- Popescu M. A model of the Smarandache paradoxist geometry. *Abstracts of Papers Presented to the American Mathematical Society*, 1996, v.17, no.1, issue 103, 265.
- 15. Smarandache F. Neutrosophy. Neutrosophic probability, set, and logic. American Research Press, Rehoboth (NM), 1998; Republished in 2000, 2003, 2005 as Smarandache F. A unifying field in logics: neutrosophic logic. Neutrosophy, neutrosophic set, neutrosophic probability and statistics. American Research Press, Rehoboth (NM).
- Smarandache F. Mixed non-Euclidean geometries. Arhivele Statului, Filiala Vâlcea, Rm. Vâlcea, 1969.
- 17. Smarandache F., Osman S. Neutrosophy in Arabic philosophy. Renaissance High Press (Ann Arbor), 2007.
- Smarandache F. Mathematical fancies and paradoxes. *The Eugene Strens Memorial on Intuitive and Recreational Mathematics and its History*, University of Calgary, Alberta, Canada, 27 July 2 August, 1986.
- Tilton H.B. Smarandache's paradoxes. *Math Power*, Tucson (AZ), 1996, v.2, no.9, 1–2.

Florentin Smarandache. Neutrosophic Diagram and Classes of Neutrosophic Paradoxes or to the Outer-Limits of Science