



International Journal of Social Economics

A family of estimators of population mean using multiauxiliary information in presence of measurement errors

Jack Allen, Housila P. Singh, Florentin Smarandache,

Article information:

To cite this document:

Jack Allen, Housila P. Singh, Florentin Smarandache, (2003) "A family of estimators of population mean using multiauxiliary information in presence of measurement errors", International Journal of Social Economics, Vol. 30 Issue: 7, pp.837-848, <https://doi.org/10.1108/03068290310478775>

Permanent link to this document:

<https://doi.org/10.1108/03068290310478775>

Downloaded on: 27 April 2018, At: 08:13 (PT)

References: this document contains references to 30 other documents.

To copy this document: permissions@emeraldinsight.com

The fulltext of this document has been downloaded 388 times since 2006*

Access to this document was granted through an Emerald subscription provided by

For Authors

If you would like to write for this, or any other Emerald publication, then please use our Emerald for Authors service information about how to choose which publication to write for and submission guidelines are available for all. Please visit www.emeraldinsight.com/authors for more information.

About Emerald www.emeraldinsight.com

Emerald is a global publisher linking research and practice to the benefit of society. The company manages a portfolio of more than 290 journals and over 2,350 books and book series volumes, as well as providing an extensive range of online products and additional customer resources and services.

Emerald is both COUNTER 4 and TRANSFER compliant. The organization is a partner of the Committee on Publication Ethics (COPE) and also works with Portico and the LOCKSS initiative for digital archive preservation.

*Related content and download information correct at time of download.



A family of estimators of population mean using multiauxiliary information in presence of measurement errors

A family of estimators

837

Jack Allen

School of Accounting and Finance, Griffith University, Gold Coast, Australia

Housila P. Singh

School of Statistics, Vikram University, Ujjain, India

Florentin Smarandache

Department of Mathematics, University of New Mexico, Gallup, New Mexico, USA

Keywords *Population, Averaging, Information, Measurement, Error estimation*

Abstract *This paper proposes a family of estimators of population mean using information on several auxiliary variables and analyzes its properties in the presence of measurement errors.*

Introduction

The discrepancies between the values exactly obtained on the variables under consideration for sampled units and the corresponding true values are termed as measurement errors. In general, standard theory of survey sampling assumes that data collected through surveys are often assumed to be free of measurement or response errors. In reality such a supposition does not hold true and the data may be contaminated with measurement errors due to various reasons (see, for example Cochran (1963) and Sukhatme *et al.* (1984)).

One of the major sources of measurement errors in survey is the nature of variables. This may happen in case of qualitative variables. Simple examples of such variables are intelligence, preference, specific abilities, utility, aggressiveness, tastes, etc. In many sample surveys it is recognized that errors of measurement can also arise from the person being interviewed, from the interviewer, from the supervisor or leader of a team of interviewers, and from the processor who transmits the information from the recorded interview on to the punched cards or tapes that will be analyzed, for instance, see Cochran (1968). Another source of measurement error is when the variable is conceptually well defined but observations can be obtained on some closely related substitutes termed as proxies or surrogates. Such a situation is



encountered when one needs to measure the economic status or the level of education of individuals, see Salabh (1997) and Sud and Srivastava (2000). In the presence of measurement errors, inferences may be misleading, see Biermer *et al.* (1991), Fuller (1995) and Manisha and Singh (2001).

There is today a great deal of research on measurement errors in surveys. An attempt has been made to study the impact of measurement errors on a family of estimators of population mean using multiauxiliary information.

The suggested family of estimators

Let Y be the study variate and its population mean μ_0 to be estimated using information on $p(> 1)$ auxiliary variates X_1, X_2, \dots, X_p . Further, let the population mean row vector $\underline{\mu}' = (\mu_1, \mu_2, \dots, \mu_p)$ of the vector $\underline{X}' = (X_1, X_2, X_p)$. Assume that a simple random sample of size n is drawn from a population, on the study character Y and auxiliary characters X_1, X_2, \dots, X_p . For the sake of simplicity we assume that the population is infinite. The recorded fallible measurements are given by:

$$y_j = Y_j + E_j$$

$$x_{ij} = X_{ij} + \eta_{ij}, i = 1, 2, \dots, p,$$

$$j = 1, 2, \dots, n.$$

where Y_j and X_{ij} are correct values of the characteristics Y and $X_i (i = 1, 2, \dots, p; j = 1, 2, \dots, n)$.

For the sake of simplicity in exposition, we assume that the error E_j s are stochastic with mean "zero" and variance $\sigma_{(0)}^2$ and uncorrelated with Y_j s. The errors η_{ij} in x_{ij} are distributed independently of each other and of the X_{ij} with mean "zero" and variance $\sigma_{(i)}^2 (i = 1, 2, \dots, p)$. Also E_j s and η_{ij} s are uncorrelated although Y_j s and X_{ij} s are correlated.

Define

$$u_i = \frac{\bar{X}_i}{\mu_i}, (i = 1, 2, \dots, p),$$

$$u^T = (u_1, u_2, \dots, u_p)_{1 \times p},$$

$$e^T = (1, 1, \dots, 1)_{1 \times p},$$

$$\bar{y} = \frac{1}{n} \sum_{j=1}^n y_j,$$

$$\bar{x}_i = \frac{1}{n} \sum_{j=1}^n x_{ij}.$$

With this background we suggest a family of estimators of μ_0 as:

$$\hat{\mu}_g = g(\bar{y}, u^T). \tag{2.1}$$

where $g(\bar{y}, u^T)$ is a function of $\bar{y}, u_1, u_2, \dots, u_p$ such that:

$$\begin{aligned} g(u_0, e^T) &= \mu_0 \\ \Rightarrow \frac{\partial g(\cdot)}{\partial \bar{y}} \Big|_{(\mu_0, e^T)} &= 1 \end{aligned}$$

and such that it satisfies the following conditions:

- the function $g(\bar{y}, u^T)$ is continuous and bounded in Q ;
- the first and second order partial derivatives of the function $g(\bar{y}, u^T)$ exist and are continuous and bounded in Q .

To obtain the mean squared error of $\hat{\mu}_g$, we expand the function $g(\bar{y}, u^T)$ about the point (μ_0, e^T) in a second order Taylor's series. We get:

$$\begin{aligned} \hat{\mu}_g &= g(\mu_0, e^T) + (\bar{y} - \mu_0) \frac{\partial g(\cdot)}{\partial \bar{y}} \Big|_{(\mu_0, e^T)} + (u - e)^T g^{(1)}(\mu_0, e^T) \tag{2.2} \\ &+ \frac{1}{2} \left\{ (\bar{y} - \mu_0)^2 \frac{\partial^2 g(\cdot)}{\partial \bar{y}^2} \Big|_{(\bar{y}^*, u^{*T})} + 2(\bar{y} - \mu_0)(u - e)^T \frac{\partial g^{(1)}(\cdot)}{\partial \bar{y}} \Big|_{(\bar{y}^*, u^{*T})} \right. \\ &\left. + (u - e)^T g^{(2)}(\bar{y}^*, u^{*T})(u - e) \right\}, \end{aligned}$$

where:

$$\bar{y}^* = \mu_0 + \theta(\bar{y} - \mu_0), u^* = e + \theta(u - e), 0 < \theta < 1; g^{(1)}(\cdot)$$

denote the p element column vector of first partial derivatives of $g(\cdot)$ and $g^{(2)}(\cdot)$ denotes a $p \times p$ matrix of second partial derivatives of $g(\cdot)$ with respect to u .

Noting that $g(\mu_0, e^T) = \mu_0$, it can be shown that:

$$E(\hat{\mu}_g) = \mu_0 + O(n^{-1}), \tag{2.3}$$

which follows that the bias of $\hat{\mu}_g$ is of the order of n^{-1} , and hence its contribution to the mean squared error of $\hat{\mu}_g$ will be of the order of n^{-2} . From (2.2), we have to terms of order n^{-1} :

$$\begin{aligned}
 \text{MSE}(\hat{\mu}_g) &= E\{(\bar{y} - \mu_0) + (u - e)^T g^{(1)}(\mu_0, e^T)\}^2 \\
 &= E\left[(\bar{y} - \mu_0)^2 + 2(\bar{y} - \mu_0)(u - e)^T g^{(1)}(\mu_0, e^T) \right. \\
 &\quad \left. + (g^{(1)}(\mu_0, e^T))^T (u - e)(u - e)^T (g^{(1)}(\mu_0, e^T))\right] \\
 &= \frac{1}{n} \left[\mu_0^2 (C_0^2 + C_{(0)}^2) + 2\mu_0 b^T g^{(1)}(\mu_0, e^T) + (g^{(1)})^T A (g^{(1)}(\mu_0, e^T)) \right]
 \end{aligned}
 \tag{2.4}$$

where $b^T = (b_1, b_2, \dots, b_p)$, $b_i = \rho_{0i} C_0 C_i$, ($i = 1, 2, \dots, p$); $C_i = \sigma_i / \mu_i$, $C_{(i)} = \sigma_i / \mu_i$, ($i = 1, 2, \dots, p$) and $C_0 = \sigma_0 / \mu_0$:

$$A = \begin{bmatrix} C_1^2 + C_{(1)}^2 & \rho_{12} C_1 C_2 & \rho_{13} C_1 C_3 & \cdots & \rho_{1p} C_1 C_p \\ \rho_{12} C_1 C_2 & C_2^2 + C_{(2)}^2 & \rho_{23} C_2 C_3 & \cdots & \rho_{2p} C_2 C_p \\ \rho_{13} C_1 C_3 & \rho_{23} C_2 C_3 & C_3^2 + C_{(3)}^2 & \cdots & \rho_{3p} C_3 C_p \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \rho_{1p} C_1 C_p & \rho_{2p} C_2 C_p & \rho_{3p} C_3 C_p & \cdots & C_p^2 + C_{(p)}^2 \end{bmatrix}_{p \times p}$$

The MSE($\hat{\mu}_g$) at equation (2.4) is minimized for

$$g^{(1)}(\mu_0, e^T) = -\mu_0 A^{-1} b.
 \tag{2.5}$$

Thus the resulting minimum MSE of $\hat{\mu}_g$ is given by:

$$\min. \text{MSE}(\hat{\mu}_g) = (\mu_0^2/n) \left[C_0^2 + C_{(0)}^2 - b^T A^{-1} b \right].
 \tag{2.6}$$

Now we have established the following theorem.

Theorem 2.1=up to terms of order n^{-1} ,

$$\text{MSE}(\hat{\mu}_g) \geq (\mu_0^2/n) \left[C_0^2 + C_{(0)}^2 - b^T A^{-1} b \right]
 \tag{2.7}$$

with equality holding if:

$$g^{(1)} = -\mu_0 A^{-1} b.$$

It is to be mentioned that the family of estimators $\hat{\mu}_g$ at equation (2.1) is very large. The following estimators may be identified as particular members of the suggested family of estimators $\hat{\mu}_g$:

- $\hat{\mu}_g^{(1)} = \bar{y} \sum_{i=1}^p \omega_i \left(\frac{\mu_i}{\bar{x}_i} \right); \sum_{i=1}^p \omega_i = 1$, (Olkin, 1958).

- $\hat{\mu}_g^{(2)} = \bar{y} \sum_{i=1}^p \omega_i \left(\frac{\bar{x}_i}{\mu_i} \right), \sum_{i=1}^p \omega_i = 1$, (Singh, 1967).
- $\hat{\mu}_g^{(3)} = \bar{y} \frac{\sum_{i=1}^p \omega_i \mu_i}{\sum_{i=1}^p \omega_i \bar{x}_i}, \sum_{i=1}^p \omega_i = 1$, (Shukla, 1966; John, 1969).
- $\hat{\mu}_g^{(4)} = \bar{y} \frac{\sum_{i=1}^p \omega_i \bar{x}_i}{\sum_{i=1}^p \omega_i \mu_i}; \sum_{i=1}^p \omega_i = 1$, (Sahai *et al.*, 1980).
- $\hat{\mu}_g^{(5)} = \bar{y} \prod_{i=1}^p \left(\frac{\mu_i}{\bar{x}_i} \right)^{\omega_i}, \sum_{i=1}^p \omega_i = 1$, (Mohanty and Pattanaik, 1984).
- $\hat{\mu}_g^{(6)} = \bar{y} \left(\sum_{i=1}^p \frac{\omega_i \bar{x}_i}{\mu_i} \right)^{-1}, \sum_{i=1}^p \omega_i = 1$, (Mohanty and Pattanaik, 1984).
- $\hat{\mu}_g^{(7)} = \bar{y} \prod_{i=1}^p \left(\frac{\bar{x}_i}{\mu_i} \right)^{\omega_i}, \sum_{i=1}^p \omega_i = 1$, (Tuteja and Bahl, 1991).
- $\hat{\mu}_g^{(8)} = \bar{y} \left[\sum_{i=1}^p \frac{\omega_i \mu_i}{\bar{x}_i} \right]^{-1}, \sum_{i=1}^p \omega_i = 1$, (Tuteja and Bahl, 1991).
- $\hat{\mu}_g^{(9)} = \bar{y} \left[\omega_{p+1} + \sum_{i=1}^p \omega_i \left(\frac{\mu_i}{\bar{x}_i} \right) \right], \sum_{i=1}^{p+1} \omega_i = 1$.
- $\hat{\mu}_g^{(10)} = \bar{y} \left[\omega_{p+1} + \sum_{i=1}^p \omega_i \left(\frac{\bar{x}_i}{\mu_i} \right) \right], \sum_{i=1}^{p+1} \omega_i = 1$.
- $\hat{\mu}_g^{(11)} = \bar{y} \left[\sum_{i=1}^q \omega_i \left(\frac{\mu_i}{x_i} \right) + \sum_{i=q+1}^p \left(\frac{\hat{x}_i}{\mu_i} \right) \right]; \left(\sum_{i=1}^q \omega_i + \sum_{i=q+1}^p \omega_i \right)^{=1}$;
(Srivastava, 1965; Rao and Mudholkar, 1967).
- $\hat{\mu}_g^{(12)} = \bar{y} \prod_{i=1}^p \left(\frac{\bar{x}_i}{\mu_i} \right)^{\alpha_i}$ (α_i s are suitably constants) (Srivastava, 1967).

- $\hat{\mu}_g^{(13)} = \bar{y} \prod_{i=1}^p \left\{ 2 - \left(\frac{\bar{x}_i}{\mu_i} \right)^{\alpha_i} \right\}$ (Sahai and Ray, 1980).
- $\hat{\mu}_g^{(14)} = \bar{y} \prod_{i=1}^p \frac{\bar{x}_i}{\{\mu_i + \alpha_i(\bar{x}_i - \mu_i)\}}$ (Walsh, 1970).
- $\hat{\mu}_g^{(15)} = \bar{y} \exp \left\{ \sum_{i=1}^p \theta_i \log u_i \right\}$ (Srivastava, 1971).
- $\hat{\mu}_g^{(16)} = \bar{y} \exp \left\{ \sum_{i=1}^p \theta_i (u_i - 1) \right\}$ (Srivastava, 1971).
- $\hat{\mu}_g^{(17)} = \bar{y} \sum_{i=1}^p \omega_i \exp \left\{ (\theta_i / \omega_i) \log u_i \right\}; \sum_{i=1}^p \omega_i = 1$, (Srivastava, 1971).
- $\hat{\mu}_g^{(18)} = \bar{y} + \sum_{i=1}^p \alpha_i (\bar{x}_i - \mu_i)$, etc.

The MSE of these estimators can be obtained from equation (2.4).
It is well known that:

$$V(\bar{y}) = (\mu_0^2/n) (C_0^2 + C_{(0)}^2). \tag{2.8}$$

It follows from equation (2.6) and equation (2.8) that the minimum variance of $\hat{\mu}_g$ is no longer than conventional unbiased estimator \bar{y} .

On substituting $\sigma_{(0)}^2 = 0, \sigma_{(i)}^2 = 0 \forall i = 1, 2, \dots, p$ in equation (2.4), we obtain the no-measurement error case. In that case, the MSE of $\hat{\mu}_g$, is given by:

$$\begin{aligned} \text{MSE}(\hat{\mu}_g) &= \frac{1}{n} \left[C_0^2 \mu_0^2 + 2\mu_0 b^T g *^{(1)} (\mu_0, e^T) + (g *^{(1)} (\mu_0, e^T))^T A * (g *^{(1)}) \right] \\ &= \text{MSE}(\hat{\mu}_g *), \end{aligned} \tag{2.9}$$

where:

$$\begin{aligned} \hat{\mu}_g &= g * \left(\bar{Y}, \frac{\bar{X}_1}{\mu_1}, \frac{\bar{X}_2}{\mu_2}, \dots, \frac{\bar{X}_p}{\mu_p} \right) \\ &= g * (\bar{Y}, U^T) \end{aligned} \tag{2.10}$$

and \bar{Y} and $\bar{X}_i (i = 1, 2, \dots, p)$ are the sample means of the characteristics Y and X_i based on true measurements. $(Y_j, X_{ij}, i = 1, 2, \dots, p; j = 1, 2, \dots, n)$. The family of estimators $\hat{\mu}_g *$ at equation (2.10) is a generalized version of Srivastava (1971, p. 80).

The MSE of $\hat{\mu}_g^*$ is minimized for:

$$g^*(\mu_0, e^T) = -A^{-1} b \mu_0. \quad (2.11)$$

A family of estimators

Thus the resulting minimum MSE of $\hat{\mu}_g^*$ is given by:

$$\min.\text{MSE}(\hat{\mu}_g^*) = \frac{\mu_0}{n} [C_0^2 - b^T A^{-1} b] \quad (2.12)$$

843

$$= \frac{\sigma_0}{n} (1 - R^2),$$

where $A^* = [a^*_{ij}]$ is a $p \times p$ matrix with $a^*_{ij} = \rho_{ij} C_i C_j$ and R stands for the multiple correlation coefficient of Y on X_1, X_2, \dots, X_p .

From equations (2.6) and (2.12) the increase in minimum MSE($\hat{\mu}_g$) due to measurement errors is obtained as:

$$\begin{aligned} \min.\text{MSE}(\hat{\mu}_g) - \min.\text{MSE}(\hat{\mu}_g^*) &= \left(\frac{\mu_0^2}{n}\right) [C_{(0)}^2 + b^T A^{-1} b - b^T A^{-1} b] \\ &> 0. \end{aligned}$$

This is due to the fact that the measurement errors introduce the variances fallible measurements of study variate Y and auxiliary variates X_i . Hence there is a need to take the contribution of measurement errors into account.

Biases and mean square errors of some particular estimators in the presence of measurement errors

To obtain the bias of the estimator $\hat{\mu}_g$, we further assume that the third partial derivatives of $g(\bar{y}, u^T)$ also exist and are continuous and bounded. Then expanding $g(\bar{y}, u^T)$ about the point $(\bar{y}, u^T) = (\mu_0, e^T)$ in a third-order Taylor's series we obtain:

$$\begin{aligned} \hat{\mu}_g &= g(\mu_0, e^T) + (\bar{y} - \mu_0) \frac{\partial g(\cdot)}{\partial \bar{y}} \Big|_{(\mu_0, e^T)} + (u - e)^T g^{(1)}(\mu_0, e^T) \\ &+ \frac{1}{2} \{ (\bar{y} - \mu_0) \frac{\partial^2 g(\cdot)}{\partial \bar{y}^2} \Big|_{(\mu_0, e^T)} + 2(\bar{y} - \mu_0)(u - e)^T g^{(1)}(\mu_0, e^T) \\ &+ (u - e)^T (g^{(2)}(\mu_0, e^T))(u - e) \} \\ &+ \frac{1}{6} \{ (\bar{y} - \mu_0) \frac{\partial}{\partial \bar{y}} + (u - e) \frac{\partial}{\partial u} \}^3 g(\bar{y}^*, u^{*T}), \end{aligned} \quad (3.1)$$

where $g^{(12)}(\mu_0, e^T)$ denotes the matrix of second partial derivatives of $g(\bar{y}, u^T)$ at the point $(\bar{y}, u^T) = (\mu_0, e^T)$.

Noting that:

$$\begin{aligned}
 g(u_0 e^T) &= \mu_0 \\
 \left. \frac{\partial g(\cdot)}{\partial \bar{y}} \right|_{(\mu_0, e^T)} &= 1 \\
 \left. \frac{\partial^2 g(\cdot)}{\partial \bar{y}^2} \right|_{(\mu_0, e^T)} &= 0,
 \end{aligned}$$

and taking expectation we obtain the bias of the family of estimators $\hat{\mu}_g$ to the first degree of approximation:

$$B(\hat{\mu}_g) = \frac{1}{2} \left[E\{(u - e)^T (g^{(2)}(\mu_0, e^T))(u - e)\} + 2\left(\frac{\mu_0}{n}\right) b^T g^{(12)}(\mu_0, e^T) \right] \tag{3.2}$$

where $b^T = (b_1, b_2, \dots, b_p)$ with $b_i = \rho_{0i} C_0 C_i$; ($i = 1, 2, \dots, p$). Thus we see that the bias of $\hat{\mu}_g$ depends also on the second order partial derivatives of the function on $g(\bar{y}, u^T)$ at the point (μ_0, e^T) , and hence will be different for different optimum estimators of the family.

The biases and mean square errors of the estimators $\hat{\mu}_{g_i}$; $i = 1$ to 18 up to terms of order n^{-1} along with the values of $g^{(1)}(\mu_0, e^T)$, $g^{(2)}(\mu_0, e^T)$ and $g^{(12)}(\mu_0, e^T)$ are given in the Table I.

Estimators based on estimated optimum

It may be noted that the minimum MSE, equation (2.6), is obtained only when the optimum values of constants involved in the estimator, which are functions of the unknown population parameters μ_0 , b and A , are known quite accurately.

To use such estimators in practice, one has to use some guessed values of the parameters μ_0 , b and A , either through past experience or through a pilot sample survey. Das and Tripathi (1978, sec. 3) have illustrated that even if the values of the parameters used in the estimator are not exactly equal to their optimum values as given by equation (2.5) but are close enough, the resulting estimator will be better than the conventional unbiased estimator \bar{y} . For further discussion on this issue, the reader is referred to Murthy (1967), Reddy (1973), Srivenkataramana and Tracy (1984) and Sahai and Sahai (1985).

On the other hand if the experimenter is unable to guess the values of population parameters due to lack of experience, it is advisable to replace the unknown population parameters by their consistent estimators. Let $\hat{\phi}$ be a consistent estimator of $\phi = A^{-1}b$. We then replace ϕ by $\hat{\phi}$ and also μ_0 by \bar{y} if necessary, in the optimum $\hat{\mu}_g$ resulting in the estimator $\hat{\mu}_{g(est)}$, say, which will now be a function of \bar{y} , u and ϕ . Thus we define a family of estimators (based on estimated optimum values) of μ_0 as:

Estimator	$g^{(1)}(\mu_0, eT)$	$g^{(2)}(\mu_0, eT)$	$g^{(12)}(\mu_0, eT)$	Bias	MSE
$\hat{\mu}_g^{(1)}$	$-\mu_0 \omega$	$2\mu_0 \tilde{W}$ $p \times p$	$-\omega$	$\left(\frac{\mu_0}{n}\right) \left(C^T \tilde{W} - b^T \tilde{\omega} \right)$ $p \times p$	$\left(\frac{\mu_0}{n}\right) \left[C_0^2 + C_{(0)}^2 - 2b^T \tilde{\omega} + \omega^T A \omega \right]$
$\hat{\mu}_g^{(2)}$	$\mu_0 \omega$	Q (null matrix) $p \times p$	ω	$\left(\frac{\mu_0}{n}\right) b^T \tilde{\omega}$	$\left(\frac{\mu_0^2}{n}\right) \left[C_0^2 + C_{(0)}^2 + 2b^T \tilde{\omega} + \omega^T A \omega \right]$
$\hat{\mu}_g^{(3)}$	$\frac{\mu_0 \omega^*}{-\omega^T \underline{\mu}}$	$\frac{2\mu_0 \omega^* \omega^{*T}}{\omega^T \underline{\mu} \underline{\mu}^T \omega}$	$-\frac{\omega^*}{\omega^T \underline{\mu}}$	$\left(\frac{\mu_0}{n}\right) \left(\frac{\omega^{*T} A \omega}{\omega^T \underline{\mu} \underline{\mu}^T \omega} - \frac{b^T \tilde{\omega}^*}{\omega^T \underline{\mu}} \right)$	$\left(\frac{\mu_0^2}{n}\right) \left[C_0^2 + C_{(0)}^2 - \frac{2b^T \tilde{\omega}^*}{\omega^T \underline{\mu}} + \frac{\omega^{*T} A \omega^*}{\omega^T \underline{\mu} \underline{\mu}^T \omega} \right]$
$\hat{\mu}_g^{(4)}$	$\frac{\mu_0 \omega}{\omega^T \underline{\mu}}$	Q (null matrix) $p \times p$	$\frac{\omega}{\omega^T \underline{\mu}}$	$\left(\frac{\mu_0}{n}\right) \frac{b^T \tilde{\omega}}{\omega^T \underline{\mu}}$	$\left(\frac{\mu_0^2}{n}\right) \left[C_0^2 + C_{(0)}^2 + \frac{2b^T \tilde{\omega}}{\omega^T \underline{\mu}} + \frac{\omega^T A \omega}{\omega^T \underline{\mu} \underline{\mu}^T \omega} \right]$
$\hat{\mu}_g^{(5)}$	$-\mu_0 \omega$	$\mu_0 \left(\omega \omega^T + \tilde{W} \right)$ $p \times p$	$-\omega$	$\left(\frac{\mu_0}{2n}\right) \left[\omega^T A \omega + C^T \tilde{W} - 2b^T \tilde{\omega} \right]$ $p \times p$	$\left(\frac{\mu_0}{n}\right) \left[C_0^2 + C_{(0)}^2 - 2b^T \tilde{\omega} + \omega^T A \omega \right]$
$\hat{\mu}_g^{(6)}$	$-\mu_0 \omega$	$2\mu_0 \omega^T \omega$	$-\omega$	$\left(\frac{\mu_0}{n}\right) \left[\omega^T A \omega - b^T \tilde{\omega} \right]$	$\left(\frac{\mu_0}{n}\right) \left[C_0^2 + C_{(0)}^2 - 2b^T \tilde{\omega} + \omega^T A \omega \right]$
$\hat{\mu}_g^{(7)}$	$\mu_0 \omega$	$\mu_0 \left(\omega \omega^T - \tilde{W} \right)$ $p \times p$	ω	$\left(\frac{\mu_0}{2n}\right) \left[\omega^T A \omega - C^T \tilde{W} + 2b^T \tilde{\omega} \right]$ $p \times p$	$\left(\frac{\mu_0}{n}\right) \left[C_0^2 + C_{(0)}^2 + 2b^T \tilde{\omega} + \omega^T A \omega \right]$
$\hat{\mu}_g^{(8)}$	$\mu_0 \omega$	$2\mu_0 \left(\omega \omega^T - \tilde{W} \right)$ $p \times p$	ω	$\left(\frac{\mu_0}{n}\right) \left[\omega^T A \omega - C^T \tilde{W} + b^T \tilde{\omega} \right]$ $p \times p$	$\left(\frac{\mu_0}{n}\right) \left[C_0^2 + C_{(0)}^2 + 2b^T \tilde{\omega} + \omega^T A \omega \right]$

(continued)

where $Wp \times p = \text{diag}(\omega_1, \omega_2, \dots, \omega_p)$ where $C^T = (C_1^2 + C_{(1)}^2, C_2^2 + C_{(2)}^2, \dots, C_p^2 + C_{(p)}^2)$

where $\omega^{*T} = (\omega_1^*, \omega_2^*, \dots, \omega_p^*)$ with $(\omega_i^*, * = \omega_i \mu)(i = 1, 2, \dots, p)$

Table I. Biases and mean squared errors of various estimators of μ_0

Estimator	$g^{(1)}(\mu_0, eT)$	$g^{(2)}(\mu_0, eT)$	$g^{(12)}(\mu_0, eT)$	Bias	MSE
$\hat{\mu}_g^{(6)}$	$-\mu_0 \omega$	$2\mu_0 W_{p \times p}$	$-\omega$	$\left(\frac{h_n}{n}\right) \left(C^T W_{p \times p} - b^T \omega \right)$	$\left(\frac{h_n^2}{n}\right) \left[C_0^2 + C_{(0)}^2 + 2b^T \omega + \omega^T A \omega \right]$
$\hat{\mu}_g^{(10)}$	$\mu_0 \omega$	Q	ω	$\left(\frac{h_n}{n}\right) b^T \omega$	$\left(\frac{h_n^2}{n}\right) \left[C_0^2 + C_{(0)}^2 + 2b^T \omega + \omega^T A \omega \right]$
$\hat{\mu}_g^{(11)}$	$\omega_{(1)} \mu_0$	$2W_{(1)} \mu_0$	$\omega_{(1)}$	$\left(\frac{h_n}{n}\right) \left(C *^T W_{(1)} - b^T \omega_{(1)} \right)$	$\left(\frac{h_n^2}{n}\right) \left[C_0^2 + C_{(0)}^2 - 2b^T \omega_{(1)} + \omega_{(1)}^T A \omega_{(1)} \right]$
where $\omega_{(1)} = (-\omega_1, -\omega_2, \dots, -\omega_q, -\omega_{q+1}, \dots, \omega_p)_{1 \times p}$					
$W_{(1) \times p} = \begin{bmatrix} \omega_1 & 0 & 0 & \dots & 0 & 0 & \dots & 0 \\ 0 & \omega_2 & 0 & \dots & 0 & 0 & \dots & 0 \\ 0 & 0 & \omega & \dots & 0 & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & \omega_q & 0 & \dots & 0 \\ 0 & 0 & 0 & \dots & 0 & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 0 & 0 & \dots & 0 \end{bmatrix}_{p \times p}$					
$C *^T = (C12 + C(1)2, \dots, Cq2 + C(q)2, \dots, 0)$					
$\hat{\mu}_g^{(12)}$	$\bar{\alpha} \mu_0$	$\mu_0 \left(\bar{\alpha} \bar{\alpha}^T - \bar{\alpha} \right)_{p \times p}$	$-\bar{\alpha}$	$\left(\frac{h_n}{2n}\right) \left[\bar{\alpha}^T A \bar{\alpha} - C^T \bar{\alpha} + 2b^T \bar{\alpha} \right]_{p \times p}$	$\left(\frac{h_n}{n}\right) \left[C_0^2 + C_{(0)}^2 + 2b^T \bar{\alpha} + \bar{\alpha}^T A \bar{\alpha} \right]$
where $\bar{\alpha}^T = (\alpha_1, \alpha_2, \dots, \alpha_p)_{1 \times p}$, $\bar{\alpha} = \text{diag}(\alpha_1, \alpha_2, \dots, \alpha_p)$					
$\hat{\mu}_g^{(13)}$	$-\bar{\alpha} \mu_0$	$-\mu_0 \left(\bar{\alpha} \bar{\alpha}^T - \bar{\alpha} \right)_{p \times p}$	$-\bar{\alpha}$	$\left(\frac{h_n}{2n}\right) \left[C^T \bar{\alpha} - \bar{\alpha}^T A \bar{\alpha} - 2b^T \bar{\alpha} \right]_{p \times p}$	$\left(\frac{h_n}{n}\right) \left[C_0^2 + C_{(0)}^2 - 2b^T \bar{\alpha} + \bar{\alpha}^T A \bar{\alpha} \right]$

(continued)

Estimator	$g^{(1)}(\mu_0, eT)$	$g^{(2)}(\mu_0, eT)$	$g^{(12)}(\mu_0, eT)$	Bias	MSE
$\hat{\mu}_g^{(14)}$	$-\underline{\alpha}\mu_0$	$2\mu_0\underline{\alpha}\underline{\alpha}$	$-\underline{\alpha}$	$\left(\frac{\mu_0}{n}\right) \left[\underline{\alpha}^T A \underline{\alpha} - C^T - b^T \underline{\alpha} \right]$	$\left(\frac{\mu_0}{n}\right) \left[C_0^2 + C_{(0)}^2 - 2b^T \underline{\alpha} + \underline{\alpha}^T A \underline{\alpha} \right]$
$\hat{\mu}_g^{(15)}$	$\mu_0 \underline{\theta}$	$\mu_0 \left(\underline{\theta} \underline{\theta}^T - \underline{\Theta} \right)_{p \times p}$	$\underline{\theta}$	$\left(\frac{\mu_0}{n}\right) \left[\underline{\theta}^T A \underline{\theta} - C^T \underline{\Theta} + 2b^T \underline{\theta} \right]_{p \times p}$	$\left(\frac{\mu_0^2}{n}\right) \left[C_0^2 + C_{(0)}^2 + 2b^T \underline{\theta} + \underline{\theta}^T A \underline{\theta} \right]$
$\hat{\mu}_g$	$\mu_0 \underline{\theta}$	where $\underline{\Theta} = \text{diag}\{\theta_1, \theta_2, \dots, \theta_p\}$	$\underline{\theta}$	$\left(\frac{\mu_0}{n}\right) \left[\underline{\theta}^T A \underline{\theta} + 2b^T \underline{\theta} \right]$	$\left(\frac{\mu_0^2}{n}\right) \left[C_0^2 + C_{(0)}^2 + 2b^T \underline{\theta} + \underline{\theta}^T A \underline{\theta} \right]$
$\hat{\mu}_g$	$\mu_0 \underline{\theta}$	$\underline{\Theta} *_{p \times p} \mu_0$	$\underline{\theta}$	$\left(\frac{\mu_0}{n}\right) \left[C^T \underline{\Theta} *_{p \times p} + 2b^T \underline{\theta} \right]$	$\left(\frac{\mu_0}{n}\right) \left[C_0^2 + C_{(0)}^2 + 2b^T \underline{\theta} + \underline{\theta}^T A \underline{\theta} \right]$
$\hat{\mu}_g$	$\underline{\alpha}^*$	$\underline{Q}_{p \times p}$	where $\underline{\Theta} *_{p \times p} = \text{diag}\left\{\theta_1 \left(\frac{\theta_1}{\alpha_1} - 1\right), \dots, \theta_p \left(\frac{\theta_p}{\alpha_p} - 1\right)\right\}$	Unbiased	$\left(\frac{1}{n}\right) \left[C_0^2 + C_{(0)}^2 + 2\mu_0 b^T \underline{\alpha}^* + \underline{\alpha}^*{}^T A \underline{\alpha}^* \right]$
			where $\underline{\alpha}^*{}^T = (\alpha_{1*}, \alpha_{2*}, \dots, \alpha_{p*})$ with $\alpha_{*i} = (\alpha_i, \mu_i, i = 1, 2, \dots, p)$		

Table I.

$$\hat{\mu}_{g(est)} = g * * (\bar{y}, u^T, \hat{\phi}^T), \quad (4.1)$$

where $g * *(\cdot)$ is a function of $(\bar{y}, u^T, \hat{\phi}^T)$ such that:

$$g * *(\mu_0, e^T, \phi^T) = \mu_0 \text{ for all } \mu_0, \Rightarrow \left. \frac{\partial g * *(\cdot)}{\partial \bar{y}} \right|_{(\mu_0, e^T, \phi^T)} = 1$$

$$\left. \frac{\partial g * *(\cdot)}{\partial u} \right|_{(\mu_0, e^T, \phi^T)} = \left. \frac{\partial g(\cdot)}{\partial u} \right|_{(\mu_0, e)^T} = -\mu_0 A^{-1} b = -\mu_0 \phi, \quad (4.2)$$

and:

$$\left. \frac{\partial g * *(\cdot)}{\partial \hat{\phi}} \right|_{(\mu_0, e^T, \phi^T)} = 0.$$

With these conditions and following Srivastava and Jhaji (1983), it can be shown to the first degree of approximation that:

$$\text{MSE}(\hat{\mu}_{g(est)}) = \min. \text{MSE}(\hat{\mu}_g) = \left(\frac{\mu_0}{n} \right) [C_0^2 + C_{(0)}^2 - b^T A^{-1} b].$$

Thus if the optimum values of constants involved in the estimator are replaced by their consistent estimators and conditions (4.2) hold true, the resulting estimator $\hat{\mu}_{g(est)}$ will have the same asymptotic mean square error, as that of optimum $\hat{\mu}_g$. Our work needs to be extended and future research will explore the computational aspects of the proposed algorithm.

References

- Biermer, P.P., Groves, R.M., Lyberg, L.E., Mathiowetz, N.A. and Sudman, S. (1991), *Measurement Errors in Surveys*, Wiley, New York, NY.
- Cochran, W.G. (1963), *Sampling Techniques*, John Wiley, New York, NY.
- Cochran, W.G. (1968), "Errors of measurement in statistics", *Technometrics*, Vol. 10 No. 4, pp. 637-66.
- Das, A.K. and Tripathi, T.P. (1978), "Use of auxiliary information in estimating the finite population variance", *Sankhya*, Vol. C No. 40, pp. 139-48.
- Fuller, W.A. (1995), "Estimation in the presence of measurement error", *International Statistical Review*, Vol. 63 No. 2, pp. 121-47.
- John, S. (1969), "On multivariate ratio and product estimators", *Biometrika*, pp. 533-6.
- Manisha and Singh, R.K. (2001), "An estimation of population mean in the presence of measurement errors", *Jour. Ind. Soc. Agri. Statist.*, Vol. 54 No. 1, pp. 13-18.
- Mohanty, S. and Pattanaik, L.M. (1984), "Alternative multivariate ratio estimators using geometric and harmonic means", *Jour. Ind. Soc. Agri. Statist.*, Vol. 36, pp. 110-18.
- Murthy, M.N. (1967), *Sampling Theory and Methods*, Statistical Publishing Society, Calcutta.
- Olkin, I. (1958), "Multivariate ratio estimation for finite population", *Biometrika*, Vol. 45, pp. 154-65.

- Rao, P.S.R.S. and Mudholkar, G.S. (1967), "Generalized multivariate estimators for the mean of a finite population", *Jour. Amer. Statist. Assoc.*, Vol. 62, pp. 1009-12.
- Reddy, V.N. (1973), "On ratio and product methods of estimation", *Sankhya*, Vol. B No. 35, pp. 307-16.
- Sahai, A. and Ray, S.K. (1980), "An efficient estimator using auxiliary information", *Metrika*, Vol. 27, pp. 271-5.
- Sahai, A. and Sahai, A. (1985), "On efficient use of auxiliary information", *Jour. Statist. Plann. Inference*, Vol. 12, pp. 203-12.
- Sahai, A., Chander, R. and Mathur, A.K. (1980), "An alternative multivariate product estimator", *Jour. Ind. Soc. Agril. Statist.*, Vol. 32 No. 2, pp. 6-12.
- Salabh (1997), "Ratio method of estimation in the presence of measurement error", *Jour. Ind. Soc. Agri. Statist.*, Vol. 52, pp. 150-5.
- Shukla, G.K. (1966), "An alternative multivariate ratio estimate for finite population", *Cal. Statist. Assoc. Bull.*, Vol. 15, pp. 127-34.
- Singh, M.P. (1967), "Multivariate product method of estimation for finite population", *Jour. Ind. Soc. Agri. Statist.*, Vol. 19 No. 2, pp. 1-10.
- Srivastava, S.K. (1965), "An estimator of the mean of a finite population using several auxiliary characters", *Jour. Ind. Statist. Assoc.*, Vol. 3, pp. 189-94.
- Srivastava, S.K. (1967), "An estimator using auxiliary information in sample surveys", *Cal. Statist. Assoc. Bull.*, Vol. 16, pp. 121-32.
- Srivastava, S.K. (1971), "A generalized estimator for the mean of a finite population using multiauxiliary information", *Jour. Amer. Statist. Assoc.*, Vol. 66, pp. 404-7.
- Srivastava, S.K. and Jhaji, H.S. (1983), "A class of estimators of the population mean using multi-auxiliary information", *Cal. Statist. Assoc. Bull.*, Vol. 32, pp. 47-56.
- Srivenkataramana, T. and Tracy, D.S. (1984), "Positive and negative valued auxiliary variates in surveys", *Metron*, Vol. 30 No. 3-4, pp. 3-13.
- Sud, U.C. and Srivastava, S.K. (2000), "Estimation of population mean in repeat surveys in the presence of measurement errors", *Jour. Ind. Soc. Ag. Statist.*, Vol. 53 No. 2, pp. 125-33.
- Sukhatme, P.V., Sukhatme, B.V., Sukhatme, S. and Ashok, C. (1984), *Sampling Theory of Surveys with Applications*, Iowa State University Press, Ames, IA.
- Tuteja, R.K. and Bahl, S. (1991), "Multivariate product estimators", *Cal. Statist. Assoc. Bull.*, Vol. 42, pp. 109-15.
- Walsh, J.E. (1970), "Generalization of ratio estimate for population total", *Sankhya*, Vol. A No. 32, pp. 99-106.

Further reading

- Reddy, V.N. and Rao, T.J. (1977), "Modified PPS method of estimation", *Sankhya*, Vol. C No. 39, pp. 185-97.
- Srivastava, S.K. (1980), "A class of estimators using auxiliary information in sample surveys", *Canad. Jour. Statist.*, Vol. 8, pp. 253-4.
- Tankou, V. and Dharmadlikari, S. (1989), "Improvement of ratio-type estimators", *Biom. Jour.*, Vol. 31 No. 7, pp. 795-802.

This article has been cited by:

1. Sadia Khalil, Sat Gupta, Muhammad Hanif. 2018. A generalized estimator for finite population mean in the presence of measurement errors in stratified random sampling. *Journal of Statistical Theory and Practice* **12**:2, 311-324. [[Crossref](#)]
2. Sadia Khalil, Muhammad Noor-ul-Amin, Muhammad Hanif. 2017. Generalized estimator of population mean by using conventional and non-conventional measures in the presence of measurement errors. *Communications in Statistics - Simulation and Computation* **31**, 1-14. [[Crossref](#)]
3. Sunil Kumar. 2016. Improved estimation of population mean in presence of nonresponse and measurement error. *Journal of Statistical Theory and Practice* **10**:4, 707-720. [[Crossref](#)]
4. Sunil Kumar, Sandeep Bhogal, N. S. Nataraja, M. Viswanathaiah. 2015. Estimation of Population Mean in the Presence of Non-Response and Measurement Error. *Revista Colombiana de Estadística* **38**:1, 145-161. [[Crossref](#)]
5. Giancarlo Diana, Marco Giordan. 2012. Finite Population Variance Estimation in Presence of Measurement Errors. *Communications in Statistics - Theory and Methods* **41**:23, 4302-4314. [[Crossref](#)]
6. 2009. A General Procedure for Estimating the General Parameter Using Auxiliary Information in Presence of Measurement Errors. *Communications of the Korean statistical society* **16**:5, 821-840. [[Crossref](#)]
7. Housila P. Singh, Namrata Karpe. 2009. A Class of Estimators Using Auxiliary Information for Estimating Finite Population Variance in Presence of Measurement Errors. *Communications in Statistics - Theory and Methods* **38**:5, 734-741. [[Crossref](#)]