

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/339787289>

Neutrosophic-Cubic Analytic Hierarchy Process with Applications

Article · April 2020

CITATIONS

0

READS

3

3 authors, including:



Muhammad Gulistan
Hazara University

71 PUBLICATIONS 339 CITATIONS

[SEE PROFILE](#)



Ismat Beg
Lahore School of Economics

282 PUBLICATIONS 3,091 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Fixed Point Theory for Set Valued Maps [View project](#)



RECENT ADVANCES IN MATHEMATICAL METHODS, MODELS & APPLICATIONS [View project](#)

NEUTROSOPHIC-CUBIC ANALYTIC HIERARCHY PROCESS WITH APPLICATIONS

MUHAMMAD GULISTAN, ISMAT BEG, AND MUHAMMAD ASIF

ABSTRACT. In this paper we extend fuzzy analytic hierarchy process into neutrosophic cubic environment. The neutrosophic cubic analytic hierarchy process can be used to manage more complex problems when the decision makers has a number of uncertainty, assigning preferences values to the considered object. We also define the concept of triangular neutrosophic cubic numbers and their operations laws. The advantages of the proposed methodology and the application of neutrosophic cubic analytic hierarchy process in decision making are shown by testing the numerical example in practical life.

THEORY OF APPROXIMATION AND APPLICATIONS (in Press)
http://msj.iau-arak.ac.ir/article_671487.html

1. Introduction

Analytic hierarchy process is initially developed by Saaty [13] and it held a very important place in the field of operation research while selecting the best alternatives. Analytic hierarchy process (AHP) is a multi criterion technique which is used to solve and analyze complex problems. AHP is also used to make mathematical and programming devices to select those alternatives [14]. AHP has various steps; In the first step, the problems are structured in order for clear understanding. This order is based on a particular pattern, it consists of the goal, decision making criteria, sub-criteria and in the last all accessible alternatives. When the hierarchy is structured, the decision makers construct pair wise comparison matrices. The scale that is used to measure criteria is called Satty's scale [15]. On the basis of measurement, alternative is first determined and then ranked. The AHP can predict both qualitative and quantitative elements. This quality is widely used in multi-criteria decision making technique. In practical life, decision criterion is a habitually hazy, difficult and conflicting. Along with that, there is uncertainty and using non-fuzzy value in a decision matrix sometimes gets inaccurate. The other thing is that the information available to the decision makers is indeterminant, see paper [16]. This is also a factor, in the way of accuracy. Many researchers started to use a new theory called the fuzzy set theory [23]. However, there were some drawbacks in that theory that it only considers truth membership degree. First time, Van Laarhoven and Pedrycz [19] introduced fuzzy AHP, in which the membership is taken in terms of triangular fuzzy number and used a logarithmic least squares method to obtain the fuzzy weight and fuzzy concert scores for ranking

1991 *Mathematics Subject Classification.* 91B06; 90B50; 91B10; 90C29; 62C86; 46S40.

Key words and phrases. Analytic hierarchy process; decision making; neutrosophic cubic set; consistency test.

the alternatives. Through this, they were able to control the indeterminacy, inconsistency and inaccuracy. Then Buckley [2] extended the classical AHP with the trapezoidal fuzzy number and gets the fuzzy weight and fuzzy concert scores in the geometric mean method. After that, Chang [4] used row mean method to derived priority for similarity ratio in the perspective of triangular fuzzy numbers. The Chang method is comparatively easier than the other fuzzy AHP approach. Even though the computation of fuzzy AHP is tiresome it is appropriate to confine and represent the human assessment of haziness when the multifarious multi-criteria decision making is considered. It has been practical as a depiction of the fuzzy AHP too many special areas, for instance public administration [8], airlines industry [12], manufacturing industry [6], textile industry [3], electronic industry [5], oil industry [7], entertainment industry [22], transportation industry [11] etc. Xu and Liao [20], provided the concept of intuitionistic fuzzy analytic hierarchy process, which is a new approach in AHP. Cubic set was developed by Jun in 2012 [9]. In 1999 Smarandache [17] define the neutrosophic set (NS). In neutrosophic set (NS) Smarandache add indeterminacy-membership function, i.e NS is composed of truth-membership $T(u)$, indeterminacy-membership $I(u)$ and falsity-membership $F(u)$, also see [18, 21]. Jun et al [10], projected the new concept of neutrosophic cubic set, which have three grade of membership true, false and indeterminacy. Basset [1], takes the fuzzy analytic hierarchy process into the neutrosophic analytic hierarchy process. In continuation of these efforts in this work we develop an algorithm for neutrosophic-cubic analytic hierarchy process for solving complicated multi-criteria decision making problems.

The arrangement of remaining paper is as follows: Section 2: review of basic concepts and properties. Section 3: we present the triangular neutrosophic-cubic numbers and operation rules. Section 4: present the algorithm for neutrosophic-cubic analytic hierarchy process. Section 5: as an application we use this in a practical example of the daily life problem. Section 6: we compare our method to other existing methods and finally we write the conclusion of this paper in section 7.

2. PRELIMINARIES

This section consists of two parts: Some notations with their descriptions and some previous definitions and results.

2.1. Notations. This section consists of some notations with their descriptions, as shown in Table 1.

Table [1]. Some notations with their descriptions.

Notation	Description
X	Ground set
x	Element of ground set (X)
Υ	Fuzzy set
$\tilde{\Upsilon} = [\Upsilon^L, \Upsilon^U]$	Interval valued fuzzy set
T_{NC}, I_{NC}, F_{NC}	Components of neutrosophic set
$\tilde{T}_{NC}, \tilde{I}_{NC}, \tilde{F}_{NC}$	Components of interval valued neutrosophic set
$\tilde{T}_{NC}, \tilde{I}_{NC}, \tilde{F}_{NC}, T_{NC}, I_{NC}, F_{NC}$	Components of neutrosophic cubic set

2.2. Pre-defined Definitions. In this section, we add some important definitions of the cubic sets which are subsequently used to prove our results.

Definition 1. [23] A mapping $\Upsilon : X \rightarrow [0, 1]$ is called a fuzzy set, and $\Upsilon(x)$ is a membership function and denoted by Υ .

Definition 2. [9] A structure $C = \left\{ \left(x; \tilde{\Upsilon}(x), \Upsilon(x) \mid x \in X \right) \right\}$ is a cubic set in X in which $\tilde{\Upsilon}(x)$ is IVF in X , and $\Upsilon(x)$ is fuzzy set in X . this is simply denoted by $C = \left(\tilde{\Upsilon}, \Upsilon \right)$. C^X denotes the collection of cubic sets in X .

Definition 3. [17] A neutrosophic set is a structure

$$N = \{ (x; T_N(x), I_N(x), F_N(x) \mid x \in X) \}$$

in X . Here $(T_N(x), I_N(x), F_N(x) \in [0, 1])$ are called truth, indeterminacy and falsity functions, respectively. Simply denoted by $N = (T_N, I_N, F_N)$.

Definition 4. [21] An interval neutrosophic set is a structure

$$N = \left\{ \left(x; \tilde{T}_{N_C}(x), \tilde{I}_{N_C}(x), \tilde{F}_{N_C}(x) \mid x \in X \right) \right\}$$

where $\left(x; \tilde{T}_{N_C}(x), \tilde{I}_{N_C}(x), \tilde{F}_{N_C}(x), x \in D[0, 1] \right)$ is called truth, indeterminacy and falsity function in X , respectively. This can be simply denoted by $N = \left(\tilde{T}_{N_C}, \tilde{I}_{N_C}, \tilde{F}_{N_C} \right)$.

Definition 5. [10] A structure

$$N_C = \left\{ \left(x; \tilde{T}_{N_C}(x), \tilde{I}_{N_C}(x), \tilde{F}_{N_C}(x), T_{N_C}(x), I_{N_C}(x), F_{N_C}(x) \mid x \in X \right) \right\}$$

is neutrosophic cubic set in X . Here

$$\left(\tilde{T}_{N_C} = [T_{N_C}^L, T_{N_C}^U], \tilde{I}_{N_C} = [I_{N_C}^L, I_{N_C}^U], \tilde{F}_{N_C} = [F_{N_C}^L, F_{N_C}^U] \right)$$

is an interval neutrosophic set and $(T_{N_C}, I_{N_C}, F_{N_C})$ is a neutrosophic set in X . Simply denoted by

$$N_C = \left(\tilde{T}_{N_C}, \tilde{I}_{N_C}, \tilde{F}_{N_C}, T_{N_C}, I_{N_C}, F_{N_C} \right),$$

$$[0, 0] \leq \tilde{T}_{N_C} + \tilde{I}_{N_C} + \tilde{F}_{N_C} \leq [3, 3],$$

$$0 \leq T_{N_C} + I_{N_C} + F_{N_C} \leq 1.$$

N_C^X denotes the collection of neutrosophic-cubic sets in X .

3. TRIANGULAR NEUTROSOPHIC-CUBIC NUMBERS

In this section we define the concept of neutrosophic-cubic numbers which is then used in next section.

Definition 6. A neutrosophic-cubic number

$$\begin{aligned} N_C &= \tilde{a} = \langle (a_1, a_2, a_3); \\ \tilde{T}_{N_C} &= [\alpha_a^L, \alpha_a^U], T_{N_C} = \alpha_a, \tilde{I}_{N_C} = [\theta_a^L, \theta_a^U], I_{N_C} = \theta_a, \tilde{F}_{N_C} = [\beta_a^L, \beta_a^U], F_{N_C} = \beta_a \rangle, \end{aligned}$$

such that $\langle \tilde{T}_{N_C}, \tilde{I}_{N_C}, \tilde{F}_{N_C} \rangle = R \rightarrow D[0, 1]$ and $\langle T_{N_C}, I_{N_C}, F_{N_C} \rangle = R \rightarrow [0, 1]$ is

1. a neutrosophic-cubic subset of the real lines,

2. Normal $\exists x_0 \in R$ such that $\langle \tilde{T}_{N_C} \rangle = [1, 1], \langle \tilde{I}_{N_C}, \tilde{F}_{N_C} \rangle = [0, 0]$ and $\langle T_{N_C} \rangle = 1, \langle I_{N_C}, F_{N_C} \rangle = [0, 0]$.

3. Convex for the truth membership: i.e

$$\begin{aligned} \tilde{T}_{N_C}(tx + (1-t)y) &\succcurlyeq \min\{\tilde{T}_{N_C}(x), \tilde{T}_{N_C}(y)\}, \\ T_{N_C}(tx + (1-t)y) &\geq \min\{T_{N_C}(x), T_{N_C}(y)\}, \end{aligned}$$

4. concave for the falsity and indeterminacy membership : i.e

$$\begin{aligned} \tilde{I}_{N_C}(tx + (1-t)y) &\preccurlyeq \max\{\tilde{I}_{N_C}(x), \tilde{I}_{N_C}(y)\}, \\ I_{N_C}(tx + (1-t)y) &\leq \max\{I_{N_C}(x), I_{N_C}(y)\}, \\ \tilde{F}_{N_C}(tx + (1-t)y) &\preccurlyeq \max\{\tilde{F}_{N_C}(x), \tilde{F}_{N_C}(y)\}, \\ F_{N_C}(tx + (1-t)y) &\leq \max\{F_{N_C}(x), F_{N_C}(y)\}. \end{aligned}$$

$\forall t \in [0, 1]$ $x, y \in R$, whose truth memberships functions are:

$$(3.1) \quad \tilde{T}_{\tilde{a}}(x) = \begin{cases} \frac{x-d}{e-d} [T_{\tilde{a}}^L, T_{\tilde{a}}^U] & d \leq x < e \\ \frac{f-x}{f-e} [T_{\tilde{a}}^L, T_{\tilde{a}}^U] & e < x \leq f \\ 0 & \text{otherwise} \end{cases}$$

$$(3.2) \quad T_{\tilde{a}}(x) = \begin{cases} \frac{x-d}{e-d} T_{\tilde{a}} & d \leq x < e \\ \frac{f-x}{f-e} T_{\tilde{a}} & e < x \leq f \\ 0 & \text{otherwise} \end{cases}$$

indeterminacy memberships functions:

$$(3.3) \quad \tilde{I}_{\tilde{a}}(x) = \begin{cases} \frac{(e-x)+(x-d)}{e-d} [I_{\tilde{a}}^L, I_{\tilde{a}}^U] & d \leq x < e \\ \frac{(x-e)+(f-x)}{f-e} [I_{\tilde{a}}^L, I_{\tilde{a}}^U] & e < x \leq f \\ 1 & \text{otherwise} \end{cases}$$

$$(3.4) \quad I_{\tilde{a}}(x) = \begin{cases} \frac{(e-x)+(x-d)}{e-d} I_{\tilde{a}} & d \leq x < e \\ \frac{(x-e)+(f-x)}{f-e} I_{\tilde{a}} & e < x \leq f \\ 1 & \text{otherwise} \end{cases}$$

falsity memberships functions:

$$(3.5) \quad \tilde{F}_{\tilde{a}}(x) = \begin{cases} \frac{e-x}{e-d} [F_{\tilde{a}}^L, F_{\tilde{a}}^U] & d \leq x < e \\ \frac{x-e}{f-e} [F_{\tilde{a}}^L, F_{\tilde{a}}^U] & e < x \leq f \\ 1 & \text{otherwise} \end{cases}$$

$$(3.6) \quad F_{\tilde{a}}(x) = \begin{cases} \frac{e-x}{e-d} F_{\tilde{a}} & d \leq x < e \\ \frac{x-e}{f-e} F_{\tilde{a}} & e < x \leq f \\ 1 & \text{otherwise} \end{cases}$$

Definition 7. Let

$$\tilde{a} = \langle (a_1, a_2, a_3), [T_{\tilde{a}}^L, T_{\tilde{a}}^U], T_{\tilde{a}}, [I_{\tilde{a}}^L, I_{\tilde{a}}^U], I_{\tilde{a}}^L, [F_{\tilde{a}}^L, F_{\tilde{a}}^U], F_{\tilde{a}}^U \rangle,$$

and

$$\tilde{b} = \langle (b_1, b_2, b_3), [T_{\tilde{b}}^L, T_{\tilde{b}}^U], T_{\tilde{b}}, [I_{\tilde{b}}^L, I_{\tilde{b}}^U], I_{\tilde{b}}^L, [F_{\tilde{b}}^L, F_{\tilde{b}}^U], F_{\tilde{b}}^U \rangle,$$

be two triangular neutrosophic cubic numbers and $\gamma \neq 0$ be any real number. Then:

1. Addition of two TNCNs are

$$(3.7) \quad \tilde{a} + \tilde{b} = \left\{ \begin{array}{l} \langle (a_1 + b_1, a_2 + b_2, a_3 + b_3), \\ \left[\min(T_a^L, T_b^L), \min(T_a^U, T_b^U) \right], \min(T_{\tilde{a}}, T_{\tilde{b}}), \\ \left[\max(I_a^L, I_b^L), \max(I_a^U, I_b^U) \right], \max(I_{\tilde{a}}, I_{\tilde{b}}), \\ \left[\max(F_a^L, F_b^L), \max(F_a^U, F_b^U) \right], \max(F_{\tilde{a}}, F_{\tilde{b}}) \rangle \end{array} \right\}$$

2. Subtraction of two TNCNs are

$$(3.8) \quad \tilde{a} - \tilde{b} = \left\{ \begin{array}{l} \langle (a_1 - b_3, a_2 - b_2, a_3 - b_1), \\ \left[\min(T_a^L, T_b^L), \min(T_a^U, T_b^U) \right], \min(T_{\tilde{a}}, T_{\tilde{b}}), \\ \left[\max(I_a^L, I_b^L), \max(I_a^U, I_b^U) \right], \max(I_{\tilde{a}}, I_{\tilde{b}}), \\ \left[\max(F_a^L, F_b^L), \max(F_a^U, F_b^U) \right], \max(F_{\tilde{a}}, F_{\tilde{b}}) \rangle \end{array} \right\}$$

3. Multiplication of a TNCNs by a scalar value are

$$(3.9) \quad \gamma \tilde{a} = \left\{ \begin{array}{l} \langle (\gamma a_1, \gamma a_2, \gamma a_3), [T_a^L, T_a^U], T_{\tilde{a}}, \\ [I_a^L, I_a^U], I_{\tilde{a}}, [F_a^L, F_a^U], F_{\tilde{a}}^U \rangle \text{ if } (\gamma > 0) \\ \langle (\gamma a_3, \gamma a_2, \gamma a_1), [T_a^L, T_a^U], T_{\tilde{a}}, [I_a^L, I_a^U], \\ I_{\tilde{a}}^L, [F_a^L, F_a^U], F_{\tilde{a}}^U \rangle \text{ if } (\gamma < 0) \end{array} \right\}$$

4. Inverse of a TNCNs are

$$(3.10) \quad \tilde{a}^{-1} = \left\{ \begin{array}{l} \langle \left(\frac{1}{a_3}, \frac{1}{a_2}, \frac{1}{a_1} \right), [T_a^L, T_a^U], T_{\tilde{a}}, \\ [I_a^L, I_a^U], I_{\tilde{a}}^L, [F_a^L, F_a^U], F_{\tilde{a}}^U \rangle \text{ where } (\tilde{a} \neq 0) \end{array} \right\}$$

5. Multiplication of two TNCNs are

$$(3.11) \quad \tilde{a}\tilde{b} = \left\{ \begin{array}{l} \langle (a_1 b_1, a_2 b_2, a_3 b_3); \\ \left[\min(T_a^L, T_b^L), \min(T_a^U, T_b^U) \right] \min(T_{\tilde{a}}, T_{\tilde{b}}), \\ \left[\max(I_a^L, I_b^L), \max(I_a^U, I_b^U) \right] \max(I_{\tilde{a}}, I_{\tilde{b}}), \\ \left[\max(F_a^L, F_b^L), \max(F_a^U, F_b^U) \right] \max(F_{\tilde{a}}, F_{\tilde{b}}) \rangle \\ \text{if } (a_3 > 0, b_3 > 0) \\ \langle (a_1 b_3, a_2 b_2, a_3 b_1); \\ \left[\min(T_a^L, T_b^L), \min(T_a^U, T_b^U) \right] \min(T_{\tilde{a}}, T_{\tilde{b}}), \\ \left[\max(I_a^L, I_b^L), \max(I_a^U, I_b^U) \right] \max(I_{\tilde{a}}, I_{\tilde{b}}), \\ \left[\max(F_a^L, F_b^L), \max(F_a^U, F_b^U) \right] \max(F_{\tilde{a}}, F_{\tilde{b}}) \rangle \\ \text{if } (a_3 < 0, b_3 > 0) \\ \langle (a_1 b_3, a_2 b_2, a_3 b_1); \\ \left[\min(T_a^L, T_b^L), \min(T_a^U, T_b^U) \right] \min(T_{\tilde{a}}, T_{\tilde{b}}), \\ \left[\max(I_a^L, I_b^L), \max(I_a^U, I_b^U) \right] \max(I_{\tilde{a}}, I_{\tilde{b}}), \\ \left[\max(F_a^L, F_b^L), \max(F_a^U, F_b^U) \right] \max(F_{\tilde{a}}, F_{\tilde{b}}) \rangle \\ \text{if } (a_3 < 0, b_3 < 0) \end{array} \right\}$$

6. Division of two TNCNs are

$$(3.12) \quad \frac{\tilde{a}}{\tilde{b}} = \left\{ \begin{array}{l} \left(\frac{a_1}{b_3}, \frac{a_2}{b_2}, \frac{a_3}{b_1} \right); \\ \left[\min(T_{\tilde{a}}^L, T_{\tilde{b}}^L), \min(T_{\tilde{a}}^U, T_{\tilde{b}}^U) \right] \min(T_{\tilde{a}}, T_{\tilde{b}}), \\ \left[\max(I_{\tilde{a}}^L, I_{\tilde{b}}^L), \max(I_{\tilde{a}}^U, I_{\tilde{b}}^U) \right] \max(I_{\tilde{a}}, I_{\tilde{b}}), \\ \left[\max(F_{\tilde{a}}^L, F_{\tilde{b}}^L), \max(F_{\tilde{a}}^U, F_{\tilde{b}}^U) \right] \max(F_{\tilde{a}}, F_{\tilde{b}}) \\ \text{if } (a_3 > 0, b_3 > 0) \\ \left(\frac{a_3}{b_3}, \frac{a_2}{b_2}, \frac{a_1}{b_1} \right); \\ \left[\min(T_{\tilde{a}}^L, T_{\tilde{b}}^L), \min(T_{\tilde{a}}^U, T_{\tilde{b}}^U) \right] \min(T_{\tilde{a}}, T_{\tilde{b}}), \\ \left[\max(I_{\tilde{a}}^L, I_{\tilde{b}}^L), \max(I_{\tilde{a}}^U, I_{\tilde{b}}^U) \right] \max(I_{\tilde{a}}, I_{\tilde{b}}), \\ \left[\max(F_{\tilde{a}}^L, F_{\tilde{b}}^L), \max(F_{\tilde{a}}^U, F_{\tilde{b}}^U) \right] \max(F_{\tilde{a}}, F_{\tilde{b}}) \\ \text{if } (a_3 > 0, b_3 > 0) \\ \left(\frac{a_3}{b_1}, \frac{a_2}{b_2}, \frac{a_1}{b_3} \right); \\ \left[\min(T_{\tilde{a}}^L, T_{\tilde{b}}^L), \min(T_{\tilde{a}}^U, T_{\tilde{b}}^U) \right] \min(T_{\tilde{a}}, T_{\tilde{b}}), \\ \left[\max(I_{\tilde{a}}^L, I_{\tilde{b}}^L), \max(I_{\tilde{a}}^U, I_{\tilde{b}}^U) \right] \max(I_{\tilde{a}}, I_{\tilde{b}}), \\ \left[\max(F_{\tilde{a}}^L, F_{\tilde{b}}^L), \max(F_{\tilde{a}}^U, F_{\tilde{b}}^U) \right] \max(F_{\tilde{a}}, F_{\tilde{b}}) \\ \text{if } (a_3 > 0, b_3 > 0) \end{array} \right.$$

7. Division of a TNCNs by a scalar value are

$$(3.13) \quad \frac{\tilde{a}}{\gamma} = \left\{ \begin{array}{l} \left(\frac{a_1}{\gamma}, \frac{a_2}{\gamma}, \frac{a_3}{\gamma} \right), [T_{\tilde{a}}^L, T_{\tilde{a}}^U], T_{\tilde{a}}, [I_{\tilde{a}}^L, I_{\tilde{a}}^U], I_{\tilde{a}}^L, [F_{\tilde{a}}^L, F_{\tilde{a}}^U], F_{\tilde{a}}^U \text{ if } (\gamma > 0) \\ \left(\frac{a_3}{\gamma}, \frac{a_2}{\gamma}, \frac{a_1}{\gamma} \right), [T_{\tilde{a}}^L, T_{\tilde{a}}^U], T_{\tilde{a}}, [I_{\tilde{a}}^L, I_{\tilde{a}}^U], I_{\tilde{a}}^L, [F_{\tilde{a}}^L, F_{\tilde{a}}^U], F_{\tilde{a}}^U \text{ if } (\gamma < 0) \end{array} \right.$$

4. NEUTROSOPHIC CUBIC ANALYTIC HIERARCHY PROCESS

In this section, we introduced the AHP in the neutrosophic cubic environment. The Analytic Hierarchy Process includes three stages:

- (1) Decomposition,
- (2) Pair-wise comparison,
- (3) Synthesis of priorities.

Next a stepwise procedure is described for the model.

Step (1) Construct the structure of hierarchy of the problem, which has three levels (i) The first level is goal which organization wants to attained. (ii) The second level that consists of the criteria and sub criteria. (iii) The third level is how the different alternatives are evaluated. The general hierarchy is presented in figure1. The next step is applied for the weighting criteria, sub-criteria and alternatives, according to expert's opinions

Step (2) First analyzing the intricate multi- criteria decision making problem into a hierarchical structure then use neutrosophic-cubic pair-wise comparison matrix. The vagueness of decision makers is represented by triangular neutrosophic-cubic number (TNCNs) and then construct neutrosophic- cubic pair-wise comparison matrix of criteria, sub-criteria and alternatives structured through the linguistic terms which are shown in Table 2.

Table [2].	Linguistic terms and the identical triangular neutrosophic-cubic numbers.	
Saaty Scale	Explanation	Neutrosophic-Cubic Triangular Scale
1	Equally influential	$\tilde{1} = \langle (1, 1, 1); [0.50, 0.50], 0.50, [0.50, 0.50], 0.50, [0.50, 0.50], 0.50 \rangle$
3	Slightly influential	$\tilde{3} = \langle (2, 3, 4); [0.25, 0.35], 0.30, [0.55, 0.65], 0.60, [0.65, 0.75], 0.70 \rangle$
5	Strongly influential	$\tilde{5} = \langle (4, 5, 6); [0.65, 0.75], 0.70, [0.15, 0.25], 0.20, [0.25, 0.35], 0.30 \rangle$
7	Very Strongly influential	$\tilde{7} = \langle (6, 7, 8); [0.85, 0.95], 0.90, [0.20, 0.20], 0.20, [0.20, 0.20], 0.20 \rangle$
9	Absolutely influential	$\tilde{9} = \langle (9, 9, 9); [1.00, 1.00], 1.00, [0.05, 0.05], 0.05, [0.10, 0.10], 0.10 \rangle$
2		$\tilde{2} = \langle (1, 2, 3); [0.50, 0.50], 0.50, [0.50, 0.50], 0.50, [0.50, 0.50], 0.50 \rangle$
4	intermediate values	$\tilde{4} = \langle (3, 4, 5); [0.50, 0.50], 0.50, [0.50, 0.50], 0.50, [0.50, 0.50], 0.50 \rangle$
6		$\tilde{6} = \langle (6, 7, 8); [0.50, 0.50], 0.50, [0.50, 0.50], 0.50, [0.50, 0.50], 0.50 \rangle$
8		$\tilde{8} = \langle (7, 8, 9); [0.50, 0.50], 0.50, [0.50, 0.50], 0.50, [0.50, 0.50], 0.50 \rangle$

The neutrosophic cubic scale is obtained on the basis of expert’s opinion. The neutrosophic cubic pair-wise comparison matrix are as follows of criteria, sub-criteria and alternatives:

$$(4.1) \quad \tilde{A} = \begin{bmatrix} \tilde{1} & \tilde{a}_{12} & \cdot & \cdot & \cdot & \tilde{a}_{1n} \\ \tilde{a}_{21} & \cdot & & & & \cdot \\ \cdot & \cdot & \cdot & & & \cdot \\ \cdot & \cdot & & \cdot & & \cdot \\ \cdot & \cdot & & & \cdot & \cdot \\ \tilde{a}_{n1} & \tilde{a}_{n2} & \cdot & \cdot & \cdot & \tilde{1} \end{bmatrix}$$

where $\tilde{a}_{ji} = \tilde{a}_{ij}^{-1}$ is the triangular neutrosophic cubic number which is used to measures the imprecision in decision.

Step (3) For calculating overall preferences of each alternative and to conclude final ranking, we should first determine weights of each standard from the corresponding pair-wise comparison matrix, by transforming it to a deterministic matrix using the following equation.

Let

$$\tilde{a}_{ij} = \langle (a_1, a_2, a_3); [T_{\tilde{a}}^L, T_{\tilde{a}}^U], T_{\tilde{a}}, [I_{\tilde{a}}^L, I_{\tilde{a}}^U], I_{\tilde{a}}, [F_{\tilde{a}}^L, F_{\tilde{a}}^U], F_{\tilde{a}} \rangle$$

be a triangular neutrosophic-cubic number; then

$$(4.2) \quad S(\tilde{a}_{ij}) = \frac{1}{21} [a_1 + a_2 + a_3] \left(+ [3 + T_{\tilde{a}}^U + I_{\tilde{a}}^U + F_{\tilde{a}}^U] + [3 + T_{\tilde{a}} + I_{\tilde{a}} + F_{\tilde{a}}] \right)$$

and

$$(4.3) \quad A(\tilde{a}_{ij}) = \frac{1}{21} [a_1 + a_2 + a_3] \left(+ [3 + T_{\tilde{a}}^U + I_{\tilde{a}}^U + F_{\tilde{a}}^U] - [3 + T_{\tilde{a}} + I_{\tilde{a}} + F_{\tilde{a}}] \right)$$

are the score function and accuracy function.

To acquire the score and accuracy function of \tilde{a}_{ji} , we use the following equation

$$(4.4) \quad S(\tilde{a}_{ji}) = \frac{1}{S(\tilde{a}_{ij})}$$

$$(4.5) \quad A(\tilde{a}_{ji}) = \frac{1}{A(\tilde{a}_{ij})}$$

With compensation by score value of each triangular neutrosophic-cubic number in the neutrosophic-cubic pair-wise comparison matrix, we get the following deterministic matrix;

$$(4.6) \quad A = \begin{bmatrix} 1 & a_{12} & \cdot & \cdot & \cdot & a_{1n} \\ a_{21} & \cdot & & & & \cdot \\ \cdot & \cdot & \cdot & & & \cdot \\ \cdot & \cdot & & \cdot & & \cdot \\ \cdot & \cdot & & & \cdot & \cdot \\ a_{n1} & a_{n2} & \cdot & \cdot & \cdot & 1 \end{bmatrix}$$

Now we can easily find ranking of priorities namely eigen vector X, from the above matrix.

- (1) Normalize the column entries by dividing each entry by the sum of the column.

(2) Take the total of the row averages.

Step (4) To measure inconsistency; within the decision in each pair-wise comparison matrix the complete hierarch, AHP methodology gives a consistency index .Check if any inconsistency in a neutrosophic-cubic comparison matrix. AHP uses consistency index and consistency ratio, If consistency ratio is greater than 10%, then judgment are unreliable because they are so closely uncertainty and the procedure is not correct and must be repeated.

To compute CI and CR do the following steps:

1. Multiply each value in the first column of the pair-wise comparison matrix by the priority of the first item; we will keep the process same for all the columns of the pair-wise comparison matrix. Sum the values across the rows to get a vector of values (weighted sum).
2. Get the elements of the weighted sum vector by the corresponding priority for each criterion.
3. Calculate the average of the values found in step 2; this average is denoted by $\lambda \max$.
4. Calculate the consistency index (CI) as follows:

$$(4.7) \quad CI = \frac{\lambda \max - n}{n - 1}$$

where n is the number of items which can be compared.

5. Calculate the consistency ratio, which is defined as:

$$(4.8) \quad CR = \frac{CI}{RI}$$

where RI is the consistency index of a randomly generated comparison matrix.

Step (5) In the last step, calculate overall weight (priority) of each alternative. Final ranking of all the alternatives is calculated using equation [22].

$$(4.9) \quad T_{w_{Alt_i}} = \sum_{j=1}^n (w_j \otimes w_{ij})$$

where $(j = 1, \dots, n)$. From previous steps, we achieve the phases of neutrosophic-cubic analytic hierarchy process, as shown in Figure 1.




Figure 1. Neutrosophic-cubic AHP schematic diagram

5. NUMERICAL APPLICATION

The model proposed in Section 5 is used to solve a real case study as follows.

The goal is to select the best among the alternatives. There are three alternatives which are Faisal Movers, Niazi Express and Bala Gujjar travels. The best alternative is selected on the basis of different criteria, which are defines as:

- (1) C_1 :Flexibility
- (2) C_2 :Safety
- (3) C_3 : Affordable
- (4) C_4 :Punctuality
- (5) C_5 :Enjoyable

Now decomposed in hierarchy structure can be shown as in figure 2. The hierarchy consists of four levels. The aim of objective is placed at Level 1, criteria are as Level 2, and the last, alternatives are as Level 3. Step 3 Structure the cubic pair-wise comparison matrix of a criteria, sub-criteria and alternative, through the linguistic terms are shown in table 1. The values in table 2 are concern to a experts opinions. The pair-wise comparison matrix of criteria is presented in **Table [3]**.




Figure 2. Structure of hierarchy problem

Table [3]The neutrosophic-cubic pair-wise comparison matrix of criteria

Criteria	Flexibility	Safety	Affordable	Punctuality	Enjoyable
Flexibility	$\tilde{1}$	$\tilde{2}$	$\tilde{4}$	$\tilde{9}$	$\tilde{5}$
Safety	$\tilde{2}^{-1}$	$\tilde{1}$	$\tilde{7}$	$\tilde{6}$	$\tilde{2}$
Affordable	$\tilde{4}^{-1}$	$\tilde{7}^{-1}$	$\tilde{1}$	$\tilde{5}$	$\tilde{3}$
Punctuality	$\tilde{9}^{-1}$	$\tilde{6}^{-1}$	$\tilde{5}^{-1}$	$\tilde{1}$	$\tilde{2}$
Enjoyable	$\tilde{5}^{-1}$	$\tilde{2}^{-1}$	$\tilde{3}^{-1}$	$\tilde{2}^{-1}$	$\tilde{1}$

Where;

$$\begin{aligned} \tilde{1} &= \langle (1, 1, 1); [0.50, 0.50], 0.50, [0.50, 0.50], 0.50, [0.50, 0.50], 0.50 \rangle, \\ \tilde{3} &= \langle (2, 3, 4); [0.25, 0.35], 0.30, [0.55, 0.65], 0.60, [0.65, 0.75], 0.70 \rangle, \\ \tilde{3} &= \langle (2, 3, 4); [0.25, 0.35], 0.30, [0.55, 0.65], 0.60, [0.65, 0.75], 0.70 \rangle, \\ \tilde{1} &= \langle (1, 1, 1); [0.50, 0.50], 0.50, [0.50, 0.50], 0.50, [0.50, 0.50], 0.50 \rangle, \end{aligned}$$

Using the equation(15), the above neutrosophic-cubic pair-wise comparison matrix transformed into deterministic pair-wise comparison matrix as shown in Table [4]

Table [4] deterministic pair-wise comparison matrix of criteria

Criteria	Flexibility	Safety	Affordable	Punctuality	Enjoyable
Flexibility	1	1.3	2.45	5.3	3
Safety	.76	1	4.3	3.6	1.3
Affordable	.41	.23	1	3	2
Punctuality	.19	.27	.33	1	1.3
Enjoyable	.33	.76	.50	.76	1

Next determined the ranking of the Criteria, namely the Eigen Vector X , from the previous matrix, as illustrated previously in the detailed steps of the proposed model. The normalized comparison matrix of Criteria is presented in Table [5].

Table[5] The normalized comparison matrix of Criteria

Criteria	Flexibility	Safety	Affordable	Punctuality	Enjoyable
Flexibility	.37	.36	.30	.40	.35
Safety	.28	.28	.50	.26	.15
Affordable	.15	.07	.17	.22	.23
Punctuality	.07	.08	.04	.07	.15
Enjoyable	.12	.21	.06	.06	.12

Now taking the total of the row averages:

$$X = \begin{bmatrix} .356 \\ .294 \\ .164 \\ .081 \\ .114 \end{bmatrix}$$

We should make sure that all the inputs given by the expert's are very consistent, now we take the consistency test as follows: Firstly calculate consistency index and consistency ratio as follows:

$$A = \begin{pmatrix} 1 & 1.3 & 2.45 & 5.3 & 3 \\ .76 & 1 & 4.3 & 3.6 & 1.3 \\ .41 & .23 & 1 & 3 & 2 \\ .19 & .27 & .33 & 1 & 1.3 \\ .33 & .76 & .50 & .76 & 1 \end{pmatrix} \begin{pmatrix} .356 \\ .294 \\ .164 \\ .081 \\ .114 \end{pmatrix} = \begin{pmatrix} 1.922 \\ .3285 \\ .1447 \\ .0755 \\ .1317 \end{pmatrix}$$

from the equation (20) and (21),

$$\lambda \max = \text{average} \left\{ \begin{array}{l} \frac{1.922}{0.356}, \frac{0.3285}{0.294}, \frac{0.1447}{0.164}, \\ \frac{0.0755}{0.081}, \frac{0.1317}{0.114} \end{array} \right\} = 5.399$$

and $CI = \frac{\lambda \max - n}{n - 1} = \frac{5.399 - 5}{5 - 1} = 0.089975$.

For the calculation of value of consistency ratio, we used *Table [6]* which was taken from Saaty book. The upper row is in random matrix order, and the lower row is the corresponding index of consistency for random decision.

Table [6]

Saaty table for calculating consistency ratio

0	2	3	4	5	6	7	8	9	10
0	0	.58	.90	1.12	1.24	1.32	1.4	1.45	1.49

$$CR = \frac{CI}{RI} = \frac{0.09975}{1.12} = 0.089 = 8.9\%$$

so evaluation are consistent i.e

$$8.9\% < 10\%.$$

The neutrosophic cubic pair-wise comparison matrix of decision criterias with respect to C_1 is presented in *Table [7]*

Table [7]

Criteria	Faisal Movers	Niazi Express	Bala Gujjar Travels
Faisal Movers	1	$\tilde{5}^{-1}$	$\tilde{2}^{-1}$
Niazi Express	$\tilde{5}$	1	$\tilde{5}$
Bala Gujjar Travels	$\tilde{2}$	$\tilde{5}^{-1}$	1

The deterministic pair-wise comparison matrix of decision criteria w.r.t C_1 shown in *Table [8]* and the normalized comparison matrix in shown in *Table [9]*

Table [8]

Criteria	Faisal Movers	Niazi Express	Bala Gujjar Travels
Faisal Movers	1	.33	.76
Niazi Express	3	1	3
Bala Gujjar Travels	1.3	.33	1

Table [9]

Criteria	Faisal Movers	Niazi Express	Bala Gujjar Travels
Faisal Movers	.18	.18	.16
Niazi Express	.57	.63	.63
Bala Gujjar Travels	.25	.18	.21

Now taking the total of the row averages;

$$X = \begin{bmatrix} .17 \\ .61 \\ .22 \end{bmatrix}$$

The neutrosophic-cubic pair-wise comparison matrix of decision criteria with respect to C_2 is presented in *Table [10]*.

Table [10]

Criteria2	Faisal Movers	Niazi Express	Bala Gujjar Travels
Faisal Movers	1	1	$\tilde{5}^{-1}$
Niazi Express	$\tilde{1}$	$\tilde{1}$	$\tilde{6}$
Bala Gujjar Travels	$\tilde{5}$	$\tilde{6}^{-1}$	$\tilde{1}$

The deterministic pair-wise comparison matrix of decision criteria w.r.t C_2 shown in table *Table [11]* and the normalized comparison matrix in shown in *Table [12]*

Table [11]

Criteria	Faisal Movers	Niazi Express	Bala Gujjar Travels
Faisal Movers	1	1	3
Niazi Express	1	1	4.3
Bala Gujjar Travels	.33	.23	1

Table [12]

Criteria2	Faisal Movers	Niazi Express	Bala Gujjar Travels
Faisal Movers	.43	.45	.36
Niazi Express	.43	.45	.52
Bala Gujjar Travels	.13	.10	.12

Now taking the total of the row averages;

$$X = \begin{bmatrix} .41 \\ .47 \\ .12 \end{bmatrix}$$

The neutrosophic-cubic pair-wise comparison matrix of decision criteria with respect to C_3 is presented in *Table [13]*

Table [13]

Criteria3	Faisal Movers	Niazi Express	Bala Gujjar Travels
Faisal Movers	1	3^{-1}	4
Niazi Express	3	1	5
Bala Gujjar Travels	4^{-1}	5^{-1}	1

The deterministic pair-wise comparison matrix of decision criteria w.r.t C_3 shown in *Table [14]* and the normalized comparison matrix in shown in *Table [15]*

Table [14]

Criteria	Faisal Movers	Niazi Express	Bala Gujjar Travels
Faisal Movers	1	.50	2.45
Niazi Express	2	1	3
Bala Gujjar Travels	.41	.33	1

Table [15]

Criteria3	Faisal Movers	Niazi Express	Bala Gujjar Travels
Faisal Movers	.29	.27	.38
Niazi Express	.58	.55	.46
Bala Gujjar Travels	.12	.17	.16

Now taking the total of the row averages;

$$X = \begin{bmatrix} .31 \\ .53 \\ .16 \end{bmatrix}$$

The neutrosophic-cubic pair-wise comparison matrix of decision criteria with respect to C_4 is presented in Table [16]

Table [16]

Criteria3	Faisal Movers	Niazi Express	Bala Gujjar Travels
Faisal Movers	1	3^{-1}	5^{-1}
Niazi Express	3	1	1
Bala Gujjar Travels	5	1	1

The deterministic pair-wise comparison matrix of decision criteria w.r.t C_4 shown in Table [17] and the normalized comparison matrix in shown in Table [18]

Table [17]

Criteria4	Faisal Movers	Niazi Express	Bala Gujjar Travels
Faisal Movers	1	.50	.33
Niazi Express	2.45	1	1
Bala Gujjar Travels	3	1	1

Table [18]

Criteria3	Faisal Movers	Niazi Express	Bala Gujjar Travels
Faisal Movers	.15	.17	.13
Niazi Express	.38	.41	.43
Bala Gujjar Travels	.47	.41	.43

Now taking the total of the row averages;

$$X = \begin{bmatrix} .15 \\ .41 \\ .44 \end{bmatrix}$$

The neutrosophic–cubic pair-wise comparison matrix of decision criteria with respect to C_5 is presented in *Table [19]*

Table [19]

Criteria3	Faisal Movers	Niazi Express	Bala Gujjar Travels
Faisal Movers	1	2^{-1}	3
Niazi Express	2	1	4
Bala Gujjar Travels	3^{-1}	4^{-1}	1

The deterministic pair-wise comparison matrix of decision criteria w.r.t C_5 shown in *Table [20]* and the normalized comparison matrix in shown in *Table [21]*

Table [20]

Criteria4	Faisal Movers	Niazi Express	Bala Gujjar Travels
Faisal Movers	1	.76	2
Niazi Express	1.3	1	2.45
Bala Gujjar Travels	.5	.41	1

Table [21]

Criteria3	Faisal Movers	Niazi Express	Bala Gujjar Travels
Faisal Movers	.36	.35	.37
Niazi Express	.46	.46	.45
Bala Gujjar Travels	.18	.19	.18

Now taking the total of the row averages;

$$X = \begin{bmatrix} .36 \\ .46 \\ .18 \end{bmatrix}$$

In *Table [22]*, Now the Total priority (weight) vector of the three alternatives with respect to Goal:


Table [22]

	C_1	C_2	C_3	C_4	C_5	Total Weight	Total Ranking of alternatives
Weights	0.356	0.294	0.164	0.081	0.114		
Faisal Movers	.17	0.41	0.31	0.15	.36	.285	2 nd
Niazi Express	.61	0.47	0.53	0.41	.46	.52	1 st
Bala Gujjar Travels	.22	0.12	0.16	0.44	.18	.196	3 rd

Hence the final ranking of the best alternative are

$$\text{Niazi Express} > \text{Faisal Movers} > \text{Bala Gujjar Travels},$$

presented in figure 3.



6. COMPARSION ANALYSIS

The comparison with other existing method is used to estimate the quality of the proposed model. In AHP we use non-fuzzy value to build a comparison matrix, which are not accurate because of ambiguity, inconsistent and doubtful information of decision maker. An idea of crisp information can be extended by many researchers in fuzzy environment for fulfilling the drawbacks of classical AHP to get the desired result, in which he takes triangular fuzzy numbers but he fails to describe the relative weights among it. In cubic environment they are only describe the membership in terms of interval valued and fuzzy point and not discuss the non-membership and indeterminacy. To overcome the drawback of fuzzy AHP the cubic AHP are introduced, which is more reliable then fuzzy AHP. Then the neutrosophic-cubic set are the generalization of neutrosophic and interval neutrosophic set, and considered the truth, false and indeterminacy membership in the forms of intervals and also points, which is a best to representation for a vague and unsure information that exist in real life. We are the first to introduce the AHP in the neutrosophic-cubic environment. We can apply this method to various fields for the solution of different problems for achieving desired goals. in Table [23]. We compare the result of the NC-AHP with the other existing methods, which are

Table [23] Comparison of other existing methods

Methods	Final ranking
AHP [14]	Faisal movers \geq Bala gujjar travels \geq Niazi express
F-AHP [2]	Bala gujjar travels \geq Niazi express \geq Faisal movers
IF-AHP[20]	Bala gujjar travels \geq Niazi express \geq Faisal movers
N-AHP[1]	Niazi express \geq Bala gujjar travels \geq Faisal movers
NC-AHP Our porposed method	Niazi express \geq Faisal movers \geq Bala gujjar travels

7. CONCLUSION AND FUTURE WORK

Decision making is vital in daily life, dealing with real life problems; one who makes decision faces the data which is incomplete and not clear. The trails of neutrosophic-cubic enable that very situation. In the result of this, for every situation we define triangle neutrosophic-cubic number (TNCNs) and their operation laws. Mostly it is difficult to compare the neutrosophic-cubic value. The score and accuracy functions are defined for the comparison of neutrosophic-cubic values. By using these operations, we define analytic hierarchy process (AHP) in neutrosophic-cubic environment, which is the more generalized version of neutrosophic sets and interval neutrosophic sets. Since neutrosophic-cubic sets are placed a better way to express inconsistent, indeterminate and obscure data. This method can be applied to the problems of daily life effectively and can be solved many intricate problems.

REFERENCES

- [1] M. A. Basset, M. Mohamed, Y. Zhou and I. Hezam, Multi-criteria group decision making based on neutrosophic analytic hierarchy process, *J. Int. & Fuzzy Syst.* 33, (2017), 4055–4066.
- [2] J. J. Buckley, Fuzzy hierarchical analysis, *Fuzzy Sets Syst.*, 17, (1985), 233–247.
- [3] U. Cebeci, Fuzzy AHP-based decision support system for selecting ERP systems in textile industry by using balanced scorecard. *Expert Syst. Appl.*, 36, (2009), 8900–8909.
- [4] D. Y. Chang, Applications of the extent analysis method on fuzzy AHP, *Eur. J. Oper. Res.*, 95, (1996), 649–655.
- [5] C. W. Chang, C. R. Wu, H. C. Chen, Using expert technology to select unstable slicing machine to control wafer slicing quality via fuzzy AHP. *Expert Syst. Appl.*, 34, (2008), 2210–2220.
- [6] O. Duran, J. Aguilo, Computer-aided machine-tool selection based on a Fuzzy-AHP approach. *Expert Syst. Appl.*, 34, (2008), 1787–1794.
- [7] Y. L. Hsu, C. H. Lee, V. B. Kreng, The application of Fuzzy Delphi Method and Fuzzy AHP in lubricant regenerative technology selection. *Expert Syst. Appl.*, 37, (2010), 419–425.
- [8] Y. Ju, A. Wang, X. Liu, Evaluating emergency response capacity by fuzzy AHP and 2-tuple fuzzy linguistic approach. *Expert Syst. Appl.*, 39, (2012), 6972–6981.
- [9] Y.B. Jun, C. S. Kim and K.O. Yang, Cubic Sets, *Ann. Fuzzy Math. Inform.* 4(1), (2012), 83-98.
- [10] Y. B. Jun, F. Smarandache and C. S. Kim, Neutrosophic Cubic Sets, *New Math. Natural Compt.* 13(1), (2015), 41-54.
- [11] O. Kulak, C. Kahraman, Fuzzy multi-attribute selection among transportation companies using axiomatic design and analytic hierarchy process, *Inform. Sci.*, 170, (2005), 191–210.
- [12] J. Rezaei, P. B. M. Fahim, L. Tavasszy, Supplier selection in the airline retail industry using a funnel methodology, Conjunctive screening method and fuzzy AHP. *Expert Syst. Appl.*, 41, (2014), 8165–8179.
- [13] T.L.Saaty, A Scaling Method for Priorities in Hierarchical Structures, *J. Mathematical Psychology*, 15, (1977), 57-68.
- [14] T.L.Saaty, *The Analytic Hierarchy Process*. McGraw-Hill International, New York, NY, U.S.A, (1980).
- [15] T.L.Saaty, What is the analytic hierarchy process. In *Mathematical Models for Decision Supported*; Springer: Berlin, Germany, (1988), 109–121.
- [16] T.L.Saaty, Axiomatic foundations of the Analytic Hierarchy Process, *Manag. Sci.* 32(7), (1986) 841-855.5.
- [17] F. Smarandache, A Unifying Field in Logics: Neutrosophic Logic. Neutrosophy, Neutrosophic Set, Neutrosophic Probability; American Reserch Press: Rehoboth, IL, USA, 1999.
- [18] F. Smarandache, Neutrosophic set-a generalization of the intuitionistic fuzzy set. *Int. J. Pure Appl. Math.* 2005, 24, 287–297.
- [19] P. J. M. Van Laarhoven and W. Pedrycz, A fuzzy extension of Saaty's priority theory, *Fuzzy Sets Syst.*, 11, (1983), 229–241.

- [20] Z. S. Xu and H.C. Liao, Intuitionistic fuzzy analytic hierarchy process, IEEE Trans. Fuzzy Syst., 22(4) (2014).
- [21] H. Wang, F. Smarandache, Y.Q. Zhang, R. Sunderraman, Interval Neutrosophic Sets and Logic: Theory and Applications in Computing; Hexis: Phoenix, AZ, USA, 2005.
- [22] L. YF, M. H. Wen, A fuzzy-AHP-based technique for the decision of design feature selection in massively multiplayer online roleplayinggame development, Expert Syst. Appl., 37, (2010), 8685–8693.
- [23] L. A. Zadeh, Fuzzy sets, Inform. Control., 8, (1965), 338–353.

DEPARTMENT OF MATHEMATICS AND STATISTICS , HAZARA UNIVERSITY MANSEH, KP PAKISTAN
E-mail address: gulistanmath@hu.edu.pk

CENTRE FOR MATHEMATICS AND STATISTICAL SCIENCES, LAHORE SCHOOL OF ECONOMICS, LAHORE, PAKISTAN
E-mail address: ibeg@lahoreschool.edu.pk

Current address: Department of Mathematics and Statistics , Hazara University Mansh, KP Pakistan
E-mail address: asiftanoli5219@gmail.com