

PAPER • OPEN ACCESS

## Neutrosophic generalized b-closed sets in Neutrosophic topological spaces

To cite this article: C. Maheswari *et al* 2018 *J. Phys.: Conf. Ser.* **1139** 012065

View the [article online](#) for updates and enhancements.



**IOP | ebooks™**

Bringing you innovative digital publishing with leading voices to create your essential collection of books in STEM research.

Start exploring the collection - download the first chapter of every title for free.

# Neutrosophic generalized b-closed sets in Neutrosophic topological spaces

C.Maheswari<sup>1</sup>, M. Sathyabama<sup>2</sup>, S.Chandrasekar<sup>3</sup>

<sup>1</sup> Department of Mathematics, Muthayammal College of Arts and Science, Rasipuram, Namakkal(DT), Tamil Nadu, India.

<sup>2</sup> Department of Mathematics, Periyar University Constituent College of Arts and Science, Idappadi, Salem(DT), Tamilnadu, India.

<sup>3</sup> PG & Research Department of Mathematics, Arignar Anna Government Arts College, Namakkal(DT), Tamil Nadu, India

Email: mahi2gobi@gmail.com<sup>1</sup>, sathyachezian@gmail.com<sup>2</sup>, chandrumat@gmail.com<sup>3</sup>

**Abstract.** Smarandache introduced and developed the new concept of Neutrosophic set from the Intuitionistic fuzzy sets. A.A. Salama introduced Neutrosophic topological spaces by using the Neutrosophic crisp sets. Aim of this paper is we introduce and study about Neutrosophic generalized b closed sets in Neutrosophic topological spaces and its properties are discussed details.

## 1. Introduction

Topology is a classical subject, as a generalization topological spaces many type of topological spaces introduced over the year. C.L. Chang[3] was introduced and developed fuzzy topological space by using L.A. Zadeh's[12] fuzzy sets. Coker[4] introduced the concepts of Intuitionistic fuzzy topological spaces by using Atanassov's[1] Intuitionistic fuzzy set Neutrality the degree of indeterminacy, as an independent concept, was introduced by Smarandache [6] in 1998. He also defined the Neutrosophic set on three component Neutrosophic topological spaces (T- Truth, F -Falsehood ,I- Indeterminacy). Neutrosophic topological spaces(N-T-S) introduced by Salama [10] et al. In 1996 D. Andrijevic [2] introduced b open sets in topological space, R.Dhavaseelan[5], Saied Jafari are introduced Neutrosophic generalized closed sets. Aim of this paper is we introduced in Neutrosophic b-open sets, Neutrosophic generalized b-open sets in Neutrosophic topological space and also discussed about properties of Neutrosophic gb-interior and Neutrosophic gb-closure in Neutrosophic topological spaces(N-T-S)

## 2. Preliminaries

In the Second section, we recall needed basic definition and operation of Neutrosophic sets and then fundamental results

*Definition 2.1 [10]*

Let X be a non-empty fixed set. A Neutrosophic set P is an object having the form

$$P = \{ \langle x, \mu_P(x), \sigma_P(x), \gamma_P(x) \rangle : x \in X \}$$

where  $\mu_P(x)$ -represents the degree of membership function,

$\sigma_P(x)$ - represents degree indeterminacy and then

$\gamma_P(x)$ - represents the degree of non-membership function

*Remark 2.2 [10]*

Neutrosophic set  $P = \{ \langle x, \mu_P(x), \sigma_P(x), \gamma_P(x) \rangle : x \in X \}$  can be write to an ordered triple lies in the interval in  $] -0, 1+ [$  on X.

*Remark 2.3 [10]*

we shall use the symbol

Neutrosophic set  $P = \{ \langle x, \mu_P(x), \sigma_P(x), \gamma_P(x) \rangle : x \in X \}$  we can be write briefly

Like as  $P = \langle x, \mu_P, \sigma_P, \gamma_P \rangle$



*Definition 2.4 [10]*

In N-T-S,  $0_N$  may be defined like as:  $\forall x \in X$

$$0_1 = \langle x, 0, 0, 1 \rangle$$

$$0_2 = \langle x, 0, 1, 1 \rangle$$

$$0_3 = \langle x, 0, 1, 0 \rangle$$

$$0_4 = \langle x, 0, 0, 0 \rangle$$

$1_N$  may be defined like as:  $\forall x \in X$

$$1_1 = \langle x, 1, 0, 0 \rangle$$

$$1_2 = \langle x, 1, 0, 1 \rangle$$

$$1_3 = \langle x, 1, 1, 0 \rangle$$

$$1_4 = \langle x, 1, 1, 1 \rangle$$

*Definition 2.5 [10]*

Neutrosophic set  $P = \{ \langle x, \mu_P(x), \sigma_P(x), \gamma_P(x) \rangle \}$  on  $X$  and  $\forall x \in X$

then complement of  $P$  is

$$P^c = \{ \langle x, \gamma_P(x), 1 - \sigma_P(x), \mu_P(x) \rangle \}$$

*Definition 2.6 [10]*

Let  $P$  and  $Q$  are two Neutrosophic sets  $\forall x \in X$

$$P = \{ \langle x, \mu_P(x), \sigma_P(x), \gamma_P(x) \rangle \}$$
 and

$$Q = \{ \langle x, \mu_Q(x), \sigma_Q(x), \gamma_Q(x) \rangle \}.$$

Then

$$P \subseteq Q \Leftrightarrow \mu_P(x) \leq \mu_Q(x), \sigma_P(x) \leq \sigma_Q(x) \text{ and } \gamma_P(x) \geq \gamma_Q(x)$$

*Proposition 2.6 [10]*

The following results are true for any Neutrosophic set  $P$

$$(i) 0_N \subseteq P, 0_N \subseteq 0_N$$

$$(ii) P \subseteq 1_N, 1_N \subseteq 1_N$$

*Definition 2.7 [10]*

Let  $X$  be a non-empty set, and

Let  $P$  and  $Q$  be two Neutrosophic sets are

$$P = \langle x, \mu_P(x), \sigma_P(x), \gamma_P(x) \rangle,$$

$$Q = \langle x, \mu_Q(x), \sigma_Q(x), \gamma_Q(x) \rangle$$
 Then

$$(i) P \cap Q = \langle x, \mu_P(x) \wedge \mu_Q(x), \sigma_P(x) \vee \sigma_Q(x) \& \gamma_P(x) \vee \gamma_Q(x) \rangle$$

$$(ii) P \cup Q = \langle x, \mu_P(x) \vee \mu_Q(x), \sigma_P(x) \wedge \sigma_Q(x) \& \gamma_P(x) \wedge \gamma_Q(x) \rangle$$

*Proposition 2.8 [10]*

The following conditions are true for all two Neutrosophic sets  $P$  and  $Q$  are

$$(i) (P \cap Q)^c = P^c \cup Q^c$$

$$(ii) (P \cup Q)^c = P^c \cap Q^c.$$

*Definition 2.9 [10]*

Let  $X$  be non-empty set and  $\tau_N$  be the collection of Neutrosophic subsets of  $X$  satisfying the following properties :

$$(i) 0_N, 1_N \in \tau_N,$$

$$(ii) T_1 \cap T_2 \in \tau_N \text{ for any } T_1, T_2 \in \tau_N,$$

$$(iii) \cup T_i \in \tau_N \text{ for every } \{T_i : i \in J\} \subseteq \tau_N$$

Then the space  $(X, \tau_N)$  is called a Neutrosophic topological space(N-T-S).

The element of  $\tau_N$  are called Neu-OS (Neutrosophic open set)

and its complement is Neu-CS(Neutrosophic closed set)

*Example 2.10 [10]*

Let  $X = \{x\}$  and  $\forall x \in X$

$$A_1 = \langle x, 0.6, 0.6, 0.5 \rangle$$

$$A_2 = \langle x, 0.5, 0.7, 0.9 \rangle$$

$$A_3 = \langle x, 0.6, 0.7, 0.5 \rangle$$

$$A_4 = \langle x, 0.5, 0.6, 0.9 \rangle$$

Then the collection  $\tau_N = \{0_N, A_1, A_2, A_3, A_4, 1_N\}$  is called a N-T-S on X.

*Definition 2.11 [10]*

$(X, \tau_N)$  be N-T-S and  $\forall x \in X$

$P = \{\langle x, \mu_P(x), \sigma_P(x), \gamma_P(x) \rangle\}$  be a Neutrosophic set in X. Then the Neutrosophic closure and

Then the Neutrosophic closure of P is

$$\text{Neu-Cl}(P) = \bigcap \{ H : H \text{ is a Neutrosophic closed set in } X \text{ and } P \subseteq H \}$$

Neutrosophic interior of P is

$$\text{Neu-Int}(P) = \bigcup \{ M : M \text{ is a Neutrosophic open set in } X \text{ and } M \subseteq P \}.$$

Then

(i) P is Neutrosophic open set iff  $P = \text{Neu-Int}(P)$ .

(ii) P is Neutrosophic closed set iff  $P = \text{Neu-Cl}(P)$ .

*Proposition 2.12 [10]*

Let  $(X, \tau_N)$  be a Neutrosophic topological spaces, Then for any Neutrosophic set P

(i)  $\text{Neu-Cl}((P)^c) = (\text{Neu-Int}(P))^c$ ,

(ii)  $\text{Neu-Int}((P)^c) = (\text{Neu-Cl}(P))^c$ .

*Proposition 2.13 [10]*

Let P, Q be two Neutrosophic sets in N-T-S  $(X, \tau_N)$ . Then the following results are true:

(i)  $\text{Neu-Int}(P) \subseteq P$ ,

(ii)  $P \subseteq \text{Neu-Cl}(P)$ ,

(iii)  $P \subseteq Q \Rightarrow \text{Neu-Int}(P) \subseteq \text{Neu-Int}(Q)$ ,

(iv)  $P \subseteq Q \Rightarrow \text{Neu-Cl}(P) \subseteq \text{Neu-Cl}(Q)$ ,

(v)  $\text{Neu-Int}(\text{Neu-Int}(P)) = \text{Neu-Int}(P)$ ,

(vi)  $\text{Neu-Cl}(\text{Neu-Cl}(P)) = \text{Neu-Cl}(P)$ ,

(vii)  $\text{Neu-Int}(P \cap Q) = \text{Neu-Int}(P) \cap \text{Neu-Int}(Q)$ ,

(viii)  $\text{Neu-Cl}(P \cup Q) = \text{Neu-Cl}(P) \cup \text{Neu-Cl}(Q)$ ,

(ix)  $\text{Neu-Int}(0_N) = 0_N$ ,

(x)  $\text{Neu-Int}(1_N) = 1_N$ ,

(xi)  $\text{Neu-Cl}(0_N) = 0_N$ ,

(xii)  $\text{Neu-Cl}(1_N) = 1_N$ ,

(xiii)  $P \subseteq Q \Rightarrow Q^c \subseteq P^c$ ,

(xiv)  $\text{Neu-Cl}(P \cap Q) \subseteq \text{Neu-Cl}(P) \cap \text{Neu-Cl}(Q)$ ,

(xv)  $\text{Neu-Int}(P \cup Q) \supseteq \text{Neu-Int}(P) \cup \text{Neu-Int}(Q)$ .

*Definition: 2.14 [5]*

Neutrosophic generalized closed set (Neu-g closed) if  $\text{Neutrosophic cl}(P) \subseteq G$  whenever  $P \subseteq G$  and G is Neutrosophic open set in  $(X, \tau_N)$ .

### 3. Neutrosophic generalized b-open sets

For this third section, we are newly introduce and study the new concept of Neutrosophic generalized b-open sets in N-T-S

*Definition: 3.1*

Let  $(X, \tau_N)$  be a N-T-S. A Neutrosophic set P is called

Neutrosophic b-open set is

$$\text{if } P \subseteq \text{Neu-cl} [\text{Neu-int}(P)] \cup \text{Neu-int}[\text{Neu-cl}(P)]$$

Neutrosophic b-closed set is

$$\text{Neu-cl} [\text{Neu-int}(P)] \cap \text{Neu-int}[\text{Neu-cl}(P)] \subseteq P$$

*Definition: 3.2*

Neutrosophic generalized b-closed Set (Neu-gb-closed set) if  $\text{Neutrosophic-bcl}(P) \subseteq G$  whenever  $P \subseteq G$  and G is Neutrosophic open set in  $(X, \tau_N)$ .

*Theorem 3.3.*

For Every Neutrosophic open sets is Neutrosophic generalized b-open sets.

*Proof.*

Now Let  $P$  is a Neu-OS in N-T-S  $(X, \tau_N)$  since  $P \subseteq \text{Neu-cl}(P)$  and  $P = \text{Neu-Int}(P)$ ,  $\text{Neu-Int}(P) \subseteq \text{Neu-Int}(\text{Neu-cl}(P))$  and then  $\text{Neu-Int}(P) \subseteq \text{Neu-cl}(\text{Neu-Int}(P))$  which implies  $\text{Neu-Int}(P) \subseteq \text{Neu-cl}(\text{Neu-Int}(P) \cup \text{Neu-Int}(\text{Neu-cl}(P)))$ . Hence  $P \subseteq \text{Neu-Int}(P) \subseteq \text{Neu-cl}(\text{Neu-Int}(P) \cup \text{Neu-Int}(\text{Neu-cl}(P)))$  and  $P$  is Neu- gb-open in  $(X, \tau_N)$ .

But the converse of this theorem is fails

i.e., For Every Neu-gbOS is not Neutrosophic open sets.

*Example 3.4*

Here  $X = \{a, b, c\}$  with  $\tau_N = \{0_N, A_1, A_2, 1_N\}$  and  $(\tau_N)^c = \{1_N, A_3, A_4, 0_N\}$  where

$A_1 = \{(0.6, 0.6, 0.4), (0.2, 0.7, 1), (1, 0.6, 0.5)\}$

$A_2 = \{(0.1, 0.4, 0.8), (0.2, 0.6, 1), (0.6, 0.5, 0.9)\}$

$A_3 = \{(0.4, 0.4, 0.6), (1, 0.3, 0.2), (0.5, 0.4, 1)\}$

$A_4 = \{(0.8, 0.6, 0.1), (1, 0.4, 0.2), (0.9, 0.5, 0.6)\}$ .

$A_5 = \{(0.3, 0.4, 1), (0.1, 0.2, 1), (0.4, 0.2, 1)\}$ .

Here the Neu-gbOSs are  $A_3, A_4$  and  $A_5$ .

Also  $A_5$  is Neu-gbCS and  $A_5$  is not Neu-CS.

*Theorem 3.5*

Consider if  $P$  and  $Q$  are Neu-gbCS, and then  $P \cup Q$  is Neu-gbCS.

*Proof:*

If  $P \cup Q \subseteq K$  and  $K$  is Neutrosophic open set, then  $P \subseteq K$  and  $Q \subseteq K$ . Since  $P$  and  $Q$  are Neu-gb closed sets,  $\text{Neu-cl}(P) \subseteq K$  and  $\text{Neu-cl}(Q) \subseteq K$  and hence  $\text{Neu-cl}(P) \cup \text{Neu-cl}(Q) \subseteq K$ . This implies  $\text{Neu-cl}(P \cup Q) \subseteq K$ . Thus  $P \cup Q$  is Neu-gbCS in  $X$ .

*Theorem 3.6*

Let  $P$  is a Neu-gb closed set and then  $\text{Neu-cl}(P) - P \not\subseteq$  any nonempty Neu-C-S.

*Proof:*

Let  $P$  is a Neu-gbCS. Let  $G$  be a Neu-CS subset of  $\text{Neu-cl}(P) - P$ . Then  $P \subseteq G^c$ . But  $P$  is Neu-gbCS. Therefore  $\text{Neu-cl}(P) \subseteq G^c$ . Consequently  $G \subseteq (\text{Neu-cl}(P))^c$ . We have  $G \subseteq \text{Neu-cl}(P)$ . Thus  $G \subseteq \text{Neu-cl}(P) \cap (\text{Neu-cl}(P))^c = \emptyset$ . Hence  $G$  is empty.

#### 4. Neutrosophic generalized b interior in a N-T-S

In this Fourth section, we newly introduce and study about the properties of Neu- gb interior in a N-T-S.

*Definition: 4.1*

Let  $(X, \tau_N)$  be a Neutrosophic topological space and  $P$  be a Neutrosophic set in  $X$ , then the Neu-gb-interior of  $P$  is defined as

$\text{Neu-gb-int}(P) = \cup \{M/M \text{ is a Neu-gbOS in } X \text{ and } M \subseteq P\}$

*Theorem: 4.2*

Neutrosophic subsets  $P$  and  $Q$  of a N-T-S  $X$  we have

(i)  $\text{Neu-gb-Int}(P) \subseteq P$

(ii)  $P$  is Neu-gb-open set in  $X \Leftrightarrow \text{Neu-gb-Int}(P) = P$

(iii)  $\text{Neu-gb-Int}(\text{Neu-gb-Int}(P)) = \text{Neu-gb-Int}(P)$

(iv) If  $P \subseteq Q$  then  $\text{Neu-gb-Int}(P) \subseteq \text{Neu-gb-Int}(Q)$

*Proof:*

Proof of (i) is directly get the result through the Definition 4.1.

Let  $P$  be Neu-gb-open set in  $X$ . Then  $P \subseteq \text{Neu-gb-Int}(P)$ . from 4.2(i) we obtain the result  $P = \text{Neu-gb-Int}(P)$ . Now Conversely we assume that  $P = \text{Neu-gb-Int}(P)$ . From the Definition 4.1, Neutrosophic set  $P$  is a Neu-gb-open set in N-T-S  $X$ . from this we get the result (ii). From the result (ii),  $\text{Neu-gb-Int}(\text{Neu-gb-Int}(P)) = \text{Neu-gb-Int}(P)$ . we get the result (iii). Since  $P \subseteq Q$ , by using (i),  $\text{Neu-gb-Int}(P) \subseteq \text{Neu-gb-Int}(Q)$ . i.e.,  $\text{Neu-gb-Int}(P) \subseteq Q$ . from the result (iii),  $\text{Neu-gb-Int}(\text{Neu-gb-Int}(P)) \subseteq \text{Neu-gb-Int}(Q)$ . Thus  $\text{Neu-gb-Int}(P) \subseteq \text{Neu-gb-Int}(Q)$ . we get the result (iv).

*Theorem 4.3*

Let  $P$  and  $Q$  are two Neutrosophic subsets of  $N$ -T-S  $(X, \tau_N)$  then

(i)  $\text{Neu-gb-Int}(P \cap Q) = \text{Neu-gb-Int}(P) \cap \text{Neu-gb-Int}(Q)$

(ii)  $\text{Neu-gb-Int}(P \cup Q) \supseteq \text{Neu-gb-Int}(P) \cup \text{Neu-gb-Int}(Q)$ .

*Proof :*

Since  $P \cap Q \subseteq P$  and  $P \cap Q \subseteq Q$ , follows from the theorem 4.2(iv),  $\text{Neu-gb-Int}(P \cap Q) \subseteq \text{Neu-gb-Int}(P)$  and  $\text{Neu-gb-Int}(P \cap Q) \subseteq \text{Neu-gb-Int}(Q)$ . This implies that  $\text{Neu-gb-Int}(P \cap Q) \subseteq \text{Neu-gb-Int}(P) \cap \text{Neu-gb-Int}(Q)$  .....(1). follows from the theorem 3.4(i),  $\text{Neu-gb-Int}(P) \subseteq P$  and  $\text{Neu-gb-Int}(Q) \subseteq Q$ . This implies that  $\text{Neu-gb-Int}(P) \cap \text{Neu-gb-Int}(Q) \subseteq P \cap Q$ . Now from theorem 4.2(iv),  $\text{Neu-gb-Int}((\text{Neu-gb-Int}(P) \cap \text{Neu-gb-Int}(Q))) \subseteq \text{Neu-gb-Int}(P \cap Q)$ . By (1),  $\text{Neu-gb-Int}(\text{Neu-gb-Int}(P)) \cap \text{Neu-gb-Int}(\text{Neu-gb-Int}(Q)) \subseteq \text{Neu-gb-Int}(P \cap Q)$ . from theorem 4.2(iii),  $\text{Neu-gb-Int}(P) \cap \text{Neu-gb-Int}(Q) \subseteq \text{Neu-gb-Int}(P \cap Q)$  .....(2). From (1) and (2),  $\text{Neu-gb-Int}(P \cap Q) = \text{Neu-gb-Int}(P) \cap \text{Neu-gb-Int}(Q)$ . This implies (i). Since  $P \subseteq P \cup Q$  and  $Q \subseteq P \cup Q$ , by from theorem 4.2(iv),  $\text{Neu-gb-Int}(P) \subseteq \text{Neu-gb-Int}(P \cup Q)$  and  $\text{Neu-gb-Int}(Q) \subseteq \text{Neu-gb-Int}(P \cup Q)$ . This implies that  $\text{Neu-gb-Int}(P) \cup \text{Neu-gb-Int}(Q) \subseteq \text{Neu-gb-Int}(P \cup Q)$ . Hence (ii).

Converse part of Theorem 4.3(ii) is need not be true

*Example 4.4*

Let  $X = \{p, q, r\}$  and  $\tau_N = \{0_N, A_1, A_2, A_3, A_4, 1_N\}$  where  $\tau_N$  is a Neutrosophic topology in  $N$ -T-S

$A_1 = \langle (0.5, 0.7, 0.2), (0.6, 0.6, 0.3), (1, 0.7, 0.4) \rangle$ ,

$A_2 = \langle (0.5, 0.6, 0.2), (0.8, 0.7, 0.3), (1, 0.5, 0.2) \rangle$ ,

$A_3 = \langle (0.5, 0.7, 0.2), (0.8, 0.7, 0.3), (1, 0.7, 0.2) \rangle$ ,

$A_4 = \langle (0.5, 0.6, 0.2), (0.6, 0.6, 0.3), (1, 0.5, 0.4) \rangle$ .

$\tau_N$  is a Neutrosophic topology in  $N$ -T-S

Consider the Neutrosophic sets

$A_5 = \langle (0.8, 0.6, 0.2), (0.8, 0.6, 0.2), (1, 0.5, 0.1) \rangle$  and

$A_6 = \langle (0.5, 0.6, 0.2), (0.6, 0.7, 0.3), (1, 0.7, 0.2) \rangle$ .

Then  $\text{Neu-gbint}(A_5) = A_4$  and  $\text{Neu-gbint}(A_6) = A_4$ .

This implies that  $\text{Neu-gbint}(A_5) \cup \text{Neu-gbint}(A_6) = A_4$ . Then

$A_5 \cup A_6 = \langle (0.8, 0.6, 0.2), (0.8, 0.7, 0.2), (1, 0.7, 0.1) \rangle$ ,

it follows that  $\text{Neu-gbint}(A_5 \cup A_6) = A_2$ . Then  $\text{Neu-gbint}(A_5 \cup A_6) \not\subseteq \text{Neu-gbint}(A_5) \cup \text{Neu-gbint}(A_6)$ .

**5. Neutrosophic generalized b-closure in N-T-S.**

Now In the fifth section, we newly introduce and study the properties and characterization of Neu- gb-closure in  $N$ -T-S.

*Definition 5.1*

Let  $P$  is a Neutrosophic subset  $P$  of Neutrosophic topological space  $(X, \tau_N)$

Neu- gb-closure defined as

$\text{Neu-gb-Cl}(P) = \cap \{H: H \text{ is a Neu-gb-closed set in } X \text{ and } H \supseteq P\}$ .

*Theorem 5.2*

Let  $P$  is a Neutrosophic subset of  $N$ -T-S  $(X, \tau_N)$

(i)  $[(\text{Neu-gb-Int}(P))^c] = \text{Neu-gb-Cl}[(P)^c]$ ,

(ii)  $[\text{Neu-gb-Cl}(P)]^c = \text{Neu-gb-Int}[(P)^c]$ .

*Proof :*

From the Definition 5.1,  $\text{Neu-gb-Int}(P) = \cup \{M: M \text{ is a Neu-gb-open set in } X \text{ and } M \subseteq P\}$ . Take complement each both sides,  $[(\text{Neu-gb-Int}(P))^c] = (\cup \{M: M \text{ is a Neu-gb open set in } X \text{ and } M \subseteq P\})^c = \cap \{M^c: M^c \text{ is a Neu-gb-closed set in } X \text{ and } [(P)^c] \subseteq M^c\}$ . Replacing  $M^c$  by  $H$ , we get  $[(\text{Neu-gb-Int}(P))^c] = \cap \{H: H \text{ is a Neu-gb-closed set in } X \text{ and } H \supseteq [(P)^c]\}$ . From the Definition 5.1,  $[(\text{Neu-gb-Int}(P))^c] = \text{Neu-gb-Cl}([(P)^c])$ . This proves (i). By using (i),  $[\text{Neu-gb-Int}((P)^c)]^c = \text{Neu-gb-Cl}[(P)^c]^c = \text{Neu-gb-Cl}(P)$ . Take complement each both sides, Then we obtain  $\text{Neu-gb-Int}((P^c)) = [\text{Neu-gb-Cl}(P)]^c$ . we obtained result(ii).

*Theorem 5.3*

If  $P$  and  $Q$  are Neutrosophic subset of N-T-S  $(X, \tau_N)$ , Then

- (i)  $P \subseteq \text{Neu-gb-Cl}(P)$
- (ii)  $P$  is Neu-gb-CS in  $X \Leftrightarrow \text{Neu-gb-Cl}(P) = P$
- (iii)  $\text{Neu-gb-Cl}(\text{Neu-gb-Cl}(P)) = \text{Neu-gb-Cl}(P)$
- (iv) Now, If  $P \subseteq Q$  and then  $\text{Neu-gb-Cl}(P) \subseteq \text{Neu-gb-Cl}(Q)$

*Proof :*

(i) We can easily get result from Definition 5.1.

Let  $P$  be Neu-gb-closed set in  $X$ . From the theorem 5.3,  $P^c$  is Neu-gb-open set in  $X$ . From the theorem 5.2(ii),  $\text{Neu-gb-Int}((P)^c) = (P)^c \Leftrightarrow [\text{Neu-gb-Cl}(P)]^c = P^c \Leftrightarrow \text{Neu-gb-Cl}(P) = P$ . we obtain the result(ii). By using(ii),  $\text{Neu-gb-Cl}(\text{Neu-gb-Cl}(P)) = \text{Neu-gb-Cl}(P)$ . we obtain the result (iii). Since  $P \subseteq Q, Q^c \subseteq P^c$ . From the theorem 4.2(iv),  $\text{Neu-gb-Int}((Q)^c) \subseteq \text{Neu-gb-Int}((P)^c)$ . apply complement each sides,  $[\text{Neu-gb-Int}((Q)^c)]^c \supseteq [\text{Neu-gb-Int}((P)^c)]^c$ . From the theorem 5.2(ii),  $\text{Neu-gb-Cl}(P) \subseteq \text{Neu-gb-Cl}(Q)$ . we obtain the result (iv).

*Theorem 5.4*

Let  $P$  be a Neutrosophic set in a N-T-S  $(X, \tau_N)$ . Then  $\text{Neu-Int}(P) \subseteq \text{Neu-gb-Int}(P) \subseteq P \subseteq \text{Neu-gb-Cl}(P) \subseteq \text{Neu-Cl}(P)$ .

*Proof :*

We can easily get result from Definition 5.1.

*Theorem 5.5*

If  $P$  and  $Q$  are Neutrosophic subset of N-T-S  $(X, \tau_N)$ , Then

- (i)  $\text{Neu-gb-Cl}(P \cup Q) = \text{Neu-gb-Cl}(P) \cup \text{Neu-gb-Cl}(Q)$  and
- (ii)  $\text{Neu-gb-Cl}(P \cap Q) \subseteq \text{Neu-gb-Cl}(P) \cap \text{Neu-gb-Cl}(Q)$ .

*Proof :*

Since  $\text{Neu-gb-Cl}(P \cup Q) = \text{Neu-gb-Cl}((P \cup Q)^c)^c$  By From theorem 5.2(i),  $\text{Neu-gb-Cl}(P \cup Q) = [\text{Neu-gb-Int}((P \cup Q)^c)]^c = [\text{Neu-gb-Int}(P^c \cap Q^c)]^c$ . once Again From theorem 3.5(i),  $\text{Neu-gb-Cl}(P \cup Q) = [\text{Neu-gb-Int}(P^c) \cap \text{Neu-gb-Int}(Q^c)]^c = [\text{Neu-gb-Int}(P^c)]^c \cup [\text{Neu-gb-Int}(Q^c)]^c$ . From theorem 5.2(i),  $\text{Neu-gb-Cl}(P \cup Q) = \text{Neu-gb-Cl}(P^c)^c \cup \text{Neu-gb-Cl}(Q^c)^c = \text{Neu-gb-Cl}(P) \cup \text{Neu-gb-Cl}(Q)$ . Thus proved(i). Since  $P \cap Q \subseteq P$  and  $P \cap Q \subseteq Q$ , From theorem 5.3(iv),  $\text{Neu-gb-Cl}(P \cap Q) \subseteq \text{Neu-gb-Cl}(P)$  and  $\text{Neu-gb-Cl}(P \cap Q) \subseteq \text{Neu-gb-Cl}(Q)$ . This implies that  $\text{Neu-gb-Cl}(P \cap Q) \subseteq \text{Neu-gb-Cl}(P) \cap \text{Neu-gb-Cl}(Q)$ . we obtain the result (ii).

Converse of (ii) is not true,  $\text{Neu-gb-Cl}(P) \cap \text{Neu-gb-Cl}(Q) \not\subseteq \text{Neu-gb-Cl}(P \cap Q)$

*Example 5.6*

$\text{Neu-gb-Cl}(P) \cap \text{Neu-gb-Cl}(Q) \not\subseteq \text{Neu-gb-Cl}(P \cap Q)$

Let  $X = \{p, q, r\}$  with  $\tau_N = \{0_N, A_1, A_2, A_3, A_4, 1_N\}$  and  $(\tau_N)^c = \{1_N, A_5, A_6, A_7, A_8, 0_N\}$  where

$A_1 = \langle (0.6, 0.6, 0.2), (0.7, 0.7, 0.2), (1, 0.5, 0.3) \rangle$

$A_2 = \langle (0.5, 0.5, 0.3), (0.9, 0.6, 0.4), (1, 0.7, 0.4) \rangle$

$A_3 = \langle (0.5, 0.5, 0.3), (0.7, 0.6, 0.4), (1, 0.5, 0.4) \rangle$

$A_4 = \langle (0.6, 0.6, 0.2), (0.9, 0.7, 0.2), (1, 0.7, 0.3) \rangle$

$A_5 = \langle (0.2, 0.4, 0.6), (0.2, 0.3, 0.7), (0.3, 0.5, 1) \rangle$ ,

$A_6 = \langle (0.3, 0.5, 0.5), (0.4, 0.4, 0.9), (0.4, 0.3, 1) \rangle$ ,

$A_7 = \langle (0.3, 0.5, 0.5), (0.4, 0.4, 0.7), (0.4, 0.5, 1) \rangle$ ,

$A_8 = \langle (0.2, 0.4, 0.6), (0.2, 0.3, 0.9), (0.3, 0.3, 1) \rangle$ .

Then  $(X, \tau_N)$  is a N-T-S.

Here we consider the some Neutrosophic sets

$A_9 = \langle (0.2, 0.2, 0.6), (0.3, 0.3, 0.8), (0.4, 0.3, 1) \rangle$  and

$A_{10} = \langle (0.3, 0.4, 0.9), (0.2, 0.2, 0.9), (0.3, 0.5, 1) \rangle$ .

Then  $\text{Neu-gbcl}(A_9) = A_7$  and  $\text{Neu-gbcl}(A_{10}) = A_7$ .

This implies that  $\text{Neu-gbcl}(A_9) \cap \text{Neu-gbcl}(A_{10}) = A_7$ .

Now,  $A_9 \cap A_{10} = \langle (0.2, 0.2, 0.9), (0.2, 0.2, 0.9), (0.3, 0.3, 1) \rangle$ , it follows that  $\text{Neu-gbcl}(A_9 \cap A_{10}) = A_8$ .

Then  $\text{Neu-gbcl}(A_9) \cap \text{Neu-gbcl}(A_{10}) \not\subseteq \text{Neu-gbcl}(A_9 \cap A_{10})$ .

*Theorem 5.7*

If  $P$  and  $Q$  are Neutrosophic subset of  $N-T-S (X, \tau_N)$  then

- (i)  $\text{Neu-gb-Cl}(P) \supseteq P \cup \text{Neu-gb-Cl}(\text{Neu-gb-Int}(P))$ ,
- (ii)  $\text{Neu-gb-Int}(P) \subseteq P \cap \text{Neu-gb-Int}(\text{Neu-gb-Cl}(P))$ ,
- (iii)  $\text{Neu-Int}(\text{Neu-gb-Cl}(P)) \subseteq \text{Neu-Int}(\text{Neu-Cl}(P))$ ,
- (iv)  $\text{Neu-Int}(\text{Neu-gb-Cl}(P)) \supseteq \text{Neu-Int}(\text{Neu-gb-Cl}(\text{Neu-gb-Int}(P)))$ .

*Proof :*

From theorem 5.3(i),  $P \subseteq \text{Neu-gb-Cl}(P)$  .....(1). We use theorem 3.4(i),  $\text{Neu-gb-Int}(P) \subseteq P$ . Then  $\text{Neu-gb-Cl}(\text{Neu-gb-Int}(P)) \subseteq \text{Neu-gb-Cl}(P)$  .....(2). From (1) & (2) we have,  $P \cup \text{Neu-gb-Cl}(\text{Neu-gb-Int}(P)) \subseteq \text{Neu-gb-Cl}(P)$ . we obtain result (i). From theorem 4.2(i),  $\text{Neu-gb-Int}(P) \subseteq P$ .....(3). We get result from theorem 5.3(i),  $P \subseteq \text{Neu-gb-Cl}(P)$ . Then  $\text{Neu-gb-Int}(P) \subseteq \text{Neu-gb-Int}(\text{Neu-gb-Cl}(P))$ .....(4). From (3) & (4), we have  $\text{Neu-gb-Int}(P) \subseteq P \cap \text{Neu-gb-Int}(\text{Neu-gb-Cl}(P))$ . We obtain (ii). From theorem 5.4,  $\text{Neu-gb-Cl}(P) \subseteq \text{Neu-Cl}(P)$ . We obtain  $\text{Neu-Int}(\text{Neu-gb-Cl}(P)) \subseteq \text{Neu-Int}(\text{Neu-Cl}(P))$ . Hence (iii). By (i),  $\text{Neu-gb-Cl}(P) \supseteq P \cup \text{Neu-gb-Cl}(\text{Neu-gb-Int}(P))$ . We have  $\text{Neu-Int}(\text{Neu-gb-Cl}(P)) \supseteq \text{Neu-Int}(P \cup \text{Neu-gb-Cl}(\text{Neu-gb-Int}(P)))$ . Since  $\text{Neu-Int}(P \cup Q) \supseteq \text{Neu-Int}(P) \cup \text{Neu-Int}(Q)$ ,  $\text{Neu-Int}(\text{Neu-gb-Cl}(P)) \supseteq \text{Neu-Int}(P) \cup \text{Neu-Int}(\text{Neu-gb-Cl}(\text{Neu-gb-Int}(P))) \supseteq \text{Neu-Int}(\text{Neu-gb-Cl}(\text{Neu-gb-Int}(P)))$ . Hence (iv).

## References

- [1] Atanassov K 1986 Intuitionistic fuzzy sets, *Fuzzy Sets and Systems* **20** pp 87-94
- [2] Andrijevic D 1996 b open sets, *Math. Vesnik* **48** 1 pp 59-64
- [3] Chang C L 1968 Fuzzy Topological Spaces, *J. Math. Anal. Appl.* **24** pp 182-190
- [4] Dogan Coker 1997 An introduction to Intuitionistic fuzzy topological spaces *Fuzzy Sets and Systems* **88** pp 81-89
- [5] Dhavaseelan R and Jafari S 2018 Generalized Neutrosophic closed sets *New trends in Neutrosophic theory and applications II* pp 261-273
- [6] Florentin Smarandache 2002 Neutrosophic and Neutrosophic Logic First International Conference on Neutrosophy , Neutrosophic Logic, Set, Probability, and Statistics *University of New Mexico, Gallup, NM 87301, USA*, smarand@unm.edu
- [7] Floretin Smaradache 2010 Neutrosophic Set A Generalization of Intuitionistic Fuzzy set, *Journal of Defense Resources Management.* **1** pp 107-114
- [8] Iswarya P and Bageerathi K 2016 On Neutrosophic semi-open sets in Neutrosophic topological spaces, *International Journal of Mathematics Trends and Technology* **37** 3 pp 24-33
- [9] Salama A A and Alblowi S A 2012 Generalized Neutrosophic set and generalized Neutrosophic topological spaces *Journal computer Sci. Engineering* **2** 7 pp 129-132
- [10] Salama A A and Alblowi S A 2012 Neutrosophic set and Neutrosophic topological space, *ISOR J.mathematics*, **3** 4 pp 31-35
- [11] Shanthi V K Chandrasekar.S , Safina Begam.K, 2018 Neutrosophic Generalized Semi Closed Sets In Neutrosophic Topological Spaces *International Journal of Research in Advent technology* **6** 7 pp 1739-1743
- [12] Zadeh L P 1965 Fuzzy Sets, *Inform and Control* **8** pp 338- 353