

Novel multiple criteria decision making methods based on bipolar neutrosophic sets and bipolar neutrosophic graphs

Muhammad Akram

*Department of Mathematics,
University of the Punjab
New Campus, Lahore- Pakistan
e-mail: m.akram@puccit.edu.pk*

Musavarah Sarwar

*Department of Mathematics,
University of the Punjab
New Campus, Lahore- Pakistan
e-mail: musavarah656@gmail.com*

Abstract. In this research article, we present certain notions of bipolar neutrosophic graphs. We study the dominating and independent sets of bipolar neutrosophic graphs. We describe novel multiple criteria decision making methods based on bipolar neutrosophic sets and bipolar neutrosophic graphs. We develop an algorithm for computing domination in bipolar neutrosophic graphs. We also show that there are some flaws in Broumi et al. [11]’s definition of complement of a bipolar neutrosophic graph.

Keywords and phrases: Bipolar neutrosophic sets, Bipolar neutrosophic graphs, Domination number, Independent number, Decision making, Algorithm

2010 Mathematics Subject Classification: 03E72, 68R10, 68R05.

1. Introduction

A fuzzy set [27] is an important mathematical structure to represent a collection of objects whose boundary is vague. Fuzzy models are becoming useful because of their aim in reducing the differences between the traditional numerical models used in engineering and sciences and the symbolic models used in expert systems. In 1994, Zhang [29] introduced the notion of bipolar fuzzy sets and relations. Bipolar fuzzy sets are extension of fuzzy sets whose membership degree ranges $[-1, 1]$. The membership degree $(0, 1]$ indicates that the object satisfies a certain property whereas the membership degree $[-1, 0)$ indicates that the element satisfies the implicit counter property. Positive information represent what is considered to be possible and negative information represent what is granted to be impossible. Actually, a variety of decision making problems are based on two-sided bipolar

judgements on a positive side and a negative side. Nowadays bipolar fuzzy sets are playing a substantial role in chemistry, economics, computer science, engineering, medicine and decision making problems. Smarandache [22] introduced the idea of neutrosophic probability, sets and logic. Peng *et al.* [19], in 2014, described some operational properties and studied a new approach for multi-criteria decision making problems using neutrosophic sets. Ye [25, 26] discussed trapezoidal neutrosophic sets and simplified neutrosophic sets with applications in multi-criteria decision making problems. The other terminologies and applications of neutrosophic sets can be seen in [23, 25, 26]. In a neutrosophic set, the membership value is associated with truth, false and indeterminacy degrees but there is no restriction on their sum. Deli et al. [12] extended the ideas of bipolar fuzzy sets and neutrosophic sets to bipolar neutrosophic sets and studied its operations and applications in decision making problems.

Graph theory has numerous applications in science and engineering. However, in some cases, some aspects of graph theoretic concepts may be uncertain. In such cases, it is important to deal with uncertainty using the methods of fuzzy sets and logics. Based on Zadeh's fuzzy relations [28], Kaufmann [14] defined a fuzzy graph. The fuzzy relations between fuzzy sets were also considered by Rosenfeld [20] and he developed the structure of fuzzy graphs, obtaining analogs of several graph theoretical concepts. The complement of a fuzzy graph was defined by Mordeson [17]. The concept of strong arcs in fuzzy graphs was discussed in [10]. The theory of fuzzy graphs has extended widely by many researchers as it can be seen in [15, 17]. The idea of domination was first arose in chessboard problem in 1862. Somasundaram and Somasundaram [24] introduced domination and independent domination in fuzzy graphs. Gani and Chandrasekaran [18] studied the notion of fuzzy domination and independent domination using strong arcs. Akram [1, 2] introduced bipolar fuzzy graphs and discuss its various properties. Akram and Dudek [3] studied regular bipolar fuzzy graphs. Several new concepts on bipolar neutrosophic graphs and bipolar neutrosophic hypergraphs have been studied in [4-7]. In this research article, we present certain notions of bipolar neutrosophic graphs. We study the dominating and independent sets of bipolar neutrosophic graphs. We describe novel multiple criteria decision making methods based on bipolar neutrosophic sets and bipolar neutrosophic graphs. We develop an algorithm for computing domination in bipolar neutrosophic graphs. We also show that there are some flaws in Broumi et al. [11]'s definition.

3. Bipolar neutrosophic graphs

In this section, we discuss the concept of a bipolar neutrosophic graph and its various properties.

Definition 2.1. [23] A neutrosophic set C on a non-empty set Y is characterized by a truth membership function $t_C : Y \rightarrow [0, 1]$, indeterminacy membership function $I_C : Y \rightarrow [0, 1]$ and a falsity membership function $f_C : Y \rightarrow [0, 1]$. There is no restriction on the sum of $t_C(x)$, $I_C(x)$ and $f_C(x)$ for all $x \in Y$.

Definition 2.2. [12] A bipolar neutrosophic set on a empty set Y is an object of the form

$$C = \{(y, t_C^p(y), I_C^p(y), f_C^p(y), t_C^n(y), I_C^n(y), f_C^n(y)) : y \in Y\}$$

where, $t_C^p, I_C^p, f_C^p : Y \rightarrow [0, 1]$ and $t_C^n, I_C^n, f_C^n : Y \rightarrow [-1, 0]$. The positive values $t_C^p(y), I_C^p(y), f_C^p(y)$ denote respectively the truth, indeterminacy and false membership degrees of an element $y \in Y$ whereas $t_C^n(y), I_C^n(y), f_C^n(y)$ denote the implicit counter property of the truth, indeterminacy and false membership degrees of the element $y \in Y$ corresponding to the bipolar neutrosophic set C .

Definition 2.3. A bipolar neutrosophic relation on a non-empty set Y is a bipolar neutrosophic subset of $Y \times Y$ of the form $D = \{(yz, t_D^p(yz), I_D^p(yz), f_D^p(yz), t_D^n(yz), I_D^n(yz), f_D^n(yz)) : yz \in Y \times Y\}$ where, $t_D^p, I_D^p, f_D^p, t_D^n, I_D^n, f_D^n$ are defined by the mappings $t_D^p, I_D^p, f_D^p : Y \times Y \rightarrow [0, 1]$ and $t_D^n, I_D^n, f_D^n : Y \times Y \rightarrow [-1, 0]$.

Definition 2.4. A bipolar neutrosophic graph on a non-empty set Y is a pair $G = (C, D)$, where C is a bipolar neutrosophic set on Y and D is a bipolar neutrosophic relation in Y such that

$$\begin{aligned} t_D^p(yz) &\leq t_C^p(y) \wedge t_C^p(z), & I_D^p(yz) &\leq I_C^p(y) \wedge I_C^p(z), & f_D^p(yz) &\leq f_C^p(y) \vee f_C^p(z), \\ t_D^n(yz) &\geq t_C^n(y) \vee t_C^n(z), & I_D^n(yz) &\geq I_C^n(y) \vee I_C^n(z), & f_D^n(yz) &\geq f_C^n(y) \wedge f_C^n(z), \end{aligned}$$

for all $y, z \in Y$. Note that $D(yz) = (0, 0, 1, 0, 0, -1)$ for all $yz \in Y \times Y \setminus E$.

Example 2.1. Here we discuss an example of a bipolar neutrosophic graph such that $Y = \{x, y, z\}$. Let C be a bipolar neutrosophic set on Y given in Table. 1 and D be a bipolar neutrosophic relation in Y given in Table. 2.


Table 1: Bipolar neutrosophic set C Table 2: Bipolar neutrosophic relation D

	x	y	z
t_C^p	0.3	0.5	0.4
I_C^p	0.4	0.4	0.3
f_C^p	0.5	0.2	0.2
t_C^n	-0.6	-0.1	-0.5
I_C^n	-0.5	-0.8	-0.5
f_C^n	-0.2	-0.2	-0.5

	xy	yz	xz
t_D^p	0.3	0.3	0.3
I_D^p	0.4	0.3	0.3
f_D^p	0.5	0.2	0.5
t_D^n	-0.1	-0.1	-0.5
I_D^n	-0.5	-0.5	-0.5
f_D^n	-0.2	-0.5	-0.5

Routine calculations show that $G = (C, D)$ is a bipolar neutrosophic graph. The bipolar neutrosophic graph G is shown in Fig. 1.

Definition 2.5. Let $G_1 = (C_1, D_1)$ and $G_2 = (C_2, D_2)$ be two bipolar neutrosophic graphs where, C_1 and C_2 are bipolar neutrosophic sets on Y_1 and Y_2 , D_1 and D_2 are bipolar neutrosophic relations in Y_1 and Y_2 , respectively . The *union*

Figure 1: Bipolar neutrosophic graph G

of G_1 and G_2 is a pair $G_1 \cup G_2 = (C_1 \cup C_2, D_1 \cup D_2)$ such that for all $y, z \in Y$,

1. If $y \in Y_1, y \notin Y_2$ then, $(C_1 \cup C_2)(y) = C_1(y)$.
2. If $y \in Y_2, y \notin Y_1$ then, $(C_1 \cup C_2)(y) = C_2(y)$.
3. If $y \in Y_1 \cap Y_2$ then,

$$(C_1 \cup C_2)(y) = (t_{C_1}^p(y) \vee t_{C_2}^p(y), \frac{I_{C_1}^p(y) + I_{C_2}^p(y)}{2}, f_{C_1}^p(y) \wedge f_{C_2}^p(y), t_{C_1}^n(y) \wedge t_{C_2}^n(y), \frac{I_{C_1}^n(y) + I_{C_2}^n(y)}{2}, f_{C_1}^n(y) \vee f_{C_2}^n(y)),$$

If E_1 and E_2 are the sets of edges in G_1 and G_2 then, $D_1 \cup D_2$ can be defined as:

1. If $yz \in E_1, yz \notin E_2$ then, $(D_1 \cup D_2)(yz) = D_1(yz)$.
2. If $yz \in E_2, yz \notin E_1$ then, $(D_1 \cup D_2)(yz) = D_2(yz)$.
3. If $yz \in E_1 \cap E_2$ then,

$$(D_1 \cup D_2)(yz) = (t_{D_1}^p(yz) \vee t_{D_2}^p(yz), \frac{I_{D_1}^p(yz) + I_{D_2}^p(yz)}{2}, f_{D_1}^p(yz) \wedge f_{D_2}^p(yz), t_{D_1}^n(yz) \wedge t_{D_2}^n(yz), \frac{I_{D_1}^n(yz) + I_{D_2}^n(yz)}{2}, f_{D_1}^n(yz) \vee f_{D_2}^n(yz)).$$

Definition 2.6. The *intersection* of two bipolar neutrosophic graphs $G_1 = (C_1, D_1)$ and $G_2 = (C_2, D_2)$ is a pair $G_1 \cap G_2 = (C_1 \cap C_2, D_1 \cap D_2)$ where, C_1, C_2, D_1 and D_2 are given in Definition 2.5. The membership values of vertices and edges in $G_1 \cap G_2$ can be defined as,

$$(C_1 \cap C_2)(y) = (t_{C_1}^p(y) \wedge t_{C_2}^p(y), \frac{I_{C_1}^p(y) + I_{C_2}^p(y)}{2}, f_{C_1}^p(y) \vee f_{C_2}^p(y), t_{C_1}^n(y) \vee t_{C_2}^n(y), \frac{I_{C_1}^n(y) + I_{C_2}^n(y)}{2}, f_{C_1}^n(y) \wedge f_{C_2}^n(y)), \text{ for all } y \in Y_1 \cap Y_2.$$

$$(D_1 \cap D_2)(yz) = (t_{D_1}^p(yz) \wedge t_{D_2}^p(yz), \frac{I_{D_1}^p(yz) + I_{D_2}^p(yz)}{2}, f_{D_1}^p(yz) \vee f_{D_2}^p(yz), t_{D_1}^n(yz) \vee t_{D_2}^n(yz), \frac{I_{D_1}^n(yz) + I_{D_2}^n(yz)}{2}, f_{D_1}^n(yz) \wedge f_{D_2}^n(yz)),$$

for all $yz \in E_1 \cap E_2$.

Definition 2.7. The *join* of two bipolar neutrosophic graphs $G_1 = (C_1, D_1)$ and $G_2 = (C_2, D_2)$ is defined by the pair $G_1 + G_2 = (C_1 + C_2, D_1 + D_2)$ such that, $C_1 + C_2 = C_1 \cup C_2$, for all $y \in Y_1 \cup Y_2$, and the membership values of the edges in $G_1 + G_2$ are defined as,

1. $D_1 + D_2 = D_1 \cup D_2$, for all $yz \in E_1 \cup E_2$.
2. Let E' be the set of all edges joining the vertices of G_1 and G_2 then for all $yz \in E'$, where $y \in Y_1$ and $z \in Y_2$,

$$(D_1 + D_2)(yz) = (t_{D_1}^p(yz) \wedge t_{D_2}^p(yz), I_{D_1}^p(yz) \wedge I_{D_2}^p(yz), f_{D_1}^p(yz) \vee f_{D_2}^p(yz), \\ t_{D_1}^n(yz) \vee t_{D_2}^n(yz), I_{D_1}^n(yz) \vee I_{D_2}^n(yz), f_{D_1}^n(yz) \wedge f_{D_2}^n(yz)).$$

Definition 2.8. The *Cartesian product* of two bipolar neutrosophic graphs G_1 and G_2 is denoted by the pair $G_1 \square G_2 = (C_1 \square C_2, D_1 \square D_2)$ and defined as,

$$t_{C_1 \square C_2}^p(y) = t_{C_1}^p(y) \wedge t_{C_2}^p(y), \quad I_{C_1 \square C_2}^p(y) = I_{C_1}^p(y) \wedge I_{C_2}^p(y), \quad f_{C_1 \square C_2}^p(y) = f_{C_1}^p(y) \vee f_{C_2}^p(y), \\ t_{C_1 \square C_2}^n(y) = t_{C_1}^n(y) \vee t_{C_2}^n(y), \quad I_{C_1 \square C_2}^n(y) = I_{C_1}^n(y) \vee I_{C_2}^n(y), \quad f_{C_1 \square C_2}^n(y) = f_{C_1}^n(y) \wedge f_{C_2}^n(y).$$

for all $y \in Y_1 \times Y_2$. The membership values of the edges in $G_1 \square G_2$ can be calculated as,

1. $t_{D_1 \square D_2}^p((y_1, y_2)(y_1, z_2)) = t_{C_1}^p(y_1) \wedge t_{D_2}^p(y_2 z_2)$, $t_{D_1 \square D_2}^n((y_1, y_2)(y_1, z_2)) = t_{C_1}^n(y_1) \vee t_{D_2}^n(y_2 z_2)$, for all $y_1 \in Y_1, y_2 z_2 \in E_2$,
2. $t_{D_1 \square D_2}^p((y_1, y_2)(z_1, y_2)) = t_{D_1}^p(y_1 z_1) \wedge t_{C_2}^p(y_2)$, $t_{D_1 \square D_2}^n((y_1, y_2)(z_1, y_2)) = t_{D_1}^n(y_1 z_1) \vee t_{C_2}^n(y_2)$, for all $y_1 z_1 \in E_1, y_2 \in Y_2$,
3. $I_{D_1 \square D_2}^p((y_1, y_2)(y_1, z_2)) = I_{C_1}^p(y_1) \wedge I_{D_2}^p(y_2 z_2)$, $I_{D_1 \square D_2}^n((y_1, y_2)(y_1, z_2)) = I_{C_1}^n(y_1) \vee I_{D_2}^n(y_2 z_2)$, for all $y_1 \in Y_1, y_2 z_2 \in E_2$,
4. $I_{D_1 \square D_2}^p((y_1, y_2)(z_1, y_2)) = I_{D_1}^p(y_1 z_1) \wedge I_{C_2}^p(y_2)$, $I_{D_1 \square D_2}^n((y_1, y_2)(z_1, y_2)) = I_{D_1}^n(y_1 z_1) \vee I_{C_2}^n(y_2)$, for all $y_1 z_1 \in E_1, y_2 \in Y_2$,
5. $f_{D_1 \square D_2}^p((y_1, y_2)(y_1, z_2)) = f_{C_1}^p(y_1) \vee f_{D_2}^p(y_2 z_2)$, $f_{D_1 \square D_2}^n((y_1, y_2)(y_1, z_2)) = f_{C_1}^n(y_1) \wedge f_{D_2}^n(y_2 z_2)$, for all $y_1 \in Y_1, y_2 z_2 \in E_2$,
6. $f_{D_1 \square D_2}^p((y_1, y_2)(z_1, y_2)) = f_{D_1}^p(y_1 z_1) \vee f_{C_2}^p(y_2)$, $f_{D_1 \square D_2}^n((y_1, y_2)(z_1, y_2)) = f_{D_1}^n(y_1 z_1) \wedge f_{C_2}^n(y_2)$, for all $y_1 z_1 \in E_1, y_2 \in Y_2$.

Definition 2.9. The *direct product* of two bipolar neutrosophic graphs $G_1 = (C_1, D_1)$ and $G_2 = (C_2, D_2)$ is denoted by the pair $G_1 \times G_2 = (C_1 \times C_2, D_1 \times D_2)$ such that,

$$t_{C_1 \times C_2}^p(y) = t_{C_1}^p(y) \wedge t_{C_2}^p(y), \quad I_{C_1 \times C_2}^p(y) = I_{C_1}^p(y) \wedge I_{C_2}^p(y), \quad f_{C_1 \times C_2}^p(y) = f_{C_1}^p(y) \vee f_{C_2}^p(y), \\ t_{C_1 \times C_2}^n(y) = t_{C_1}^n(y) \vee t_{C_2}^n(y), \quad I_{C_1 \times C_2}^n(y) = I_{C_1}^n(y) \vee I_{C_2}^n(y), \quad f_{C_1 \times C_2}^n(y) = f_{C_1}^n(y) \wedge f_{C_2}^n(y),$$

for all $y \in Y_1 \times Y_2$.

1. $t_{D_1 \times D_2}^p((y_1, y_2)(z_1, z_2)) = t_{D_1}^p(y_1 z_1) \wedge t_{D_2}^p(y_2 z_2)$, $t_{D_1 \times D_2}^n((y_1, y_2)(z_1, z_2)) = t_{D_1}^n(y_1 z_1) \vee t_{D_2}^n(y_2 z_2)$, for all $y_1 z_1 \in E_1, y_2 z_2 \in E_2$,
2. $I_{D_1 \times D_2}^p((y_1, y_2)(z_1, z_2)) = I_{D_1}^p(y_1 z_1) \wedge I_{D_2}^p(y_2 z_2)$, $I_{D_1 \times D_2}^n((y_1, y_2)(z_1, z_2)) = I_{D_1}^n(y_1 z_1) \vee I_{D_2}^n(y_2 z_2)$, for all $y_1 z_1 \in E_1, y_2 z_2 \in E_2$,
3. $f_{D_1 \times D_2}^p((y_1, y_2)(z_1, z_2)) = f_{D_1}^p(y_1 z_1) \vee f_{D_2}^p(y_2 z_2)$, $f_{D_1 \times D_2}^n((y_1, y_2)(z_1, z_2)) = f_{D_1}^n(y_1 z_1) \wedge f_{D_2}^n(y_2 z_2)$, for all $y_1 z_1 \in E_1, y_2 z_2 \in E_2$.

Proposition 2.1. *Let G_1 and G_2 be any two bipolar neutrosophic graphs then $G_1 \cup G_2, G_1 \cap G_2, G_1 + G_2, G_1 \square G_2$ and $G_1 \times G_2$ are bipolar neutrosophic graphs.*

Definition 2.10. A bipolar neutrosophic graph $G = (C, D)$ is called *strong bipolar neutrosophic graph* if

$$\begin{aligned} t_D^p(yz) &= t_C^p(y) \wedge t_C^p(z), & I_D^p(yz) &= I_C^p(y) \wedge I_C^p(z), & f_D^p(yz) &= f_C^p(y) \vee f_C^p(z), \\ t_D^n(yz) &= t_C^n(y) \vee t_C^n(z), & I_D^n(yz) &= I_C^n(y) \vee I_C^n(z), & f_D^n(yz) &= f_C^n(y) \wedge f_C^n(z), \end{aligned}$$

for all $yz \in E$, E is the set of edges.

Definition 2.11. A bipolar neutrosophic graph $G = (C, D)$ is called *complete bipolar neutrosophic graph* if

$$\begin{aligned} t_D^p(yz) &= t_C^p(y) \wedge t_C^p(z), & I_D^p(yz) &= I_C^p(y) \wedge I_C^p(z), & f_D^p(yz) &= f_C^p(y) \vee f_C^p(z), \\ t_D^n(yz) &= t_C^n(y) \vee t_C^n(z), & I_D^n(yz) &= I_C^n(y) \vee I_C^n(z), & f_D^n(yz) &= f_C^n(y) \wedge f_C^n(z), \end{aligned}$$

for all $y, z \in Y$.

Definition 2.12. The *complement* of a bipolar neutrosophic graph $G = (C, D)$ is defined as a pair $G^c = (C^c, D^c)$ such that, $C^c(y) = C(y)$, for all $y \in Y$, and the membership values of the edges in G^c can be calculated as,

$$\begin{aligned} t_{D^c}^p(yz) &= t_C^p(y) \wedge t_C^p(z) - t_D^p(yz), & I_{D^c}^p(yz) &= I_C^p(y) \wedge I_C^p(z) - I_D^p(yz), \\ f_{D^c}^p(yz) &= f_C^p(y) \vee f_C^p(z) - f_D^p(yz), & t_{D^c}^n(yz) &= t_C^n(y) \vee t_C^n(z) - t_D^n(yz), \\ I_{D^c}^n(yz) &= I_C^n(y) \vee I_C^n(z) - I_D^n(yz), & f_{D^c}^n(yz) &= f_C^n(y) \wedge f_C^n(z) - f_D^n(yz), \end{aligned}$$

for all $yz \in \tilde{Y}^2$.

Remark 2.1. A bipolar neutrosophic graph G is called *self complementary* if $G = G^c$.

Theorem 2.1. *Let G be a self complementary bipolar neutrosophic graph then,*

$$\begin{aligned} \sum_{y \neq z} t_D^p(yz) &= \frac{1}{2} \sum_{y \neq z} t_C^p(y) \wedge t_C^p(z), & \sum_{y \neq z} I_D^p(yz) &= \frac{1}{2} \sum_{y \neq z} I_C^p(y) \wedge I_C^p(z), \\ \sum_{y \neq z} f_D^p(yz) &= \frac{1}{2} \sum_{y \neq z} f_C^p(y) \vee f_C^p(z), & \sum_{y \neq z} t_D^n(yz) &= \frac{1}{2} \sum_{y \neq z} t_C^n(y) \vee t_C^n(z), \\ \sum_{y \neq z} I_D^n(yz) &= \frac{1}{2} \sum_{y \neq z} I_C^n(y) \vee I_C^n(z), & \sum_{y \neq z} f_D^n(yz) &= \frac{1}{2} \sum_{y \neq z} f_C^n(y) \wedge f_C^n(z). \end{aligned}$$

Theorem 2.2. Let $G = (C, D)$ be a bipolar neutrosophic graph such that for all $y, z \in Y$,

$$\begin{aligned} t_{D^c}^p(yz) &= \frac{1}{2}(t_C^p(y) \wedge t_C^p(z)), & I_{D^c}^p(yz) &= \frac{1}{2}(I_C^p(y) \wedge I_C^p(z)), & f_{D^c}^p(yz) &= \frac{1}{2}(f_C^p(y) \vee f_C^p(z)), \\ t_{D^c}^n(yz) &= \frac{1}{2}(t_C^n(y) \vee t_C^n(z)), & I_{D^c}^n(yz) &= \frac{1}{2}(I_C^n(y) \vee I_C^n(z)), & f_{D^c}^n(yz) &= \frac{1}{2}(f_C^n(y) \wedge f_C^n(z)). \end{aligned}$$

Then G is self complementary bipolar neutrosophic graph.

Proof. Let $G^c = (C^c, D^c)$ be the complement of a bipolar neutrosophic graph $G = (C, D)$ then, by Definition. 2.12,

$$\begin{aligned} t_{D^c}^p(yz) &= t_C^p(y) \wedge t_C^p(z) - t_D^p(yz) \\ t_{D^c}^p(yz) &= t_C^p(y) \wedge t_C^p(z) - \frac{1}{2}(t_C^p(y) \wedge t_C^p(z)) \\ t_{D^c}^p(yz) &= \frac{1}{2}(t_C^p(y) \wedge t_C^p(z)) \\ t_{D^c}^p(yz) &= t_D^p(yz) \end{aligned}$$

Similarly, it can be proved that $t_{D^c}^n(yz) = t_D^n(yz)$, $I_{D^c}^p(yz) = I_D^p(yz)$, $I_{D^c}^n(yz) = I_D^n(yz)$, $f_{D^c}^p(yz) = f_D^p(yz)$ and $f_{D^c}^n(yz) = f_D^n(yz)$. Hence, G is self complementary. **q.e.d.**

Definition 2.13. The *degree* of a vertex y in a bipolar neutrosophic graph $G = (C, D)$ is denoted by $\deg(y)$ and defined by the 6-tuple as,

$$\begin{aligned} \deg(y) &= (\deg_t^p(y), \deg_I^p(y), \deg_f^p(y), \deg_t^n(y), \deg_I^n(y), \deg_f^n(y)), \\ &= \left(\sum_{yz \in E} t_D^p(yz), \sum_{yz \in E} I_D^p(yz), \sum_{yz \in E} f_D^p(yz), \sum_{yz \in E} t_D^n(yz), \sum_{yz \in E} I_D^n(yz), \sum_{yz \in E} f_D^n(yz) \right). \end{aligned}$$

The term degree is also referred as *neighborhood degree*.

Definition 2.14. The closed neighborhood degree of a vertex y in a bipolar neutrosophic graph is denoted by $\deg[y]$ and defined as, $\deg[y] = \deg(y) + C(y)$.

Definition 2.15. A bipolar neutrosophic graph G is known as a *regular* bipolar neutrosophic graph if all vertices of G have same degree. A bipolar neutrosophic graph G is known as a *totally regular* bipolar neutrosophic graph if all vertices of G have same closed neighborhood degree.

Theorem 2.3. A complete bipolar neutrosophic graph is regular.

Theorem 2.4. Let $G = (C, D)$ be a bipolar neutrosophic graph then $C = (t^p, I^p, f^p, t^n, I^n, f^n)$ is a constant function if and only if the following statements are equivalent:

- (1) G is a regular bipolar neutrosophic graph,
- (2) G is totally regular bipolar neutrosophic graph.

Proof. Assume that C is a constant function and for all $y \in Y$,

$$t_C^p(y) = k_t, I_C^p(y) = k_I, f_C^p(y) = k_f, t_C^n(y) = k'_t, I_C^n(y) = k'_I, f_C^n(y) = k'_f$$

where, $k_t, k_I, k_f, k'_t, k'_I, k'_f$ are constants.

(1) \Rightarrow (2) Suppose that G is a regular bipolar neutrosophic graph and $\deg(y) = (p_t, p_I, p_f, n_t, n_I, n_f)$ for all $y \in Y$.

Now consider,

$$\deg[y] = \deg(y) + C(y) = (p_t + k_t, p_I + k_I, p_f + k_f, n_t + k'_t, n_I + k'_I, n_f + k'_f), \quad \text{for all } y \in Y.$$

It is proved that G is totally regular bipolar neutrosophic graph.

(2) \Rightarrow (1) Suppose that G is totally regular bipolar neutrosophic graph and for all $y \in Y$

$$\deg[y] = (p'_t, p'_I, p'_f, n'_t, n'_I, n'_f).$$

$$\Rightarrow \deg(y) + C(y) = (p'_t, p'_I, p'_f, n'_t, n'_I, n'_f),$$

$$\Rightarrow \deg(y) = (p'_t - k_t, p'_I - k_I, p'_f - k_f, n'_t - k'_t, n'_I - k'_I, n'_f - k'_f),$$

for all $y \in Y$. Thus G is a regular bipolar neutrosophic graph.

Conversely, assume that the conditions are equivalent. Let $\deg(y) = (c_t, c_I, c_f, d_t, d_I, d_f)$ and $\deg[y] = (c'_t, c'_I, c'_f, d'_t, d'_I, d'_f)$.

Since by definition of closed neighborhood degree for all $y \in Y$,

$$\deg[y] = \deg(y) + C(y),$$

$$\Rightarrow C(y) = (c'_t - c_t, c'_I - c_I, c'_f - c_f, d'_t - d_t, d'_I - d_I, d'_f - d_f),$$

for all $y \in Y$. Hence $C = (c'_t - c_t, c'_I - c_I, c'_f - c_f, d'_t - d_t, d'_I - d_I, d'_f - d_f)$, a constant function which completes the proof. **q.e.d.**

Definition 2.16. A bipolar neutrosophic graph G is said to be *irregular* if at least two vertices have distinct degrees. If all vertices do not have same closed neighborhood degrees then G is known as *totally irregular* bipolar neutrosophic graph.

Theorem 2.5. Let $G = (C, D)$ be a bipolar neutrosophic graph and $C = (t_C^p, I_C^p, f_C^p, t_C^n, I_C^n, f_C^n)$ be a constant function then G is an irregular bipolar neutrosophic graph if and only if G is a totally irregular bipolar neutrosophic graph.

Proof. Assume that G is an irregular bipolar neutrosophic graph then at least two vertices of G have distinct degrees. Let y and z be two vertices such that $\deg(y) = (r_1, r_2, r_3, s_1, s_2, s_3)$ and $\deg(z) = (r'_1, r'_2, r'_3, s'_1, s'_2, s'_3)$ where, $r_i \neq r'_i$, for some $i = 1, 2, 3$.

Since, C is a constant function let $C = (k_1, k_2, k_3, l_1, l_2, l_3)$. Therefore,

$$\deg[y] = \deg(y) + (k_1, k_2, k_3, l_1, l_2, l_3)$$

$$\deg[y] = (r_1 + k_1, r_2 + k_2, r_3 + k_3, s_1 + l_1, s_2 + l_2, s_3 + l_3)$$

$$\text{and } \deg[z] = (r'_1 + k_1, r'_2 + k_2, r'_3 + k_3, s'_1 + l_1, s'_2 + l_2, s'_3 + l_3).$$

Clearly $r_i + k_i \neq r'_i + k_i$, for some $i = 1, 2, 3$ therefore y and z have distinct closed neighborhood degrees. Hence G is a totally irregular bipolar neutrosophic graph.

The converse part is similar.

q.e.d.

Definition 2.17. If $G = (C, D)$ is a bipolar neutrosophic graph and y, z are two vertices in G then we say that y *dominates* z if

$$\begin{aligned} t_D^p(yz) &= t_C^p(y) \wedge t_C^p(z), & I_D^p(yz) &= I_C^p(y) \wedge I_C^p(z), & f_D^p(yz) &= f_C^p(y) \vee f_C^p(z), \\ t_D^n(yz) &= t_C^n(y) \vee t_C^n(z), & I_D^n(yz) &= I_C^n(y) \vee I_C^n(z), & f_D^n(yz) &= f_C^n(y) \wedge f_C^n(z). \end{aligned}$$

A subset $D' \subseteq Y$ is a *dominating set* if for each $z \in Y \setminus D'$ there exists $y \in D'$ such that y dominates z . A dominating set D' is minimal if for every $y \in D'$, $D' \setminus \{y\}$ is not a dominating set. The *domination number* of G is the minimum cardinality among all minimal dominating sets of G , denoted by $\lambda(G)$.

Example 2.2. Consider a bipolar neutrosophic graph as shown in Fig. 2. The set $\{x, w\}$ is a minimal dominating set and $\lambda(G) = 2$.




Figure 2: Bipolar neutrosophic graph G .

Theorem 2.6. Let G_1 and G_2 be two bipolar neutrosophic graphs with D'_1 and D'_2 as dominating sets then following conditions hold,

- (1) If $Y_1 \cap Y_2 = \emptyset$ then, $\lambda(G_1 \cup G_2) = \lambda(G_1) + \lambda(G_2)$.
- (2) If for every $y \in Y_1 \cap Y_2 \neq \emptyset$, $C_1(y) = C_2(y)$ and for $yz \in E_1 \cap E_2$, $D_1(yz) = D_2(yz)$ then, $\lambda(G_1 \cup G_2) = \lambda(G_1) + \lambda(G_2) - |D'_1 \cap D'_2|$.

Proof. (1). The proof is obvious.

(2). Since D'_1 and D'_2 are dominating sets of G_1 and G_2 , $D'_1 \cup D'_2$ is a dominating set of $G_1 \cup G_2$. Therefore, $\lambda(G_1 \cup G_2) \leq |D'_1 \cup D'_2|$. It only remains to show that $D'_1 \cup D'_2$ is a minimal dominating set. On contrary, assume that $D' = D'_1 \cup D'_2 \setminus \{y\}$ is a minimal dominating set of $G_1 \cup G_2$. There are two cases,

Case 1. If $y \in D'_1$ and $y \notin D'_2$, then $D'_1 \setminus \{y\}$ is not a dominating set of G_1 which implies that $D'_1 \cup D'_2 \setminus \{y\} = D'$ is not a dominating set of $G_1 \cup G_2$. A contradiction, hence $D'_1 \cup D'_2$ is a minimal dominating set and

$$\begin{aligned} \lambda(G_1 \cup G_2) &= |D'_1 \cup D'_2|, \\ \Rightarrow \lambda(G_1 \cup G_2) &= \lambda(G_1) + \lambda(G_2) - |D'_1 \cap D'_2|. \end{aligned}$$

Case 2. If $y \in D'_2$ and $y \notin D'_1$, same contradiction can be obtained. **q.e.d.**

Theorem 2.7. *If G_1 and G_2 are two bipolar neutrosophic graphs then the following conditions are satisfied,*

- (1) *If $Y_1 \cap Y_2 = \emptyset$ then, $\lambda(G_1 + G_2) = 2$.*
- (2) *If for every $y \in Y_1 \cap Y_2 \neq \emptyset$, $C_1(y) = C_2(y)$ and for $yz \in E_1 \cap E_2$, $D_1(yz) = D_2(yz)$ then, $\lambda(G_1 + G_2) = \min\{\lambda(G_1), \lambda(G_2), 2\}$.*

Proof. (1). Let $y_1 \in Y_1$ and $y_2 \in Y_2$, since $G_1 + G_2$ is a bipolar neutrosophic graph, we have

$$\begin{aligned} t_{D_1+D_2}^p(y_1y_2) &= t_{C_1+C_2}^p(y_1) \wedge t_{C_1+C_2}^p(y_2), & t_{D_1+D_2}^n(y_1y_2) &= t_{C_1+C_2}^n(y_1) \vee t_{C_1+C_2}^n(y_2) \\ I_{D_1+D_2}^p(y_1y_2) &= I_{C_1+C_2}^p(y_1) \wedge I_{C_1+C_2}^p(y_2), & I_{D_1+D_2}^n(y_1y_2) &= I_{C_1+C_2}^n(y_1) \vee I_{C_1+C_2}^n(y_2) \\ f_{D_1+D_2}^p(y_1y_2) &= f_{C_1+C_2}^p(y_1) \vee f_{C_1+C_2}^p(y_2), & f_{D_1+D_2}^n(y_1y_2) &= f_{C_1+C_2}^n(y_1) \wedge f_{C_1+C_2}^n(y_2). \end{aligned}$$

Hence any vertex of G_1 dominates all vertices of G_2 and similarly any vertex of G_2 dominates all vertices of G_1 . So, $\{y_1, y_2\}$ is a dominating set of $G_1 + G_2$.

(2). If D is a minimal dominating set of $G_1 + G_2$ then, D is one of the following forms,

1. $D = D_1$ where, $\lambda(G_1) = |D_1|$,
2. $D = D_2$ where, $\lambda(G_2) = |D_2|$,
3. $D = \{y_1, y_2\}$ where, $y_1 \in Y_1$ and $y_2 \in Y_2$. $\{y_1\}$ and $\{y_2\}$ are not dominating sets of G_1 or G_2 , respectively.

Hence, $\lambda(G_1 + G_2) = \min\{\lambda(G_1), \lambda(G_2), 2\}$.

q.e.d.

Theorem 2.8. *Let $G_1 = (C_1, D_1)$ and $G_2 = (C_2, D_2)$ be two bipolar neutrosophic graphs. If for $y_1 \in Y_1$, $C_1(y_1) \neq (0, 0, 1, 0, 0, -1)$, and y_2 dominates z_2 in G_2 then, (y_1, y_2) dominates (y_1, z_2) in $G_1 \square G_2$.*

Proof. Since y_2 dominates z_2 therefore,

$$\begin{aligned} t_{D_2}^p(y_2z_2) &= t_{C_2}^p(y_2) \wedge t_{C_2}^p(z_2), & I_{D_2}^p(y_2z_2) &= I_{C_2}^p(y_2) \wedge I_{C_2}^p(z_2), \\ f_{D_2}^p(y_2z_2) &= f_{C_2}^p(y_2) \vee f_{C_2}^p(z_2), & t_{D_2}^n(y_2z_2) &= t_{C_2}^n(y_2) \vee t_{C_2}^n(z_2), \\ I_{D_2}^n(y_2z_2) &= I_{C_2}^n(y_2) \vee I_{C_2}^n(z_2), & f_{D_2}^n(y_2z_2) &= f_{C_2}^n(y_2) \wedge f_{C_2}^n(z_2). \end{aligned}$$

For $y_1 \in Y_1$, take $(y_1, z_2) \in Y_1 \times Y_2$. By Definition 2.8,

$$\begin{aligned} t_{D_1 \square D_2}^p((y_1, y_2)(y_1, z_2)) &= t_{C_1}^p(y_1) \wedge t_{D_2}^p(y_2z_2), \\ &= t_{C_1}^p(y_1) \wedge \{t_{C_2}^p(y_2) \wedge t_{C_2}^p(z_2)\}, \\ &= \{t_{C_1}^p(y_1) \wedge t_{C_2}^p(y_2)\} \wedge \{t_{C_1}^p(y_1) \wedge t_{C_2}^p(z_2)\}, \\ &= t_{C_1 \square C_2}^p(y_1, y_2) \wedge t_{C_1 \square C_2}^p(y_1, z_2). \end{aligned}$$

$$\begin{aligned}
t_{D_1 \square D_2}^n((y_1, y_2)(y_1, z_2)) &= t_{C_1}^n(y_1) \vee t_{D_2}^n(y_2 z_2), \\
&= t_{C_1}^n(y_1) \vee \{t_{C_2}^n(y_2) \vee t_{C_2}^n(z_2)\}, \\
&= \{t_{C_1}^n(y_1) \vee t_{C_2}^n(y_2)\} \vee \{t_{C_1}^n(y_1) \vee t_{C_2}^n(z_2)\}, \\
&= t_{C_1 \square C_2}^n(y_1, y_2) \vee t_{C_1 \square C_2}^n(y_1, z_2).
\end{aligned}$$

Similarly, it can be proved that

$$\begin{aligned}
I_{D_1 \square D_2}^p((y_1, y_2)(y_1, z_2)) &= I_{C_1 \square C_2}^p(y_1, y_2) \wedge I_{C_1 \square C_2}^p(y_1, z_2), \\
I_{D_1 \square D_2}^n((y_1, y_2)(y_1, z_2)) &= I_{C_1 \square C_2}^n(y_1, y_2) \vee I_{C_1 \square C_2}^n(y_1, z_2), \\
f_{D_1 \square D_2}^p((y_1, y_2)(y_1, z_2)) &= f_{C_1 \square C_2}^p(y_1, y_2) \vee f_{C_1 \square C_2}^p(y_1, z_2), \\
f_{D_1 \square D_2}^n((y_1, y_2)(y_1, z_2)) &= f_{C_1 \square C_2}^n(y_1, y_2) \wedge f_{C_1 \square C_2}^n(y_1, z_2).
\end{aligned}$$

Hence (y_1, y_2) dominates (y_1, z_2) and the proof is complete. **q.e.d.**

Proposition 2.2. *If G_1 and G_2 are bipolar neutrosophic graphs and for $z_2 \in Y_2$, $C_2(z_2) \neq (0, 0, 1, 0, 0, -1)$, y_1 dominates z_1 in G_1 then (y_1, z_2) dominates (z_1, z_2) in $G_1 \square G_2$.*

Theorem 2.9. *If D'_1 and D'_2 are minimal dominating sets of $G_1 = (C_1, D_1)$ and $G_2 = (C_2, D_2)$, respectively. Then $D'_1 \times Y_2$ and $Y_1 \times D'_2$ are dominating sets of $G_1 \square G_2$ and*

$$(2.1) \quad \lambda(G_1 \square G_2) \leq |D'_1 \times Y_2| \wedge |Y_1 \times D'_2|.$$

Proof. To prove Inequality. (2.1), we need to show that $D'_1 \times Y_2$ and $Y_1 \times D'_2$ are dominating sets of $G_1 \square G_2$. Let $(z_1, z_2) \notin D'_1 \times Y_2$ then, $z_1 \notin D'_1$. Since D'_1 is a dominating set of G_1 , there exists $y_1 \in D'_1$ that dominates z_1 . By Theorem 2.2, (y_1, z_2) dominates (z_1, z_2) in $G_1 \square G_2$. Since (z_1, z_2) was taken to be arbitrary therefore, $D'_1 \times Y_2$ is a dominating set of $G_1 \square G_2$. Similarly, $Y_1 \times D'_2$ is a dominating set if $G_1 \square G_2$. Hence the proof. **q.e.d.**

Theorem 2.10. *Let D'_1 and D'_2 be the dominating sets of $G_1 = (C_1, D_1)$ and $G_2 = (C_2, D_2)$, respectively then, $D'_1 \times D'_2$ is a dominating set of the direct product $G_1 \times G_2$ and*

$$(2.2) \quad \lambda(G_1 \times G_2) = |D'_x| = |D'_1 \times D'_2 \cup \{(t_1, t_2) : t_1 y_1 \in E_1, y_1 \in D'_1, t_2 \in D'_2\}|.$$

Proof. Let $(z_1, z_2) \in Y_1 \times Y_2 \setminus D'_x$ then there are two cases.

Case 1: If $z_1 \in Y_1 \setminus D'_1$ and $z_2 \in Y_2 \setminus D'_2$. Since, D'_1 and D'_2 are dominating sets there exist $y_1 \in D'_1$ and $y_2 \in D'_2$ such that y_1 dominates z_1 and y_2 dominates z_2 . Consider,

$$\begin{aligned}
t_{D_1 \times D_2}^p((y_1, y_2)(z_1, z_2)) &= t_{D_1}^p(y_1 z_1) \wedge t_{D_2}^p(y_2 z_2), \\
&= \{t_{C_1}^p(y_1) \wedge t_{C_1}^p(z_1)\} \wedge \{t_{C_2}^p(y_2) \wedge t_{C_2}^p(z_2)\}, \\
&= \{t_{C_1}^p(y_1) \wedge t_{C_2}^p(y_2)\} \wedge \{t_{C_1}^p(z_1) \wedge t_{C_2}^p(z_2)\}, \\
&= t_{C_1 \times C_2}^p(y_1, y_2) \wedge t_{C_1 \times C_2}^p(z_1, z_2).
\end{aligned}$$

Similarly, it can be proved for other truth, indeterminacy and falsity membership degrees. Hence (y_1, y_2) dominates (z_1, z_2) .

Case 2: If $z_1 \in D'_1$ and $z_2 \in Y_2 \setminus D'_2$ then, there exists $(t_1, t_2) \in \{(t_1, t_2) : t_1 z_1 \in E_1, z_1 \in D'_1, t_2 \in D'_2\}$ such that z_1 dominates t_1 and t_2 dominates z_2 . Consider,

$$\begin{aligned} t_{D_1 \times D_2}^p((t_1, t_2)(z_1, z_2)) &= t_{D_1}^p(t_1 z_1) \wedge t_{D_2}^p(t_2 z_2), \\ &= \{t_{C_1}^p(t_1) \wedge t_{C_1}^p(z_1)\} \wedge \{t_{C_2}^p(t_2) \wedge t_{C_2}^p(z_2)\}, \\ &= \{t_{C_1}^p(t_1) \wedge t_{C_2}^p(t_2)\} \wedge \{t_{C_1}^p(z_1) \wedge t_{C_2}^p(z_2)\}, \\ &= t_{C_1 \times C_2}^p(t_1, t_2) \wedge t_{C_1 \times C_2}^p(z_1, z_2). \end{aligned}$$

On the same lines, the result can be obtained for other truth, indeterminacy and falsity membership degrees. Hence (z_1, z_2) is dominated by (t_1, t_2) .

Since (y_1, y_2) was taken to be arbitrary therefore, every element of $Y_1 \times Y_2 \setminus D'_\times$ is dominated by some element of D'_\times .

Clearly, no vertex in $\{(t_1, t_2) : t_1 y_1 \in E_1, y_1 \in D'_1, t_2 \in D'_2\}$ is dominated by any other vertex in D'_\times . Therefore, it only remains to show that $D'_1 \times D'_2$ is minimal. On contrary, assume that D' is a minimal such that $|D'| < |D'_1 \times D'_2|$. Let $(t_1, t_2) \in D'_1 \times D'_2$ such that $(t_1, t_2) \notin D'$ i.e., $t_1 \in D'_1$ and $t_2 \in D'_2$ then there exist $t'_1 \in Y_1 \setminus D'_1$ and $t'_2 \in Y_2 \setminus D'_2$ which are only dominated by t_1 and t_2 , respectively. Hence no element other than (t_1, t_2) dominates (t'_1, t'_2) so $(t_1, t_2) \in D'$. A contradiction, thus $D'_1 \times D'_2$ is minimal. **q.e.d.**

Corollary 2.1. *If G_1 and G_2 are two bipolar neutrosophic graphs, y_1 dominates z_1 in G_1 and y_2 dominates z_2 in G_2 then (y_1, z_1) dominates (y_2, z_2) in $G_1 \times G_2$.*

Definition 2.18. In a bipolar neutrosophic graph two vertices y and z are *independent* if

$$(2.3) \quad \begin{aligned} t_D^p(yz) &< t_C^p(y) \wedge t_C^p(z), & I_D^p(yz) &< I_C^p(y) \wedge I_C^p(z), & f_D^p(yz) &< f_C^p(y) \vee f_C^p(z), \\ t_D^n(yz) &> t_C^n(y) \vee t_C^n(z), & I_D^n(yz) &> I_C^n(y) \vee I_C^n(z), & f_D^n(yz) &> f_C^n(y) \wedge f_C^n(z). \end{aligned}$$

An *independent set* N of a bipolar neutrosophic graph is a subset N of Y such that for all $y, z \in N$, Equations (2.3) are satisfied. An independent set is *maximal* if for every $t \in Y \setminus N$, $N \cup \{t\}$ is not an independent set. An *independent number* is the maximal cardinality among all maximal independent sets of a bipolar neutrosophic graph. It is denoted by $\alpha(G)$.

Theorem 2.11. *If G_1 and G_2 are bipolar neutrosophic graphs on Y_1 and Y_2 , respectively such that $Y_1 \cap Y_2 = \emptyset$ then $\alpha(G_1 \cup G_2) = \alpha(G_1) + \alpha(G_2)$.*

Proof. Let N_1 and N_2 be maximal independent sets of G_1 and G_2 . Since $N_1 \cap N_2 = \emptyset$ therefore, $N_1 \cup N_2$ is a maximal independent set of $G_1 \cup G_2$. Hence $\alpha(G_1 \cup G_2) = \alpha(G_1) + \alpha(G_2)$. **q.e.d.**

Theorem 2.12. *Let G_1 and G_2 be two bipolar neutrosophic graphs then $\alpha(G_1 + G_2) = \alpha(G_1) \vee \alpha(G_2)$.*

Proof. Let N_1 and N_2 be maximal independent sets. Since every vertex of G_1 dominates every vertex of G_2 in $G_1 + G_2$. Hence, maximal independent set of $G_1 + G_2$ is either N_1 or N_2 . Thus, $\alpha(G_1 + G_2) = \alpha(G_1) \vee \alpha(G_2)$. **q.e.d.**

Theorem 2.13. *If N_1 and N_2 are maximal independent sets of G_1 and G_2 , respectively and $Y_1 \cap Y_2 = \emptyset$. Then $\alpha(G_1 \square G_2) = |N_1 \times N_2| + |N|$ where, $N = \{(y_i, z_i) : y_i \in Y_1 \setminus N_1, z_i \in Y_2 \setminus N_2, y_i y_{i+1} \in E_1, z_i z_{i+1} \in E_2, i = 1, 2, 3, \dots\}$.*

Proof. N_1 and N_2 are maximal independent sets of G_1 and G_2 , respectively. Clearly, $N_1 \times N_2$ is an independent set of $G_1 \square G_2$ as no vertex of $N_1 \times N_2$ dominates any other vertex of $N_1 \times N_2$.

Consider the set of vertices $N = \{(y_i, z_i) : y_i \in Y_1 \setminus N_1, z_i \in Y_2 \setminus N_2, y_i y_{i+1} \in E_1, z_i z_{i+1} \in E_2\}$. It can be seen that no vertex $(y_i, z_i) \in N$ for each $i = 1, 2, 3, \dots$ dominates $(y_{i+1}, z_{i+1}) \in N$ for each $i = 1, 2, 3, \dots$. Hence $N' = (N_1 \times N_2) \cup N$ is an independent set of $G_1 \square G_2$.

Assume that $S = N' \cup \{(y_i, z_j)\}$, for some $i \neq j$, $y_i \in Y_1 \setminus N_1$ and $z_j \in Y_2 \setminus N_2$, is a maximal independent set. Without loss of generality, assume that $j = i + 1$ then, (y_i, z_j) is dominated by (y_i, z_i) . A contradiction, hence N' is a maximal independent set and $\alpha(G_1 \square G_2) = |N'| = |N_1 \times N_2| + |N|$ **q.e.d.**

Theorem 2.14. *If D'_\times is a minimal dominating sets of $G_1 \times G_2$ then, $Y_1 \times Y_2 \setminus D'_\times$ is a maximal independent set of $G_1 \times G_2$ and $\alpha(G_1 \times G_2) = n_1 n_2 - \lambda(G_1 \times G_2)$ where, n_1 and n_2 are the number of vertices in G_1 and G_2 .*

The proof is obvious.

Theorem 2.15. *An independent set of a bipolar neutrosophic graph $G = (C, D)$ is maximal if and only if it is independent and dominating.*

Proof. If N is a maximal independent set of G , then for every $y \in Y \setminus N$, $N \cup \{y\}$ is not an independent set. For every vertex $y \in Y \setminus N$, there exists some $z \in N$ such that

$$\begin{aligned} t_D^p(yz) &= t_C^p(y) \wedge t_C^p(z), & I_D^p(yz) &= I_C^p(y) \wedge I_C^p(z), & f_D^p(yz) &= f_C^p(y) \vee f_C^p(z), \\ t_D^n(yz) &= t_C^n(y) \vee t_C^n(z), & I_D^n(yz) &= I_C^n(y) \vee I_C^n(z), & f_D^n(yz) &= f_C^n(y) \wedge f_C^n(z). \end{aligned}$$

Thus y dominates z and hence N is both independent and dominating set. Conversely, assume that D is both independent and dominating set but not maximal independent set. So there exists a vertex $y \in Y \setminus N$ such that $N \cup \{y\}$ is an independent set i.e., no vertex in N dominates y , a contradiction to the fact that N is a dominating set. Hence N is maximal. **q.e.d.**

Theorem 2.16. *Any maximal independent set of a bipolar neutrosophic graph is a minimal dominating set.*

Proof. If N is a maximal independent set of a bipolar neutrosophic graph then by Theorem 2.15, N is a dominating set. Assume that N is not a minimal dominating set then, there always exist at least one $z \in N$ for which $N \setminus \{z\}$ is a dominating set. On the other hand if $N \setminus \{z\}$ dominates $Y \setminus \{N \setminus \{z\}\}$, at least one vertex in $N \setminus \{z\}$ dominates z . A contradiction to the fact that N is an independent set of bipolar neutrosophic graph G . Hence N is a minimal dominating set. **q.e.d.**

3 Multi-criteria decision making methods

Multiple criteria decision making refers to making decisions in the presence of multiple, usually conflicting criteria. Multi-criteria decision making problems are common in everyday life. In this section, we present multi-criteria decision making methods for the identification of risk in decision support systems. The method is explained by an example for prevention of accidental hazards in chemical industry. The application of domination in bipolar neutrosophic graphs is given for the construction of transmission stations.

(1) An outranking approach for safety analysis using bipolar neutrosophic information

The proposed methodology can be implemented in various fields in different ways e.g., multi-criteria decision making problems with bipolar neutrosophic information. However, our main focus is the identification of risk assessments in industry which is described in the following steps.

The bipolar neutrosophic information consists of a group of risks\alternatives $R = \{r_1, r_2, \dots, r_n\}$ evaluated on the basis of criteria $C = \{c_1, c_2, \dots, c_m\}$. Here r_i , $i = 1, 2, \dots, n$ is the possibility for the criteria c_k , $k = 1, 2, \dots, m$ and r_{ik} are in the form of bipolar neutrosophic values. This method is suitable if we have a small set of data and experts are able to evaluate the data in the form of bipolar neutrosophic information. Take the values of r_{ik} as $r_{ik} = (t_{ik}^p, I_{ik}^p, f_{ik}^p, t_{ik}^n, I_{ik}^n, f_{ik}^n)$.

Step 1. Construct the table of the given data.

Step 2. Determine the average values using the following bipolar neutrosophic average operator,

$$(3.1) \quad A_i = \frac{1}{n} \left(\sum_{j=1}^m t_{ij}^p - \prod_{j=1}^m t_{ij}^p, \prod_{j=1}^m I_{ij}^p, \prod_{j=1}^m f_{ij}^p, \prod_{j=1}^m t_{ij}^n, \sum_{j=1}^m I_{ij}^n - \prod_{j=1}^m I_{ij}^n, \sum_{j=1}^m f_{ij}^n - \prod_{j=1}^m f_{ij}^n \right),$$

for each $i = 1, 2, \dots, n$.

Step 3. Construct the weighted average matrix.

Choose the weight vector $\mathbf{w} = (w_1, w_2, \dots, w_n)$. According to the weights for each alternative, the weighted average table can be calculated by multiplying each average value with the corresponding weight as:

$$\beta_i = A_i w_i, \quad i = 1, 2, \dots, n.$$

Step 4. Calculate the normalized value for each alternative\risk β_i using the formula,

$$(3.2) \quad \alpha_i = \sqrt{(t_i^p)^2 + (I_i^p)^2 + (f_i^p)^2 + (-1 + t_i^n)^2 + (-1 + I_i^n)^2 + (-1 + f_i^n)^2},$$

for each $i = 1, 2, \dots, n$. The resulting table indicate the preference ordering of the alternatives\risks. The alternative\risk with maximum α_i value is most dangerous or more preferable.

Example 3.1. Chemical industry is a very important part of human society. These industries contain large amount of organic and inorganic chemicals and materials. Many chemical products have a high risk of fire due to flammable materials, large explosions and oxygen deficiency etc. These accidents can cause the death of employs, damages to building, destruction of machines and transports, economical losses etc. Therefore, it is very important to prevent these accidental losses by identifying the major risks of fire, explosions and oxygen deficiency. A manager of a chemical industry Y wants to prevent such types of accidents that caused the major loss to company in the past. He collected data from witness reports, investigation teams and near by chemical industries and found that the major causes could be the chemical reactions, oxidizing materials, formation of toxic substances, electric hazards, oil spill, hydrocarbon gas leakage and energy systems. The witness reports, investigation teams and industries have different opinions. There is a bipolarity in people’s thinking and judgement. The data can be considered as bipolar neutrosophic information. The bipolar neutrosophic information about company Y old accidents are given in Table. 3 and Table. 4.

Table 3: Bipolar neutrosophic Data

	Fire	Oxygen Deficiency
Chemical Exposures	(0.5,0.7,0.2,-0.6,-0.3,-0.7)	(0.1,0.5,0.7,-0.5,-0.2,-0.8)
Oxidizing materials	(0.9,0.7,0.2,-0.8,-0.6,-0.1)	(0.3,0.5,0.2,-0.5,-0.5,-0.2)
Toxic vapour cloud	(0.7,0.3,0.1,-0.4,-0.1,-0.3)	(0.6,0.3,0.2,-0.5,-0.3,-0.3)
Electric Hazard	(0.3,0.4,0.2,-0.6,-0.3,-0.7)	(0.9,0.4,0.6,-0.1,-0.7,-0.5)
Oil Spill	(0.7,0.5,0.3,-0.4,-0.2,-0.2)	(0.2,0.2,0.2,-0.7,-0.4,-0.4)
Hydrocarbon gas leakage	(0.5,0.3,0.2,-0.5,-0.2,-0.2)	(0.3,0.2,0.3,-0.7,-0.4,-0.3)
Ammonium Nitrate	(0.3,0.2,0.3,-0.5,-0.6,-0.5)	(0.9,0.2,0.1,0.0,-0.6,-0.5)

Table 4: Bipolar neutrosophic Data

	Large Explosion
Chemical Exposures	(0.6,0.2,0.3,-0.4,0.0,-0.1)
Oxidizing materials	(0.9,0.5,0.5,-0.6,-0.5,-0.2)
Toxic vapour cloud	(0.5,0.1,0.2,-0.6,-0.2,-0.2)
Electric Hazard	(0.7,0.6,0.8,-0.7,-0.5,-0.1)
Oil Spill	(0.9,0.2,0.7,-0.1,-0.6,-0.8)

Continued on next page

Table 4 – *Continued from previous page*

Hydrocarbon gas leakage	(0.8,0.2,0.1,-0.1,-0.9,-0.2)
Ammonium Nitrate	(0.6,0.2,0.1,-0.2,-0.3,-0.5)

By applying the bipolar neutrosophic average operator 3.1 on Table. 3 and Table. 4, the average values are given in Table. 5.

Table 5: Bipolar neutrosophic average values

	Average Value
Chemical Exposures	(0.39,0.023,0.014,-0.04,-0.167,-0.515)
Oxidizing materials	(0.619,0.032,0.001,-0.08,-0.483,-0.165)
Toxic vapour cloud	(0.53,0.003,0.001,-0.04,-0.198,-0.261)
Electric Hazard	(0.570,0.032,0.032,-0.014,-0.465,-0.422)
Oil Spill	(0.558,0.007,0.014,-0.009,-0.384,-0.445)
Hydrocarbon gas leakage	(0.493,0.004,0.002,-0.011,-0.543,-0.229)
Ammonium Nitrate	(0.546,0.003,0.001,0.0,-0.464,-0.417)

With regard to the weight vector (0.35, 0.80, 0.30, 0.275, 0.65, 0.75, 0.50) associated to each cause of accident, the weighted average values are obtained by multiplying each average value with corresponding weight and are given in Table. 6.

Table 6: Bipolar neutrosophic weighted average table

	Weighted Value
Chemical Exposures	(0.1365,0.0081,0.0049,-0.0140,-0.0585,-0.1803)
Oxidizing materials	(0.4952,0.0256,0.0008,-0.0640,-0.3864,-0.1320)
Toxic vapour cloud	(0.1590,0.0009,0.0003,-0.012,-0.0594,-0.0783)
Electric Hazard	(0.2850,0.0160,0.0160,-0.0070,-0.2325,-0.2110)
Oil Spill	(0.1535,0.0019,0.0039,-0.0025,-0.1056,-0.1224)
Hydrocarbon gas leakage	(0.3205,0.0026,0.0013,-0.0072,-0.3530,-0.1489)
Ammonium Nitrate	(0.4095,0.0023,0.0008,0.0,-0.3480,-0.2110)

Using Formula. 3.2, the resulting normalized values are shown in Table. 8.

Table 7: Normalized values

	Normalized value
Chemical Exposures	1.5966
Oxidizing materials	1.5006
Toxic vapour cloud	1.6540
Electric Hazard	1.6090
Oil Spill	1.4938
Hydrocarbon gas leakage	1.6036

Continued on next page

Table 7 – <i>Continued from previous page</i>	
Ammonium Nitrate	1.5089

The accident possibilities can be placed in the following order: toxic vapour cloud \succ electric hazard \succ hydrocarbon gas leakage \succ chemical exposures \succ ammonium nitrate \succ oxidizing materials \succ Oil spill where, the symbol \succ represents partial ordering of objects. It can be easily seen that the formation of toxic vapour clouds, electrical and energy systems and hydrocarbon gas leakage are the major dangers to the chemical industry. There is a very little danger due to oil spill. Chemical exposures, oxidizing materials and ammonium nitrate has an average accidental danger. Therefore, industry needs special precautions to prevent the major hazards that could happen due the formation of toxic vapour clouds.

(2) Domination in bipolar neutrosophic graphs

Domination has a wide variety of applications in communication networks, coding theory, fixing surveillance cameras, detecting biological proteins and social networks etc. Consider the example of a TV channel that wants to set up transmission stations in a number of cities such that every city in the country get access to the channel signals from at least one of the stations. To reduce the cost for building large stations it is required to set up minimum number of stations. This problem can be represented by a bipolar neutrosophic graph in which vertices represent the cities and there is an edge between two cities if they can communicate directly with each other. Consider a network of ten cities $\{C_1, C_2, \dots, C_{10}\}$. In the bipolar neutrosophic graph, the positive degree of each vertex represents the level of truth, indeterminacy and falsity of strong signals it can transmit to other cities and the negative degree of each vertex represents the level of truth, indeterminacy and falsity of weaker signals it can transmit to other cities. The bipolar neutrosophic value of each edge represents the degree of truth, indeterminacy and falsity of strong and weak communication between the cities. The graph is shown in Fig. 3. $D = \{C_8, C_{10}\}$ is the minimum dominating set. It is concluded that building only two large transmitting stations in C_8 and C_{10} , a high economical benefit can be achieved. The method of calculating the minimum number of stations is described in the following Algorithm. 1.

Table 8: Algorithm for the selection of minimum locations

Algorithm 1
1. Begin
2. Enter the membership values $B(x_i)$ of n number of possible locations $A = \{C_1, C_2, \dots, C_n\}$.
3. Input the adjacency matrix $[C_{ij}]_{n \times n}$ of transmission stations.
4. $k = 0$
5. $D = \emptyset$
6. do i from 1 $\rightarrow n$
7. do j from $i + 1 \rightarrow n$
8. if $C_{ij} = A(C_i) \cap A(C_j)$ then

Continued on next page

Table 8 – *Continued from previous page*

9.	$C_i \in D, k = k + 1, x_k = C_i$
10.	end if
11.	end do
12.	end do
13.	Arrange $X \setminus D = \{x_{k+1}, x_{k+2}, \dots, x_n\} = J$
14.	do i from 1 $\rightarrow k$
15.	$D' = D \setminus \{x_i\}$
16.	if D' is a dominating set then
17.	$D = D'$
18.	$J = J \cup \{x_i\}$
19.	end if
20.	end do
21.	if $D \cup J = Y$ then
22.	print: D is a minimal dominating set.
23.	else
24.	print: There is no dominating set.
25.	end if

4 Comments on Broumi et al.'s Bipolar Neutrosophic Graphs

Broumi et al. [11] defined bipolar single-valued neutrosophic graphs in the following way:

Definition 4.1. [11] Let $A = (T_A^P, I_A^P, F_A^P, T_A^N, I_A^N, F_A^N)$ and $B = (T_B^P, I_B^P, F_B^P, T_B^N, I_B^N, F_B^N)$ be bipolar single valued neutrosophic graph on set Y . If $B = (T_B^P, I_B^P, F_B^P, T_B^N, I_B^N, F_B^N)$ is a bipolar single valued neutrosophic relation on $A = (T_A^P, I_A^P, F_A^P, T_A^N, I_A^N, F_A^N)$ then

$$T_B^P(xy) \leq T_A^P(x) \wedge T_A^P(y), \quad I_B^P(xy) \geq I_A^P(x) \vee I_A^P(y), \quad F_B^P(xy) \geq F_A^P(x) \vee F_A^P(y),$$

$$T_B^N(xy) \geq T_A^N(x) \vee T_A^N(y), \quad I_B^N(xy) \leq I_A^N(x) \wedge I_A^N(y), \quad F_B^N(xy) \leq F_A^N(x) \wedge F_A^N(y),$$

for all $x, y \in Y$.

Broumi et al. [11] defined complement of a bipolar neutrosophic graph as follows:

Definition 4.2. [11] The complement of a bipolar neutrosophic graph $G = (A, B)$ is a bipolar single valued neutrosophic graph $\overline{G} = (\overline{A}, \overline{B})$ where, $\overline{A} = A = (T_A^P, I_A^P, F_A^P, T_A^N, I_A^N, F_A^N)$, $\overline{B} = (\overline{T}_B^P, \overline{I}_B^P, \overline{F}_B^P, \overline{T}_B^N, \overline{I}_B^N, \overline{F}_B^N)$ and is defined by

$$\overline{T}_B^P(xy) = T_A^P(x) \wedge T_A^P(y) - T_B^P(xy), \quad \overline{I}_B^P(xy) = I_A^P(x) \vee I_A^P(y) - I_B^P(xy),$$

$$\overline{F}_B^P(xy) = F_A^P(x) \vee F_A^P(y) - F_B^P(xy), \quad \overline{T}_B^N(xy) = T_A^N(x) \vee T_A^N(y) - T_B^N(xy),$$

$$\overline{I}_B^N(xy) = I_A^N(x) \wedge I_A^N(y) - I_B^N(xy), \quad \overline{F}_B^N(xy) = F_A^N(x) \wedge F_A^N(y) - F_B^N(xy),$$

for all $xy \in Y \times Y$.




Figure 3: Domination in bipolar neutrosophic graph




Figure 4: Bipolar single valued neutrosophic graph G according to [11]




Figure 5: Complement of G according to [11]

We illustrate Definitions 5.1-5.2 [11] by the following example.

Example 4.1. The complement \overline{G} of G is obtained by using Definitions 5.1-5.2 [11] as shown in Fig. 5. It can be seen from Fig. 5 that $I_B^p(yz), f_B^p(yz), I_B^p(yt), f_B^p(yt) \notin [0, 1]$ and $I_B^n(yz), f_B^n(yz), I_B^n(yt), f_B^n(yt) \notin [-1, 0]$. Hence \overline{G} is not a bipolar single valued neutrosophic graph. ■

4. Conclusion and Future Work

Bipolar fuzzy graph theory has many applications in science and technology. A bipolar neutrosophic graph is a generalization of the notion bipolar fuzzy graph. We have introduced the idea of bipolar neutrosophic graph and operations on bipolar neutrosophic graphs. We have investigated the dominating and independent sets of certain graph products. Two applications of bipolar neutrosophic sets and bipolar neutrosophic graphs are studied in chemical industry and construction of radio channels. We are planning to extend our research of fuzzification to (1) Bipolar fuzzy rough graphs; (2) Bipolar fuzzy rough hypergraphs, (3) Bipolar fuzzy rough neutrosophic graphs, and (4) Decision support systems based on bipolar neutrosophic graphs.

References

- [1] AKRAM, M., *Bipolar fuzzy graphs*, Information Sciences, 181(24)(2011), 5548-5564.
- [2] AKRAM, M., *Bipolar fuzzy graphs with application*, Knowledge Based Systems, 39(2013), 1-8.
- [3] AKRAM, M., DUDEK, W. A. , *Regular bipolar fuzzy graphs*, Neural Computing and Applications, 21(1)(2012), 197-205.
- [4] AKRAM, M., SITARA, M., *Bipolar neutrosophic graph structures*, Journal of the Indonesian Mathematical Society, 23(1)(2017), 55-76.

- [5] AKRAM, M., NASIR, N., *Certain Bipolar Neutrosophic Competition Graphs*, Journal of the Indonesian Mathematical Society, 23(2)(2017), 1-23.
- [6] AKRAM, M., LUQMAN, A., *Novel applications of bipolar neutrosophic hypergraphs*, Journal of intelligent and fuzzy systems, 2017 (In Press).
- [7] AKRAM, M., SHUM, K. P., *Bipolar neutrosophic planar graphs*, Journal of Mathematical Research with Applications, 2017 (In Press).
- [8] BHATTACHARYA, P., *Some remarks on fuzzy graphs*, Pattern Recognition Letters, 6(5)(1987), 297-302.
- [9] BHUTANI, K. R., BATTOU, A., *On M -strong fuzzy graphs*, Information Sciences, 155(1)(2003), 103-109.
- [10] BHUTANI, K. R., ROSENFELD, A., *Strong arcs in fuzzy graphs*, Information Sciences, 152(2003), 319-322.
- [11] BROUMI, S., TALEA, M., BAKALI, A., SMARANDACHE, F., *On bipolar single valued neutrosophic graphs*, Journal of New Theory, 11(2016), 84-102.
- [12] DELI, I., ALI, M., SMARANDACHE, F., *Bipolar neutrosophic sets and their application based on multi-criteria decision making problems*, In Advanced Mechatronic Systems (ICAMechS), International Conference IEEE, (2015), 249-254.
- [13] DELI, I., ÇAĞMAN, N., *Intuitionistic fuzzy parameterized soft set theory and its decision making*, Applied Soft Computing, 28(2015) 109-113.
- [14] KAUFMANN, A., *Introduction à la théorie des sous-ensembles flous à l'usage des ingénieurs (fuzzy sets theory)*, Masson, Paris, 1975.
- [15] MATHEW, S., SUNITHA, M., *Strongest strong cycles and theta fuzzy graphs*, IEEE Transactions on Fuzzy Systems, 21(6)(2013), 1096-1104.
- [16] MORDESON J. N., CHANG-SHYH, P., *Operations on fuzzy graphs*, Information Sciences, 79(3)(1994), 159-170.
- [17] MORDESON, J. N., MALIK, D. S., RICHARDS, C. D., TREBBIAN, J.A., BOYCE, M. A., BYRNE M. P., COUSINO, B.J., *Fuzzy graphs and complementary fuzzy graphs*, The Journal of Fuzzy Mathematics, 24(2)(2016) 271-288.
- [18] NAGOORGANI, A., CHANDRASEKARAN, V. T., *Domination in fuzzy graphs*, Advances in fuzzy sets and system, I(1)(2006) 17-26.
- [19] PENG, J. J., WANG, J. Q., ZHANG, H. Y., CHEN, X. H., *An outranking approach for multi-criteria decision-making problems with simplified neutrosophic set*, Applied Soft Computing, 25(2014), 336-346.

- [20] ROSENFELD, A., *Fuzzy graphs*, in: L.A. Zadeh, K.S. Fu, M. Shimura (Eds.), *Fuzzy Sets and their Applications*, Academic Press, New York, 513(1975), 77-95.
- [21] SARWAR, M., AKRAM, M., *Novel concepts bipolar fuzzy competition graphs*. *JJournal of Applied Mathematics and Computing*, (2016), DOI 10.1007/s12190-016-1021-z.
- [22] SMARANDACHE, F., *A unifying field in logics: Neutrosophic logic*, *Philosophy*, 1 – 141, 1999.
- [23] SMARANDACHE, F., *A Unifying Field in Logics: Neutrosophic Logic. Neutrosophy, Neutrosophic Set, Neutrosophic Probability: Neutrosophic Logic. Neutrosophy, Neutrosophic Set, Neutrosophic Probability*, Infinite Study, 2005.
- [24] SOMASUNDARAM, A., SOMASUNDARAM, S., *Domination in fuzzy graphs-I*, *Pattern Recognition Letters*, 19(9)(1998), 787-791.
- [25] YE, J., *Trapezoidal neutrosophic set and its application to multiple attribute decision-making*, *Neural Computing and Applications*, 26(5)(2015), 1157-1166.
- [26] YE, J., *Vector similarity measures of simplified neutrosophic sets and their application in multicriteria decision making*, *International Journal of Fuzzy Systems*, 16(2)(2014), 204-215, 2014.
- [27] ZADEH, L. A., *Fuzzy sets*, *Information and control*, 8(3)(1965), 338-353.
- [28] ZADEH, L. A., *Similarity relations and fuzzy orderings*, *Information sciences*, 3(2)(1971), 177-200.
- [29] ZHANG, W.–R., *Bipolar fuzzy sets and relations: a computational framework for cognitive modeling and multiagent decision analysis*, In *Proc. of IEEE conf. Fuzzy Information Processing Society Biannual Conference*, 1994, 305-309.

Accepted: 12.04.2017