

Article

Symmetry in Hyperstructure: Neutrosophic Extended Triplet Semihypergroups and Regular Hypergroups

Xiaohong Zhang ^{1,*}, Florentin Smarandache ² and Yingcang Ma ³

¹ Department of Mathematics, Shaanxi University of Science & Technology, Xi'an 710021, China

² Department of Mathematics, University of New Mexico, 705 Gurley Avenue, Gallup, NM 87301, USA; smarand@unm.edu

³ School of Science, Xi'an Polytechnic University, Xi'an 710048, China; mayingcang@xpu.edu.cn

* Correspondence: zhangxiaohong@sust.edu.cn

Received: 28 August 2019; Accepted: 25 September 2019; Published: 1 October 2019

Abstract: The symmetry of hyperoperation is expressed by hypergroup, more extensive hyperalgebraic structures than hypergroups are studied in this paper. The new concepts of neutrosophic extended triplet semihypergroup (NET- semihypergroup) and neutrosophic extended triplet hypergroup (NET-hypergroup) are firstly introduced, some basic properties are obtained, and the relationships among NET- semihypergroups, regular semihypergroups, NET-hypergroups and regular hypergroups are systematically are investigated. Moreover, pure NET-semihypergroup and pure NET-hypergroup are investigated, and a structure theorem of commutative pure NET-semihypergroup is established. Finally, a new notion of weak commutative NET-semihypergroup is proposed, some important examples are obtained by software MATLAB, and the following important result is proved: every pure and weak commutative NET-semihypergroup is a disjoint union of some regular hypergroups which are its subhypergroups.

Keywords: hypergroup; semihypergroup; neutrosophic extended triplet group; neutrosophic extended triplet semihypergroup (NET-semihypergroup); NET-hypergroup

1. Introduction and Preliminaries

As a generalization of traditional algebraic structures, hyper algebraic structures (or hypercompositional structures) have been extensively studied and applied [1–7]. Especially, hypergroups and semihypergroups are basic hyper structures which are extensions of groups and semigroups [8]. In fact, hypergroups characterize the symmetry of hyperoperations.

On the other hand, as an extension of fuzzy set and intuitionistic fuzzy set, the concept of neutrosophic set firstly proposed by F. Smarandache in [9], has been applied to many fields [10–12]. Moreover, as an application of the ideal of neutrosophic sets, a new notion of neutrosophic triplet group (NTG) was proposed by F. Smarandache and Ali in [13], while the new notion of neutrosophic extended group (NETG) was proposed by Smarandache in [14]. Furthermore, the basic properties and structural characteristics of neutrosophic extended groups (NETGs) are studied in [15,16]; the closed connection between between NETG and regular semigroup investigated, and the new notion of neutrosophic extended triplet Abel-Grassmann's Groupoid is proposed in [17]; the decomposition theorem of NETG is proved in [18]; the generalized neutrosophic extended groups are presented in [19]; the relationship and difference between NETGs and generalized groups are systematically studied in [20]. From these research results, we know that NETG is a typical algebraic system with important research value.

In this paper, we combine the two directions mentioned above to study the hyperalgebraic structures related to neutrosophic extended triplet groups (NETGs), which can be regarded as a further development of the research ideas in [21].

At first, we recall some concepts and results on hypergroups, semigroups and NETGs.

Let H be a non-empty set and $P^*(H)$ the set of all non-empty subsets of H . A map $\circ: H \times H \rightarrow P^*(H)$ is called (binary) hyperoperation (or hypercomposition), and (H, \circ) is called a hypergroupoid. If $A, B \in P^*(H)$, $x \in H$, then

$$A \circ B = \bigcup_{a \in A, b \in B} (a \circ b), A \circ x = A \circ \{x\}, x \circ B = \{x\} \circ B.$$

Definition 1. ([1–4]) Let (H, \circ) be a hypergroupoid. If $(\forall x, y, z \in H) (x \circ y) \circ z = x \circ (y \circ z)$, then (H, \circ) is called a semihypergroup. That is,

$$\bigcup_{u \in x \circ y} (u \circ z) = \bigcup_{v \in y \circ z} (x \circ v).$$

For a semihypergroup (H, \circ) , if $(\forall x, y \in H) x \circ y = y \circ x$, then we call that H is commutative.

Note that, if (H, \circ) is a semihypergroup, then $(A \circ B) \circ C = A \circ (B \circ C)$, $\forall A, B, C \in P^*(H)$.

Definition 2. ([1–4]) Assume that (H, \circ) is a semihypergroup. (1) If $a \in H$ satisfies $(\forall x \in H) |a \circ x| = |x \circ a| = 1$, then a is called to be scalar. (2) If $e \in H$ satisfies $(\forall x \in H) x \circ e = e \circ x = \{x\}$, then e is called scalar identity. (3) If $e \in H$ satisfies $(\forall x \in H) x \in (e \circ x) \cap (x \circ e)$, then e is called identity. (4) Let $a, b \in H$. If there exists an identity $e \in H$ satisfies $e \in (a \circ b) \cap (b \circ a)$, then b is called an inverse of a . (5) If $0 \in H$ satisfies $(\forall x \in H) x \circ 0 = 0 \circ x = \{0\}$, then 0 is called zero element.

Definition 3. ([1–4]) Let (H, \circ) be a semihypergroup. (1) If $(\forall x \in H) a \circ H = H \circ a = H$ (reproductive axiom), then (H, \circ) is called a hypergroup. (2) If (H, \circ) is a hypergroup and (H, \circ) has at least one identity and each element has at least one inverse, then (H, \circ) is called to be regular.

Definition 4. ([1–4]) Let (H, \circ) be a semihypergroup. If $x \in H$ satisfies $x \in x \circ H \circ x$, i.e., there exists an element $y \in H$, $x \in x \circ y \circ x$, then x is said to be regular. If $(\forall x \in H) x$ is regular, then (H, \circ) is called to be regular.

Note that, Every regular semigroup is a regular semihypergroup, and every hypergroup is a regular semihypergroup.

Definition 5. ([14]) Let N be a non-empty set, and $*$ a binary operation on N . If $(\forall a \in N)$ there exist $neut(a) \in N$, $anti(a) \in N$ satisfy

$$\begin{aligned} neut(a) * a &= a * neut(a) = a, \text{ and} \\ anti(a) * a &= a * anti(a) = neut(a). \end{aligned}$$

Then N is called a neutrosophic extended triplet set (NETS). Moreover, for $a \in N$, $(a, neut(a), anti(a))$ is called a neutrosophic extend triplet, $neut(a)$ is called an extend neutral of “ a ”, and $anti(a)$ is called an opposite of “ a ”.

For a neutrosophic extended triplet set N , $a \in N$, the set of $neut(a)$ is denoted by $\{neut(a)\}$, and the set of $anti(a)$ is denoted by $\{anti(a)\}$.

Definition 6. ([13,14]) Let $(N, *)$ be a NETS. If $(N, *)$ is a semigroup, then $(N, *)$ is called to be a neutrosophic extended triplet group (NETG).

About some basic properties of neutrosophic extended triplet groups, please see [15,17,20].

2. Neutrosophic Extended Triplet Semihypergroups (NET-Semihypergroups) and Neutrosophic Extended Triplet Hypergroups (NET-Hypergroups)

In this section, we propose the new concepts of neutrosophic extended triplet semihypergroup (NET-semihypergroup) and neutrosophic extended triplet hypergroup (NET-hypergroup), and give some typical examples to illustrate their wide representativeness.

Definition 7. Let $(H, *)$ be a semihypergroup (i.e., $*$ be a binary hyperoperation on nonempty set H such that $(x*y)*z = x*(y*z)$, for all $x, y, z \in H$). $(H, *)$ is called a neutrosophic extended triplet semihypergroup (shortened form, NET-semihypergroup), if for every $x \in H$, there exist $neut(x)$ and $anti(x)$ such that

$$x \in (neut(x)*x) \cap (x*neut(x)), \text{ and}$$

$$neut(x) \in (anti(x)*x) \cap (x*anti(x)).$$

Here, we call that $(x, neut(x), anti(x))$ to be a hyper-neutrosophic-triplet.

Example 1. Denote $H = \{a, b, c\}$, define hyperoperations $*$ on H as shown in Table 1. We can verify that $(H, *)$ is semihypergroup by software MATLAB (see Figural 1).

Table 1. The hyperoperation $*$ on H .

$*$	a	b	c
a	a	$\{a, b\}$	$\{a, b, c\}$
b	a	$\{a, b\}$	$\{a, b, c\}$
c	a	$\{a, b\}$	c

Moreover,

$$a \in (a*a) \cap (a*a);$$

$$b \in (b*b) \cap (b*b);$$

$$c \in (c*c) \cap (c*c).$$

This means that $(H, *)$ is neutrosophic extended triplet semihypergroup (NET-semihypergroup) and (a, a, a) , (b, b, b) , (c, c, c) are hyper-neutrosophic-triplets.




Figure 1. A program by Matlab to verify hyperoperation.

Example 2. Denote $H = \{a, b, c, d\}$, define hyperoperations $*$ on H as shown in Table 2. We can verify that $(H, *)$ is semihypergroup by software MATLAB (see Figural 2).

Table 2. The hyperoperation $*$ on H .

$*$	a	b	c	d
a	$\{a, b\}$	$\{a, b\}$	$\{c, d\}$	$\{c, d\}$
b	$\{a, b\}$	$\{a, b\}$	$\{c, d\}$	$\{c, d\}$
c	$\{c, d\}$	$\{c, d\}$	a	b
d	$\{c, d\}$	$\{c, d\}$	b	a




Figure 2. Verify hyperoperation by Matlab.

Moreover,

$$a \in (a*a) \cap (a*a); a \in (b*a) \cap (a*b), b \in (b*a) \cap (a*b).$$

$$b \in (b*b) \cap (b*b).$$

$$c \in (a*c) \cap (c*a), a \in (c*c) \cap (c*c); c \in (b*c) \cap (c*b), b \in (d*c) \cap (c*d).$$

$$d \in (a*d) \cap (d*a), a \in (d*d) \cap (d*d); d \in (b*d) \cap (d*b), b \in (c*d) \cap (d*c).$$

This means that $(H, *)$ is neutrosophic extended triplet semihypergroup (NET-semihypergroup) and (a, a) , (a, b, b) , (b, b, b) , (c, a, c) , (c, b, d) , (d, a, d) , (d, b, c) are hyper-neutrosophic-triplets.

Remark 1. From Example 2 we know that $neut(x)$ may be not unique for an element x in a neutrosophic extended triplet semihypergroup (NET-semihypergroup). In fact, in Example 2, we have

$$\{neut(a)\} = \{a, b\}, neut(b) = b, \{neut(c)\} = \{a, b\}, \{neut(d)\} = \{a, b\}.$$

Example 3. Let H be the set of all nonnegative integers, and define a hyperoperation $*$ on H as following:

$$x*y = \{z \in H \mid z \geq \max\{x, y\}\}.$$

For examples,

$$3*5 = \{5, 6, 7, 8, \dots\}; 9*9 = \{9, 10, 11, 12, \dots\}; 2019*0 = \{2019, 2020, 2021, 2022, \dots\}.$$

Then $(H, *)$ is a commutative semihypergroup. Moreover, for any $x \in H$, we have

$$x \in (x*x) \cap (x*x); x \in (x*x) \cap (x*x).$$

This means that $(H, *)$ is a neutrosophic extended triplet semihypergroup (NET-semihypergroup). In fact, we have

$$neut(0)=0; \{neut(1)\}=\{0,1\}; \{neut(2)\}=\{0, 1, 2\}; \{neut(3)\}=\{0, 1, 2, 3\}...$$

Example 4. Let R be the set of all real numbers, and Z the set of integers. We use the modulo of real numbers (that we denote by mod_R) in the following way:

$$\forall a, b \in R, \text{ then } a = b \pmod{6}, \text{ if and only if } a - b = 6n, \text{ where } n \text{ is an integer.}$$

For examples, $14.73 = 2.73 \pmod{6}$, since $14.73 - 2.73 = 12 = 6 \times 2$; but $18 \neq 15 \pmod{6}$, since $18 - 15 = 3 \neq 6n$ with n integer. Now, we define a hyperoperation $\#$ on R as following:

$$a\#b = \{x \in R \mid x = 4ab \pmod{6}\}.$$

Then $(R, \#)$ is a commutative semihypergroup, since $a\#b = b\#a = 4ab \pmod{6}$, and associative because:

$$(a\#b)\#c = (4ab)\#c = 4(4ab)c = 16abc \pmod{6}, \text{ and}$$

$$a\#(b\#c) = a\#(4bc) = 4a(4bc) = 16abc \pmod{6}.$$

Moreover, for any $a \in R$, we have

- (1) when $a=0$, $(a, 6m, r)$ are hyper-triplets for any integer number m and real number r ;
- (2) when $a \neq 0$, $\left(a, \frac{1}{4} + \frac{3m}{2a}, \frac{1}{16a} + \frac{3m}{8a} + \frac{3n}{2a}\right)$ are hyper-neutrosophic-triplets for any integer numbers m, n .

This means that $(R, \#)$ is a neutrosophic extended triplet semihypergroup (NET-semihypergroup), and infinitely many $\text{neut}(a)$ and infinitely many $\text{anti}(a)$ for any element a in R .

Remark 2. The following example shows that a sub-semihypergroup of a NET-semihypergroup may be not a NET-semihypergroup.

Example 5. Denote $H = \{a, b, c, d, e\}$, define hyperoperations $*$ on H as shown in Table 3. We can verify that $(H, *)$ is semihypergroup by software MATLAB (see Figural 3).

Table 3. The hyperoperation $*$ on H .

$*$	a	b	c	d	e
a	a	a	a	d	$\{a, b, c, d, e\}$
b	a	$\{a, b\}$	$\{a, c\}$	d	$\{a, b, c, d, e\}$
c	a	a	a	d	$\{a, b, c, d, e\}$
d	d	d	d	d	$\{a, b, c, d, e\}$
e	$\{a, b, c, d, e\}$	$\{a, b, c, d, e\}$	$\{a, b, c, d, e\}$	$\{a, b, c, d, e\}$	$\{a, b, c, d, e\}$




Figure 3. Verify the hyperoperation by Matlab.

Moreover, $(a, a, a), (a, e, e), (b, b, b), (b, e, e), (c, e, e), (d, d, d), (d, e, e), (e, e, e), (e, a, e), (e, b, e), (e, c, e), (e, d, e)$ are hyper-neutrosophic-triplets. This means that $(H, *)$ is a NET-semihypergroup. For $S = \{a, b, c\} \subseteq H$, $(S, *)$ is sub-semihypergroup of $(H, *)$. But, $(S, *)$ is not a NET-semihypergroup.

Remark 3. For the traditional algebraic structures, we have the conclusion that any group must be a neutrosophic extended triplet group (NETG). For hyper algebraic structures, we know from Example 1 that a NET-semihypergroup is not necessarily a hypergroup (since $a^*H \neq H$ in Example 1). Moreover, the following example shows that a hypergroup may be not a NET-semihypergroup. Therefore, hypergroup and NET-semihypergroup are two non-inclusion hyperalgebraic systems.

Example 6. Denote $H = \{1, 2, 3\}$, define hyperoperations $*$ on H as shown in Table 4. We can verify that $(H, *)$ is semihypergroup by software MATLAB.

Table 4. The hyperoperation $*$ on H .

$*$	1	2	3
1	2	2	{1, 3}
2	{1, 2, 3}	{2, 3}	{1, 2, 3}
3	2	{1, 2, 3}	{1, 3}

Moreover,

$$1^*H = H^*1 = H, 2^*H = H^*2 = H, 3^*H = H^*3 = H.$$

This means that $(H, *)$ is a hypergroup. But, for $1 \in H$, we cannot find $x, y \in H$ such that $1 \in (x^*1) \cap (1^*x)$, and $x \in (y^*1) \cap (1^*y)$. That is, $(H, *)$ is not a NET-semihypergroup.

Definition 8. Let $(H, *)$ be a semihypergroup. $(H, *)$ is called a neutrosophic extended triplet hypergroup (shortened form, NET-hypergroup), if $(H, *)$ is both a NET-semihypergroup and a hypergroup.

Obviously, the NET-semihypergroups in Example 2 and Example 5 are all NET-hypergroups. And, the following proposition is true (the proof is omitted).

Proposition 1. Every regular hypergroup is a NET-hypergroup.

The NET-hypergroup in Example 2 is not a regular hypergroup, it shows that the inverse of Proposition 1 is not true.

Proposition 2. Let $(H, *)$ be a NET-semihypergroup (or a NET-hypergroup). Then $(H, *)$ is a regular semihypergroup.

Proof. Assume that $(H, *)$ is a NET-semihypergroup. For any $x \in H$, by Definition 7 we get that there exist $neut(x)$ and $anti(x)$ such that

$$x \in (neut(x)^*x) \cap (x^*neut(x)), \text{ and } neut(x) \in (anti(x)^*x) \cap (x^*anti(x)).$$

Then,

$$x \in neut(x)^*x \subseteq (x^*anti(x))^*x.$$

That is, $x \in x^*anti(x)^*x$. From this, by Definition 4, we know that $(H, *)$ is a regular semihypergroup.

If $(H, *)$ is a NET-hypergroup, by Definition 8, it follows that $(H, *)$ is a NET-semihypergroup. Then, by the proof above, $(H, *)$ is a regular semihypergroup. \square

The following example shows that the inverse of Proposition 2 is not true. Moreover, it also shows that a regular semihypergroup may be not a hypergroup.


Example 7. Denote $H = \{a, b, c\}$, define hyperoperations $*$ on H as shown in Table 5. We can verify that $(H, *)$ is semihypergroup.

Table 5. The hyperoperation $*$ on H .

$*$	a	b	c
a	a	a	a
b	$\{a, b, c\}$	$\{a, b, c\}$	$\{a, b, c\}$
c	$\{a, b, c\}$	$\{a, b\}$	$\{a, b\}$

Moreover, $a \in a*a*a$; $b \in b*b*b$; $c \in c*a*c$. This means that $(H, *)$ is a regular semihypergroup. But it is not a NET-semihypergroup, since there is not any $x \in H$ such that $c \in x*c$ and $c \in c*x$. Obviously, $(H, *)$ is not a hypergroup.

Therefore, the relationships among semihypergroup, NET-semihypergroup, NET-hypergroup, (regular) hypergroup and regular semihypergroup can be expressed by Figure 4.

**Figure 4.** The relationships among some kinds of semihypergroups.

For basic properties of NET-semihypergroups and NET-hypergroups, we can get following results.

Theorem 1. Let $(H, *)$ be a semihypergroup. Then

- (1) if $(H, *)$ is commutative NET-semihypergroup, then for any $x \in H$ and hyper-neutrosophic-triplet $(x, neut(x), anti(x))$, there exists $p \in neut(x)*neut(x)$ and $q \in anti(x)*neut(x)$ such that (x, p, q) is also a hyper-neutrosophic-triplet.
- (2) if $(H, *)$ is commutative NET-semihypergroup, then for any $x \in H$ and $neut(x) \in \{neut(x)\}$, there exists $p \in neut(x)*neut(x)$ such that $p \in \{neut(x)\}$.
- (3) if $(H, *)$ is NET-semihypergroup and $x \in H$ is scalar, then $|\{neut(x)\}| = 1$, that is, the neutral element of x is unique; Moreover, if x is scalar, then $neut(x)*neut(x) = neut(x)$.
- (4) if $(H, *)$ is commutative hypergroup, then $(H, *)$ is NET-hypergroup.

Proof. (1) Assume that $x \in H$ and $(x, neut(x), anti(x))$ is a hyper-neutrosophic-triplet. By Definition 7:

$$x \in (neut(x)*x) \cap (x*neut(x)), \text{ and } neut(x) \in (anti(x)*x) \cap (x*anti(x)).$$

Since $(H, *)$ is commutative, then:

$$x \in neut(x)*x \subseteq neut(x)*(neut(x)*x) = (neut(x)*neut(x))*x = x*(neut(x)*neut(x)).$$

This means that there exists $p \in neut(x)*neut(x)$ such that $x \in p*x = x*p$. Moreover:

$$p \in neut(x)*neut(x) \subseteq (x*anti(x))*neut(x) = x*(anti(x)*neut(x)) = (anti(x)*neut(x))*x.$$

It follows that there exists $q \in anti(x)*neut(x)$ such that $p \in q*x = x*q$. By Definition 7 we know that (x, p, q) is also a hyper-neutrosophic-triplet.

(2) It follows from (1).

(3) Suppose that $x \in H$ and x is scalar. Using Definition 2, $|x^*a| = |a^*x| = 1$ for any $a \in H$. From this, for a hyper-neutrosophic-triplet $(x, neut(x), anti(x))$, applying Definition 7, we have:

$$x = neut(x)^*x = x^*neut(x), \text{ and } neut(x) = anti(x)^*x = x^*anti(x).$$

Assume $p_1, p_2 \in \{neut(x)\}$, then there exists $q_1, q_2 \in H$ such that:

$$x = p_1^*x = x^*p_1, p_1 = q_1^*x = x^*q_1; x = p_2^*x = x^*p_2, p_2 = q_2^*x = x^*q_2.$$

Then:

$$p_1 = q_1^*x = q_1^*(x^*p_2) = (q_1^*x)^*p_2 = p_1^*p_2;$$

$$p_2 = x^*q_2 = (x^*p_1)^*q_2 = (x^*(q_1^*x))^*q_2 = (x^*q_1)^*(x^*q_2) = p_1^*p_2.$$

It follows that $p_1 = p_2$ and $p_1 = p_1^*p_1$. That is, $|\{neut(x)\}| = 1$ and $neut(x)^*neut(x) = neut(x)$.

(4) Let $(H, *)$ be a commutative hypergroup. By Definition 3, for any $x \in H$, $x^*H = H^*x = H$. Then, for any $x \in H$, there exists $h \in H$ such that $x = h^*x = x^*h$. Moreover, for $h \in H$, there exists $u \in H$ such that $h = u^*x = x^*u$. Thus, (x, h, u) is a hyper-neutrosophic-triplet, and it means that $(H, *)$ is a NET-semihypergroup by Definition 7. On the other hand, since $(H, *)$ is a hypergroup, so $(H, *)$ is a NET-hypergroup by Definition 8.

3. Pure NET-semihypergroups and Regular hypergroups

In this section, we discuss some properties of NET-semihypergroups. We'll propose the new notion of pure NET-semihypergroup, investigate the structure of pure NET-semihypergroups.

Definition 9. Let $(H, *)$ be a NET-semihypergroup. $(H, *)$ is called a pure NET-semihypergroup, if for every $x \in H$, there exist $neut(x)$ and $anti(x)$ such that

$$x = (neut(x)^*x) \cap (x^*neut(x)), \text{ and } neut(x) = (anti(x)^*x) \cap (x^*anti(x)).$$

Obviously, the following proposition is true and the proof is omitted.

Proposition 3. (1) Every neutrosophic extended triplet group (NETG) is pure NET-semihypergroup. (2) If $(H, *)$ is a pure NET-semihypergroup and the hyper operation $*$ is commutative, then for every $x \in H$, there exists $y, z \in H$ such that

$$x = y^*x = x^*y, \text{ and } y = z^*x = x^*z.$$

Example 8. Denote $H = \{a, b, c\}$, define hyperoperations $*$ on H as shown in Table 6. We can verify that $(H, *)$ is semihypergroup.

Table 6. The hyperoperation $*$ on H .

$*$	a	b	c
a	a	$\{a, b, c\}$	$\{a, b, c\}$
b	$\{a, b, c\}$	c	b
c	$\{a, b, c\}$	b	c

Moreover,

$$a = (a^*a) \cap (a^*a); b = (c^*b) \cap (b^*c), c = (b^*b) \cap (b^*b); c = (c^*c) \cap (c^*c).$$

This means that $(H, *)$ is a pure NET-semihypergroup.

Example 9. Denote $H = \{a, b, c, d, e\}$, define hyperoperations $*$ on H as shown in Table 7. We can verify that $(H, *)$ is semihypergroup.

Table 7. The hyperoperation $*$ on H .

$*$	a	b	c	d	e
a	a	$\{a, b, c\}$	$\{a, b, c\}$	d	a

b	$\{a, b, c\}$	b	c	d	b
c	$\{a, b, c\}$	c	b	d	c
d	d	d	d	d	d
e	a	b	c	d	e

Moreover:

$$a=(a^*a)\cap(a^*a); b=(b^*b)\cap(b^*b); c=(b^*c)\cap(c^*b), b=(c^*c)\cap(c^*c); d=(d^*d)\cap(d^*d); e=(e^*e)\cap(e^*e).$$

This means that $(H, *)$ is a pure NET-semihypergroup.

Remark 4. From Example 8 and Example 9, we have:

$$a=(a^*a)\cap(a^*a);$$

$$a\in(b^*a)\cap(a^*b), b\in(b^*a)\cap(a^*b); a\in(c^*a)\cap(a^*c), c\in(c^*a)\cap(a^*c).$$

This means that $\{neut(a)\} = \{a, b, c\}$. But, $b\in\{neut(a)\}$ and $c\in\{neut(a)\}$ are different to $a\in\{neut(a)\}$, since one is “ \in ” and the other is “ $=$ ”. In order to clearly express the difference between the two kinds of neutral elements, we introduce a new concept: pure neutral element.

Definition 10. Let $(H, *)$ be a NET-semihypergroup and $x\in H$. An element $y\in H$ is called a pure neutral element of the element x , if there exist $z\in H$ such that:

$$x = y^*x = x^*y, \text{ and } y = z^*x = x^*z.$$

Here, we denote y by $pneut(x)$.

Proposition 4. Let $(H, *)$ be a NET-semihypergroup and $x\in H$. If there exists a pure neutral element of x , then the pure neutral element of x , that is, $pneut(x)$, is unique.

Proof. Assume that there exists two pure neutral elements y_1, y_2 for $x\in H$. Then there exists $z_1, z_2\in H$ such that:

$$x = y_1^*x = x^*y_1, \text{ and } y_1 = z_1^*x = x^*z_1;$$

$$x = y_2^*x = x^*y_2, \text{ and } y_2 = z_2^*x = x^*z_2.$$

Therefore,

$$y_1 = z_1^*x = z_1^*(x^*y_2) = (z_1^*x)^*y_2 = y_1^*y_2;$$

$$y_2 = x^*z_2 = (x^*y_1)^*z_2 = (x^*(z_1^*x))^*z_2 = (x^*z_1)^*(x^*z_2) = y_1^*y_2.$$

Hence, $y_1 = y_2$. That is, $pneut(x)$ is unique. \square

By the proof of Proposition 4, we know that $y_1 = y_2 = y_1^*y_2$, it follows that $y_1 = y_1^*y_1$. Therefore, we have the following corollary.

Corollary 1. Let $(H, *)$ be a NET-semihypergroup and $x\in H$. If there exists a pure neutral element of x , then the pure neutral element of x is idempotent, that is, $pneut(x)^*pneut(x) = pneut(x)$.

Remark 5. From Proposition 4, we know that the pure neutral element of an element x is unique when there exists one pure neutral element of x . Particularly, for commutative pure NET-semihypergroups, applying Proposition 3 (2), we get following Proposition 5 (the proof is omitted).

Proposition 5. Let $(H, *)$ be a commutative pure NET-semihypergroup. Then for any $x\in H$, $pneut(x)$ is unique.

Proposition 6. Let $(H, *)$ be a commutative pure NET-semihypergroup. Then for any $x, y\in H$, $pneut(x^*y) = pneut(x)^*pneut(y)$ when $|x^*y|=1$. Moreover, if $pneut(x) = z_1^*x = x^*z_1$ and $pneut(y) = z_2^*y = y^*z_2$, $z_1, z_2\in H$, then:

$$pneut(x^*y) = (z_1^*z_2)^*(x^*y) = (x^*y)^*(z_1^*z_2).$$

Proof. Assume that $x, y\in H$ and $|x^*y|=1$. Since $(H, *)$ be a commutative pure NET-semihypergroup, then:

$$\begin{aligned}
(x*y)*(pneut(x)*pneut(y)) &= (x*y)*(pneut(y)*pneut(x)) \\
&= x*(y*pneut(y))*pneut(x) \\
&= x*y*pneut(x) \\
&= (x*pneut(x))*y \\
&= x*y; \\
(pneut(x)*pneut(y))*(x*y) &= (pneut(y)*pneut(x))*(x*y) \\
&= pneut(y)*(pneut(x)*x)*y \\
&= pneut(y)*x*y \\
&= x*(pneut(y)*y) \\
&= x*y.
\end{aligned}$$

On the other hand, assume that $(x, pneut(x), anti(x))$ and $(y, pneut(y), anti(y))$ are hyper-neutrosophic-triplets, then:

$$\begin{aligned}
(x*y)*(anti(x)*anti(y)) &= (x*y)*(anti(y)*anti(x)) \\
&= x*(y*anti(y))*anti(x) \\
&= x*pneut(y)*anti(x) \\
&= (x*anti(x))*pneut(y) \\
&= pneut(x)*pneut(y); \\
(anti(x)*anti(y))*(x*y) &= (anti(x)*anti(y))*(y*x) \\
&= anti(x)*(anti(y)*y)*x \\
&= anti(x)*pneut(y)*x \\
&= (anti(x)*x)*pneut(y) \\
&= pneut(x)*pneut(y).
\end{aligned}$$

Applying Proposition 5 we get that $pneut(x*y) = pneut(x)*pneut(y)$.

Moreover, assume $pneut(x) = z_1*x = x*z_1$, $pneut(y) = z_2*y = y*z_2$. Then, by commutativity of the hyper operation $*$:

$$\begin{aligned}
(z_1*z_2)*(x*y) &= (z_1*x)*(z_2*y) \\
&= pneut(x)*pneut(y) \\
&= pneut(x*y); \\
(x*y)*(z_1*z_2) &= (x*z_1)*(y*z_2) \\
&= pneut(x)*pneut(y) \\
&= pneut(x*y).
\end{aligned}$$

Therefore, the proof is completed.

Theorem 2. Let $(H, *)$ be a commutative pure NET-semihypergroup and H satisfies:

$$\forall x, y \in H, pneut(x) = pneut(y) \Rightarrow |x*y| = 1. \quad (C1)$$

Define a binary relation \approx on H as following:

$$\forall x, y \in H, x \approx y \text{ if and only if } pneut(x) = pneut(y).$$

Then:

- (1) The binary relation is a equivalent relation on H ;
- (2) For any $x \in H$, $[x]_{\approx}$ is a sub-NET-semihypergroup of H , where $[x]_{\approx}$ is the equivalent class of x based on equivalent relation \approx ;
- (3) For any $x \in H$, $[x]_{\approx}$ is a regular hypergroup.

Proof. (1) It is obviously.

(2) Assume $a, b \in [x]_{\approx}$, then $pneut(a) = pneut(b) = pneut(x)$. Applying Proposition 6 and Corollary 1, we have

$$\begin{aligned} pneut(a*b) &= pneut(a)*pneut(b) \\ &= pneut(x)*pneut(x) \\ &= pneut(x). \end{aligned}$$

This means that $[x]_{\approx}$ is closed on the hyper operation $*$.

Moreover, by Corollary 1, we have $pneut(x)*pneut(x) = pneut(x)$. From this and using Proposition 5, we get that $pneut(pneut(x)) = pneut(x)$. It follows that $pneut(a) \in [x]_{\approx}$ for any $a \in [x]_{\approx}$. Moreover, assume that $a \in [x]_{\approx}$, by the definition of commutative pure NET-semihypergroup, there exists $r \in H$ such that:

$$pneut(a) = r*a = a*r.$$

It follows that:

$$pneut(a) = (r*pneut(a))*a = a*(r*pneut(a)). \quad (C2)$$

Applying Proposition 6 and Corollary 1:

$$\begin{aligned} &pneut(r*pneut(a)) \\ &= pneut(r)*pneut(pneut(a)) \\ &= pneut(r)*pneut(a) \\ &= pneut(r*a) \\ &= pneut(pneut(a)) \\ &= pneut(a). \end{aligned}$$

That is, $pneut(r*pneut(a)) = pneut(a) = pneut(x)$. This means that $r*pneut(a) \in [x]_{\approx}$. Therefore, by (C2), there exists $anti(a)$ (see Definition 7), it is in $[x]_{\approx}$. This means that $[x]_{\approx}$ is a sub-NET-semihypergroup of H .

(3) For any $x \in H$, from (2) we know that $[x]_{\approx}$ is a sub-NET-semihypergroup of H . By the definition of \approx , for any $a \in [x]_{\approx}$, $pneut(a) = pneut(x)$. Then, $a*[x]_{\approx} = [x]_{\approx}$, and $pneut(x)$ is a (local) identity in $[x]_{\approx}$. By Definition 3, we get that $[x]_{\approx}$ is a regular hypergroup. \square

From Theorem 2 we know that for a commutative pure NET-semihypergroup (it satisfies the condition in Theorem 2), it is a union of some regular hypergroups. The following picture (Figure 5) shows this special structure.




Figure 5. The structure of a commutative pure NET-semihypergroups.

Example 10. Denote $H = \{a, b, c, d, e\}$, define hyperoperations $*$ on H as shown in Table 8. We can verify that $(H, *)$ is commutative pure NET-semihypergroup.

Table 8. The hyperoperation $*$ on H .

$*$	a	b	c	d	e
a	a	$\{a, b, c\}$	$\{a, b, c\}$	d	$\{a, d, e\}$
b	$\{a, b, c\}$	b	c	d	$\{b, c, d, e\}$
c	$\{a, b, c\}$	c	b	d	$\{b, c, d, e\}$
d	d	d	d	d	d
e	$\{a, d, e\}$	$\{b, c, d, e\}$	$\{b, c, d, e\}$	d	e

Moreover:

$$\begin{aligned}
 H_1 &= \{a\} = [a]_{\neq}; \\
 H_2 &= \{b, c\} = [b]_{\neq} = [c]_{\neq}; \\
 H_3 &= \{d\} = [d]_{\neq}; \\
 H_4 &= \{e\} = [e]_{\neq};
 \end{aligned}$$

and $H = H_1 \cup H_2 \cup H_3 \cup H_4$, where, $H_i (i=1, 2, 3, 4)$ are regular hypergroups.

Remark 6. The above example shows that a commutative pure NET-semihypergroup may be not a hypergroup (since $d * H \neq H$ in Example 10).

4. Weak Commutative NET-Semihypergroups and Their Structures

In this section, we discuss generalized commutativity in NET-semihypergroups. We propose a new notion of weak commutative NET-semihypergroup, and prove the structure theorem of weak commutative pure NET-semihypergroup (WCP-NET-semihypergroup), which can be regarded as a generalization of Clifford Theorem in semigroup theory.

Definition 11. Let $(H, *)$ be a NET-semihypergroup. $(H, *)$ is called a weak commutative NET-semihypergroup, if for every $x \in H$, every hyper-neutrosophic-triplet $(x, \text{neut}(x), \text{anti}(x))$, the following conditions are satisfied:

$(H, *)$ is called a weak commutative pure NET-semihypergroup (shortly, WCP-NET-semihypergroup), if it both weak commutative and pure.

Obviously, the following proposition is true and the proof is omitted.

Proposition 7. Every commutative NET-semihypergroup is weak commutative.

The following examples show that there exists some weak commutative NET-semihypergroups which are not commutative.

Example 11. Denote $H = \{1, 2, 3, 4, 5, 6, 7, 8\}$, define hyperoperations $*$ on H as shown in Table 9. We can verify that $(H, *)$ is NET-semihypergroup.

Table 9. The hyperoperation $*$ on H .

$*$	1	2	3	4	5	6	7	8
1	1	$\{1, 2\}$	1	1	1	1	1	1
2	$\{1, 2\}$	2	1	1	1	1	1	1
3	1	1	3	4	5	6	7	8
4	1	1	4	3	8	7	6	5
5	1	1	5	7	3	8	4	6
6	1	1	6	8	7	3	5	4
7	1	1	7	5	6	4	8	3
8	1	1	8	6	4	5	3	7

Moreover, $(1, 1, 1)$, $(2, 2, 2)$, $(3, 3, 3)$, $(4, 3, 4)$, $(5, 3, 5)$, $(6, 3, 6)$, $(7, 3, 8)$ and $(8, 3, 7)$ are hyper-neutrosophic-triplets, and $(\forall x \in H) 1 * x = x * 1, 2 * x = x * 2$ and $3 * x = x * 3, 7 * 8 = 8 * 7$. This means that $(H, *)$ is a weak commutative NET-semihypergroup. Since $4 * 5 \neq 5 * 4$, $(H, *)$ is not commutative.

Remark 7. The above example shows that there exists WCP-NET-semihypergroup (by Definition 9, we know that the NET-semihypergroup in Example 11 is pure).

Example 12. Denote $H = \{1, 2, 3, 4, 5, 6, 7, 8, 9\}$, define hyperoperations $*$ on H as shown in Table 10. We can verify that $(H, *)$ is NET-semihypergroup.

Table 10. The hyperoperation $*$ on H .

*	1	2	3	4	5	6	7	8	9
1	2	{1, 3}	3	1	1	1	1	1	1
2	{1, 3}	2	{1, 3}	{1, 2, 3}	{1, 2, 3}	{1, 2, 3}	{1, 2, 3}	{1, 2, 3}	{1, 2, 3}
3	3	{1, 3}	1	{1, 3}	{1, 3}	{1, 3}	{1, 3}	{1, 3}	{1, 3}
4	1	{1, 2, 3}	{1, 3}	4	5	6	7	8	9
5	1	{1, 2, 3}	{1, 3}	5	4	9	8	7	6
6	1	{1, 2, 3}	{1, 3}	6	8	4	9	5	7
7	1	{1, 2, 3}	{1, 3}	7	9	8	4	6	5
8	1	{1, 2, 3}	{1, 3}	8	6	7	5	9	4
9	1	{1, 2, 3}	{1, 3}	9	7	5	6	4	8

Moreover, $(1, 2, 1), (2, 2, 2), (3, 1, 3), (4, 4, 4), (5, 4, 5), (6, 4, 6), (7, 4, 7), (8, 4, 9)$ and $(9, 4, 8)$ are hyper-neutrosophic-triplets, and $(\forall x \in H) 2*x = x*2, 1*x = x*1$ and $4*x = x*4, 8*9 = 9*8$. This means that $(H, *)$ is a weak commutative NET-semihypergroup. Since $5*6 \neq 6*5$, $(H, *)$ is not commutative.

Proposition 8. Let $(H, *)$ be a weak commutative pure NET-semihypergroup (WCP-NET-semihypergroup). Then for any $x \in H$, there exists a pure neutral element of x , and $pneut(x)$ is unique, $pneut(x)*pneut(x) = pneut(x)$.

Proof. For any $x \in H$. Since $(H, *)$ is pure, by Definition 9, there exists hyper-neutrosophic-triplet $(x, neut(x), anti(x))$ such that

$$x = (neut(x)*x) \cap (x*neut(x)), \text{ and } neut(x) = (anti(x)*x) \cap (x*anti(x)).$$

Moreover, since $(H, *)$ is weak commutative, by Definition 11, $neut(x)*x = x*neut(x)$, and $anti(x)*x = x*anti(x)$. Thus

$$x = neut(x)*x = x*neut(x), \text{ and } neut(x) = anti(x)*x = x*anti(x).$$

Therefore, by Definition 10, $neut(x)$ is a pure neutral element of x . Applying Proposition 4 we know that pure neutral element of x is unique. Moreover, using Corollary 1, $pneut(x)*pneut(x) = pneut(x)$. \square

Proposition 9. Let $(H, *)$ be a weak commutative pure NET-semihypergroup (WCP-NET-semihypergroup). Then for any $x, y \in H$, $pneut(x*y) = pneut(x)*pneut(y)$ when $|x*y| = 1$. Moreover, if $pneut(x) = z_1*x = x*z_1$ and $pneut(y) = z_2*y = y*z_2$, $z_1, z_2 \in H$, then

$$pneut(x*y) = (z_2*z_1)*(x*y) = (x*y)*(z_2*z_1).$$

Proof. Since $(H, *)$ be a WCP-NET-semihypergroup, then for any $x, y \in H$ and $|x*y| = 1$, $pneut(x)*y = y*pneut(x)$ by Definition 11. Then

$$x*y*(pneut(x)*pneut(y)) = (x*y)*(pneut(y)*pneut(x)) = x*y*pneut(x) = (x*pneut(x))*y = x*y;$$

$$(pneut(x)*pneut(y))*(x*y) = (pneut(y)*pneut(x))*(x*y) = pneut(y)*x*y = x*(pneut(y)*y) = x*y.$$

On the other hand, let $(x, pneut(x), anti(x))$ and $(y, pneut(y), anti(y))$ are hyper-neutrosophic-triplets, then

$$\begin{aligned} & xx*y*(anti(y)*anti(x)) \\ &= x*(y*anti(y))*anti(x) \\ &= x*pneut(y)*anti(x) \end{aligned}$$

$$\begin{aligned}
&= p_{neut}(y) * x * anti(x) \\
&= p_{neut}(y) * p_{neut}(x) \\
&= p_{neut}(x) * p_{neut}(y);
\end{aligned}$$

$$(anti(y) * anti(x)) * (x * y) = anti(y) * (anti(x) * x) * y = anti(y) * p_{neut}(x) * y = p_{neut}(x) * anti(y) * y = p_{neut}(x) * p_{neut}(y).$$

Thus, $p_{neut}(x) * p_{neut}(y)$ is a pure neutral element of $x * y$ by Definition 7 and Definition 10. Applying Proposition 8 we get that $p_{neut}(x * y) = p_{neut}(x) * p_{neut}(y)$.

Moreover, assume $p_{neut}(x) = z_1 * x = x * z_1$, $p_{neut}(y) = z_2 * y = y * z_2$. Then, by weak commutativity (Definition 11) we have

$$(z_2 * z_1) * (x * y) = z_2 * (z_1 * x) * y = z_2 * p_{neut}(x) * y = p_{neut}(x) * (z_2 * y) = p_{neut}(x) * p_{neut}(y) = p_{neut}(x * y);$$

$$(x * y) * (z_2 * z_1) = x * (y * z_2) * z_1 = x * p_{neut}(y) * z_1 = (x * z_1) * p_{neut}(y) = p_{neut}(x) * p_{neut}(y) = p_{neut}(x * y).$$

Therefore, the proof is completed. \square

Theorem 3. Let $(H, *)$ be a WCP-NET-semihypergroup and H satisfies

$$(\forall x, y \in H, p_{neut}(x) = p_{neut}(y) \Rightarrow |x * y| = 1). \quad (C1)$$

Define a binary relation \approx on H as following:

$$\forall x, y \in H, x \approx y \text{ if and only if } p_{neut}(x) = p_{neut}(y).$$

Then

- (1) The binary relation \approx is a equivalent relation on H ;
- (2) For any $x \in H$, $[x]_{\approx}$ is a sub-NET-semihypergroup of H , where $[x]_{\approx}$ is the equivalent class of x based on equivalent relation \approx ;
- (3) For any $x \in H$, $[x]_{\approx}$ is a regular hypergroup.

Proof. (1) From the definition of \approx , by Proposition 8 and Proposition 9, we know that the binary relation \approx is a equivalent relation.

(2) Suppose $a, b \in [x]_{\approx}$. By the definition of \approx , $p_{neut}(a) = p_{neut}(b) = p_{neut}(x)$. Using Proposition 8 and Proposition 9, we have

$$p_{neut}(a * b) = p_{neut}(a) * p_{neut}(b) = p_{neut}(x) * p_{neut}(x) = p_{neut}(x).$$

It follows that $[x]_{\approx}$ is closed on the hyper operation $*$.

And, applying Proposition 8, we have $p_{neut}(x) * p_{neut}(x) = p_{neut}(x)$. From this and using Proposition 8, we get that $p_{neut}(p_{neut}(x)) = p_{neut}(x)$. It follows that $p_{neut}(a) \in [x]_{\approx}$ for any $a \in [x]_{\approx}$. Moreover, assume that $a \in [x]_{\approx}$, by the definition of WCP-NET-semihypergroup, there exists $r \in H$ such that $p_{neut}(a) = r * a = a * r$. Thus (by Proposition 9)

$$\begin{aligned}
p_{neut}(a) &= (r * p_{neut}(a)) * a = a * (r * p_{neut}(a)) \\
&\Rightarrow r * p_{neut}(a) \in \{anti(a)\}. \\
&p_{neut}(r * p_{neut}(a)) \\
&= p_{neut}(r) * p_{neut}(p_{neut}(a)) \\
&= p_{neut}(r) * p_{neut}(a) \\
&= p_{neut}(r * a) \\
&= p_{neut}(p_{neut}(a)) \\
&= p_{neut}(a).
\end{aligned}$$

That is, $pneut(r^*pneut(a)) = pneut(a) = pneut(x)$. This means that $r^*pneut(a) \in [x]_{\approx}$. Combining this and $r^*pneut(a) \in \{anti(a)\}$, we know that there exists $anti(a)$ which is in $[x]_{\approx}$. This means that $[x]_{\approx}$ is a sub-NET-semihypergroup of H .

(3) Assume $x \in H$, from (2) we know that $[x]_{\approx}$ is a sub-NET-semihypergroup of H . By the definition of \approx , for any $a \in [x]_{\approx}$, $pneut(a) = pneut(x)$. From the proof of (2), there exists $anti(a) \in \{anti(a)\}$ and $anti(a) \in [x]_{\approx}$. Then, $[x]_{\approx} \subseteq a^*[x]_{\approx}^*a$. Obviously, $a^*[x]_{\approx}^*a \subseteq [x]_{\approx}$. Thus, $a^*[x]_{\approx}^*a = [x]_{\approx}$.

On the other hand, $pneut(x)$ is a (local) identity in $[x]_{\approx}$. Therefore, by Definition 3, we get that $[x]_{\approx}$ is a regular hypergroup. □

Example 13. Denote $H = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11\}$, define hyperoperations $*$ on H as shown in Table 11. We can verify that $(H, *)$ is WCP-NET-semihypergroup, and not commutative.

Table 11. The hyperoperation $*$ on H .

$*$	1	2	3	4	5	6	7	8	9	10	11
1	1	{1,2,3}	{1,2,3}	1	1	1	1	1	1	1	1
2	{1,2,3}	3	2	3	2	{1,2,3}	{1,2,3}	{1,2,3}	{1,2,3}	{1,2,3}	{1,2,3}
3	{1,2,3}	2	3	2	3	{1,2,3}	{1,2,3}	{1,2,3}	{1,2,3}	{1,2,3}	{1,2,3}
4	1	3	2	5	4	{6,7,8, 9,10,11}	{6,7,8, 9,10,11}	{6,7,8, 9,10,11}	{6,7,8, 9,10,11}	{6,7,8, 9,10,11}	{6,7,8, 9,10,11}
5	1	2	3	4	5	{6,7,8, 9,10,11}	{6,7,8, 9,10,11}	{6,7,8, 9,10,11}	{6,7,8, 9,10,11}	{6,7,8, 9,10,11}	{6,7,8, 9,10,11}
6	1	{1,2,3}	{1,2,3}	{6,7,8, 9,10,11}	{6,7,8, 9,10,11}	6	7	8	9	10	11
7	1	{1,2,3}	{1,2,3}	{6,7,8, 9,10,11}	{6,7,8, 9,10,11}	7	6	11	10	9	8
8	1	{1,2,3}	{1,2,3}	{6,7,8, 9,10,11}	{6,7,8, 9,10,11}	8	10	6	11	7	9
9	1	{1,2,3}	{1,2,3}	{6,7,8, 9,10,11}	{6,7,8, 9,10,11}	9	11	10	6	8	7
10	1	{1,2,3}	{1,2,3}	{6,7,8, 9,10,11}	{6,7,8, 9,10,11}	10	8	9	7	11	6
11	1	{1,2,3}	{1,2,3}	{6,7,8, 9,10,11}	{6,7,8, 9,10,11}	11	9	7	8	6	10

Moreover,

$$\begin{aligned}
 H_1 &= \{1\} = [1]_{\approx}; \\
 H_2 &= \{2, 3\} = [2]_{\approx} = [3]_{\approx}; \\
 H_3 &= \{4, 5\} = [4]_{\approx} = [5]_{\approx}; \\
 H_4 &= \{6, 7, 8, 9, 10, 11\} = [6]_{\approx} = [7]_{\approx} = [8]_{\approx} = [9]_{\approx} = [10]_{\approx} = [11]_{\approx};
 \end{aligned}$$

and $H = H_1 \cup H_2 \cup H_3 \cup H_4$, where, $H_i (i=1, 2, 3, 4)$ are regular hypergroups.

5. Conclusions

In this paper, we propose some new notions of neutrosophic extended triplet semihypergroup (NET-semihypergroup), neutrosophic extended triplet hypergroup (NET-hypergroup), pure NET-semihypergroup and weak commutative NET-semihypergroup, investigate some basic properties and the relationships among them (see Figure 6), study their close connections with regular hypergroups and regular semihypergroups. Particularly, we prove two structure theorems of commutative pure NET-semihypergroup (CP-NET-semihypergroup) and weak commutative pure NET-semihypergroup (WCP-NET-semihypergroup) under the condition (C1) (see Theorem 2 and Theorem 3). From these results, we know that NET-semihypergroup is a hyperalgebraic structure independent of hypergroup, and NET-semihypergroup is also a generalization of group concept in hyperstructures. The research results in this paper show that NET-semihypergroups and NET-hypergroups have important theoretical research value, which greatly enriches the traditional theory of hyperalgebraic structures.




Figure 6. The relationships among some kinds of NET-semihypergroups.

In the future, we will investigate the combinations of NET-semihypergroups and related algebraic systems ([22–24]).

Author Contributions: F.S. and X.Z. proposed the idea of the research, X.Z. and Y.M. wrote the paper.

Funding: This research was funded by National Natural Science Foundation of China (Grant Nos. 61573240 and 61976130).

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Corsini, P.; Leoreanu, V. *Applications of Hyperstructure Theory*; Springer: Berlin/Heidelberg, Germany, 2003.
2. Wall, H.S. Hypergroups. *Am. J. Math.* **1937**, *59*, 77–98.
3. Freni, D. A new characterization of the derived hypergroup via strongly regular equivalences. *Commun. Algebra* **2002**, *30*, 3977–3989.
4. Davvaz, B. *Semihypergroup Theory*; Elsevier: Amsterdam, The Netherlands, 2016.
5. Jafarabadi, H.M.; Sarmin, N.H.; Molaei, M.R. Simple semihypergroups. *Aust. J. Basic Appl. Sci.* **2011**, *5*, 51–55.
6. Salvo, M.D.; Freni, D.; Faro, G.L. Fully simple semihypergroups. *J. Algebra* **2014**, *399*, 358–377.
7. Jafarabadi, H.M.; Sarmin, N.H.; Molaei, M.R. Completely simple and regular semihypergroups. *Bull. Malays. Math. Sci. Soc.* **2012**, *35*, 335–343.
8. Howie, J.M. *Fundamentals of Semigroup Theory*; Oxford University Press: Oxford, UK, 1995.
9. Smarandache, F. Neutrosophic set—A generalization of the intuitionistic fuzzy sets. *Int. J. Pure Appl. Math.* **2005**, *3*, 287–297.
10. Zhang, X.H.; Bo, C.X.; Smarandache, F.; Dai, J.H. New inclusion relation of neutrosophic sets with applications and related lattice structure. *Int. J. Mach. Learn. Cybern.* **2018**, *9*, 1753–1783.
11. Zhang, X.H.; Bo, C.X.; Smarandache, F.; Park, C. New operations of totally dependent-neutrosophic sets and totally dependent-neutrosophic soft sets. *Symmetry* **2018**, *10*, 187, doi:10.3390/sym10060187.
12. Zhang, X.H.; Mao, X.Y.; Wu, Y.T.; Zhai, X.H. Neutrosophic filters in pseudo-BCI algebras. *Int. J. Uncertain. Quantif.* **2018**, *8*, 511–526.
13. Smarandache, F.; Ali, M. Neutrosophic triplet group. *Neural Comput. Appl.* **2018**, *29*, 595–601.
14. Smarandache, F. *Neutrosophic Perspectives: Triplets, Duplets, Multisets, Hybrid Operators, Modal Logic, Hedge Algebras. And Applications*; Pons Publishing House: Brussels, Belgium, 2017.
15. Zhang, X.H.; Hu, Q.Q.; Smarandache, F.; An, X.G. On neutrosophic triplet groups: Basic properties, NT-subgroups, and some notes. *Symmetry* **2018**, *10*, 289, doi:10.3390/sym10070289.

16. Jaíyéolá, T.G.; Smarandache, F. Some results on neutrosophic triplet group and their applications. *Symmetry* **2018**, *10*, 202.
17. Zhang, X.H.; Wu, X.Y.; Mao, X.Y.; Smarandache, F.; Park, C. On neutrosophic extended triplet groups (Loops) and Abel-Grassmann's Groupoids (AG-Groupoids). *J. Intell. Fuzzy Syst.* **2019**, doi: 10.3233/JIFS-181742.
18. Wu, X.Y.; Zhang, X.H. The decomposition theorems of AG-neutrosophic extended triplet loops and strong AG-(l, l)-loops. *Mathematics* **2019**, *7*, 268, doi:10.3390/math7030268.
19. Ma, Y.C.; Zhang, X.H.; Yang, X.F.; Zhou, X. Generalized neutrosophic extended triplet group. *Symmetry* **2019**, *11*, 327, doi:10.3390/sym11030327.
20. Zhang, X.H.; Wang, X.J.; Smarandache, F.; Jaíyéolá, T.G.; Lian, T.Y. Singular neutrosophic extended triplet groups and generalized groups. *Cogn. Syst. Res.* **2019**, *57*, 32–40.
21. Gulistan, M.; Nawaz, S.; Hassan, N. Neutrosophic triplet non-associative semihypergroups with application. *Symmetry* **2018**, *10*, 613, doi:10.3390/sym10110613.
22. Zhang, X.H.; Borzooei, R.A.; Jun, Y.B. Q-filters of quantum B-algebras and basic implication algebras. *Symmetry* **2018**, *10*, 573, doi:10.3390/sym10110573.
23. Omid, S.; Davvaz, B.; Zhan, J.M. An investigation on ordered algebraic hyperstructures. *Acta Math. Sin. Engl. Ser.* **2017**, *33*, 1107–1124.
24. Zhang, X.H.; Mao, X.Y.; Smarandache, F.; Park, C. On homomorphism theorem for perfect neutrosophic extended triplet groups. *Information* **2018**, *9*, 237, doi:10.3390/info9090237.



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).