

## Global Equitable Domination Number of Some Wheel Related Graphs

S.K.Vaidya and R.M.Pandit

(Department of Mathematics, Saurashtra University, RAJKOT-360 005 (Gujarat) India)

E-mail: samirkvaidya@yahoo.co.in

**Abstract:** A dominating set is called a global dominating set if it is a dominating set of a graph  $G$  and its complement  $\overline{G}$ . A subset  $D$  of  $V(G)$  is called an equitable dominating set if for every  $v \in V(G) - D$ , there exists a vertex  $u \in D$  such that  $uv \in E(G)$  and  $|d_G(u) - d_G(v)| \leq 1$ . An equitable dominating set  $D$  of a graph  $G$  is a global equitable dominating set if it is also an equitable dominating set of the complement of  $G$ . The global equitable domination number  $\gamma_g^e(G)$  of  $G$  is the minimum cardinality of a global equitable dominating set of  $G$ . In this paper, we investigate the global equitable domination number of some wheel related graphs.

**Key Words:** Global dominating set, equitable dominating set, Smarandachely equitable dominating set, global equitable dominating set, global equitable domination number.

**AMS(2010):** 05C69, 05C76.

### §1. Introduction

The study of domination in graphs is one of the fastest growing areas within graph theory. An excellent survey on the concept of domination and its related parameters can be found in the book by Haynes *et al.* [4] while some advanced topics on domination are explored in Haynes *et al.* [5]. The concept of domination has interesting applications in the study of social networks which motivated Prof. E. Sampathkumar to introduce the concept of equitable domination in graphs.

Secondly, let  $G$  be a graph of road network linking various locations. It is desirable to maintain the supply to these locations uninterruptedly by using the alternative links even if the original links get disturbed. Then the problem of finding the minimum number of supplying stations needed to accomplish this task is equivalent to find the global domination number. The concept of global domination was introduced by Sampathkumar [9].

Many domination models are introduced by combining two different domination parameters. Independent domination, global domination, equitable domination, connected domination are among worth to mention. Motivated through the concepts of global domination and equitable domination, a new concept of global equitable domination was conceived by Basavanagoud

---

<sup>1</sup>Received December 01, 2015, Accepted August 12, 2016.

and Teli [2] and formalized by Vaidya and Pandit [14]. In the present paper, we obtain the global equitable domination number of some wheel related graphs.

Throughout the paper, a graph  $G = (V(G), E(G))$  we mean a finite and undirected graph without loops and multiple edges. The set  $D \subseteq V(G)$  of vertices in a graph  $G$  is called a dominating set if every vertex  $v \in V(G)$  is either an element of  $D$  or is adjacent to an element of  $D$ . The minimum cardinality of a dominating set of  $G$  is called the domination number of  $G$  which is denoted by  $\gamma(G)$ .

The complement  $\overline{G}$  of  $G$  is the graph with vertex set  $V(G)$  in which two vertices are adjacent in  $\overline{G}$  if they are not adjacent in  $G$ .

For a vertex  $v \in V(G)$ , the open neighborhood of  $v$ , denoted by  $N(v)$ , is  $\{u \in V(G) : uv \in E(G)\}$ . We denote the degree of a vertex  $v$  in  $G$  by  $d_G(v)$ . A vertex of degree one is called a pendant vertex and a vertex which is not the end of any edge is called an isolated vertex. An edge  $e$  of a graph  $G$  is said to be incident with the vertex  $v$  if  $v$  is an end vertex of  $e$ . An edge incident with a pendant vertex is called a pendant edge.

A set  $D \subseteq V(G)$  is called a global dominating set of  $G$  if  $D$  is a dominating set of both  $G$  and  $\overline{G}$ . The global domination number  $\gamma_g(G)$  is the minimum cardinality of a global dominating set in  $G$ . Many researchers have explored this concept. For example, Gangadharappa and Desai [3] have discussed the global domination in graphs of small diameters. Vaidya and Pandit [12, 13] have investigated the global domination number of the larger graphs obtained by some graph operations on a given graph while Kulli and Janakiram [6] have introduced the concept of total global dominating sets.

A subset  $D$  of  $V(G)$  is called an equitable dominating set if for every  $v \in V(G) - D$ , there exists a vertex  $u \in D$  such that  $uv \in E(G)$  and  $|d_G(u) - d_G(v)| \leq 1$ , otherwise, a Smarandachely equitable dominating set, i.e.,  $|d_G(u) - d_G(v)| \geq 2$  for each edge  $uv \in E(G)$  with  $u \in D$  and  $v \in V(G) - D$ . The minimum cardinality of such a dominating set is called the equitable domination number of  $G$  which is denoted by  $\gamma^e(G)$ . Swaminathan and Dharmalingam [11] have studied the equitable domination in graphs and characterized the minimal equitable dominating sets. Sivakumar *et al.* [10] have discussed the connected equitable domination in graphs while Murugan and Emmanuel [7] have identified the inter relationship among domination, equitable domination and independent domination in graphs. Revathi and Harinarayanan [8] have studied the equitable domination in fuzzy graphs while Basavanagoud *et al.* [1] have studied the equitable total domination in graphs.

A vertex  $v \in V(G)$  is equitably adjacent with a vertex  $u \in V(G)$  if  $|d_G(u) - d_G(v)| \leq 1$  and  $uv \in E(G)$ . A vertex  $u \in V(G)$  is called an equitable isolate if  $|d_G(u) - d_G(v)| \geq 2$  for all  $v \in N(u)$ . Analogous to the characteristic of an isolated vertex in a dominating set, an equitable isolate must belong to any equitable dominating set of  $G$ . Clearly, the isolated vertices are the equitable isolates. Hence,  $I_s \subseteq I_e \subseteq D$  for every equitable dominating set  $D$  where  $I_s$  and  $I_e$  denote the sets of all isolated vertices and all equitable isolates of  $G$  respectively.

A subset  $D$  of  $V(G)$  is called a global equitable dominating set of  $G$  if  $D$  is an equitable dominating set of both  $G$  and  $\overline{G}$ . The minimum cardinality of a global equitable dominating set of  $G$  is called the global equitable domination number of  $G$  and it is denoted by  $\gamma_g^e(G)$ .

Since at least two vertices are required to equitably dominate both  $G$  and  $\overline{G}$ , we have

$2 \leq \gamma_g^e(G) \leq n$  for every graph of order  $n > 1$ . Both of these bounds are sharp. In particular, the equality of the lower bound is attained by  $P_n$  ( $2 \leq n \leq 6$ ) and  $K_{r,s}$  ( $|r - s| \leq 1$ ) while the upper bound is achieved by  $K_n$ ,  $K_{1,p}$  and  $K_{r,s}$  ( $|r - s| \geq 2$ ).

The wheel  $W_n$  is defined to be the join  $C_{n-1} + K_1$  where  $n \geq 4$ . The vertex corresponding to  $K_1$  is known as the apex vertex and the vertices corresponding to cycle  $C_{n-1}$  are known as the rim vertices. For any real number  $n$ ,  $\lceil n \rceil$  denotes the smallest integer not less than  $n$  and  $\lfloor n \rfloor$  denotes the greatest integer not greater than  $n$ .

For notations and graph theoretic terminology not defined herein, we refer the readers to West [15] while the terms related to the concept of domination are used in the sense of Haynes *et al.* [4].

## §2. Main Results

**Definition 2.1** *The helm  $H_n$  is the graph obtained from a wheel  $W_n$  by attaching a pendant edge to each of its rim vertices.*

**Proposition 2.2** ([2])

- (i) For the path  $P_n$  ( $n \geq 4$ ),  $\gamma_g^e(P_n) = \lceil \frac{n}{3} \rceil$ ;  
(ii) For the cycle  $C_n$ ,  $\gamma_g^e(C_n) = \begin{cases} 3 & \text{if } n = 3, 5 \\ \lceil \frac{n}{3} \rceil & \text{otherwise.} \end{cases}$

**Theorem 2.3** For the helm,  $\gamma_g^e(H_n) = \begin{cases} 7 & \text{if } n = 4 \\ n + 2 & \text{if } n = 5, 6 \\ \lceil \frac{4n-1}{3} \rceil & \text{otherwise.} \end{cases}$

*Proof* Let  $v_1, v_2, \dots, v_{n-1}$  be the rim vertices of wheel  $W_n$  and let  $c$  denotes the apex vertex of the helm  $H_n$ . Let  $u_1, u_2, \dots, u_{n-1}$  be the pendant vertices of  $H_n$ . Then,  $|V(H_n)| = 2n - 1$  and  $|E(H_n)| = 3(n - 1)$ .

**Case 1.**  $n = 4$

For  $n = 4$ , the pendant vertices of  $H_n$  are equitable isolates in  $H_n$  while the remaining vertices of  $H_n$  are equitable isolates in  $\overline{H_n}$ . Hence, the vertex set of  $H_n$  is the only global equitable dominating set of  $H_n$  implying that  $\gamma_g^e(H_n) = |V(H_n)| = 7$ .

**Case 2.**  $n = 5, 6$

Since the  $n - 1$  pendant vertices of  $H_n$  are equitable isolates in  $H_n$  and the apex vertex  $c$  is an equitable isolate in  $\overline{H_n}$ , it follows that every global equitable dominating set of  $H_n$  must contain these vertices. Now, these vertices equitably dominate all the vertices of  $H_n$  but do not equitably dominate all the vertices of  $\overline{H_n}$ . Moreover, any two adjacent rim vertices of  $W_n$  can equitably dominate the remaining vertices of  $\overline{H_n}$ . Hence, every global equitable dominating set of  $H_n$  must contain at least  $n + 2$  vertices of  $H_n$ . Therefore,  $\gamma_g^e(H_n) = n + 2$ .

**Case 3.**  $n \geq 7$

In this case, the apex vertex is an equitable isolate in  $H_n$  as well as in  $\overline{H_n}$  while the  $n - 1$  pendant vertices of  $H_n$  are equitable isolates in  $H_n$  only. Therefore, these vertices must belong to every global equitable dominating set of  $H_n$ . Now, the remaining vertices induce a cycle  $C_{n-1}$  and by Proposition 2.2,  $\gamma_g^e(C_{n-1}) = \lceil \frac{n-1}{3} \rceil$ . Therefore,  $\gamma_g^e(H_n) = n - 1 + 1 + \lceil \frac{n-1}{3} \rceil = \lceil \frac{4n-1}{3} \rceil$ .

Thus, we have proved that

$$\gamma_g^e(H_n) = \begin{cases} 7 & \text{if } n = 4 \\ n + 2 & \text{if } n = 5, 6 \\ \lceil \frac{4n-1}{3} \rceil & \text{otherwise.} \end{cases}$$

□

**Definition 2.4** The flower graph  $Fl_n$  is the graph obtained from the helm  $H_n$  by joining each pendant vertex to the apex vertex of the helm  $H_n$ .

**Theorem 2.5** For the flower graph,

$$\gamma_g^e(Fl_n) = \begin{cases} n + 3 & \text{if } n = 4, 6 \\ \lceil \frac{4n-1}{3} \rceil & \text{otherwise.} \end{cases}$$

*Proof* Let  $v_1, v_2, \dots, v_{n-1}$  be the rim vertices of wheel  $W_n$  and let  $u_1, u_2, \dots, u_{n-1}$  be the pendant vertices of the helm  $H_n$ . Let  $c$  denotes the apex vertex of  $Fl_n$ . Then  $|V(Fl_n)| = 2n - 1$ . Here,  $d_G(v_i) = 4$ ,  $d_G(u_i) = 2$  for  $1 \leq i \leq n - 1$  and  $d_G(c) = 2(n - 1)$  where  $G = Fl_n$ .

Now, the vertex  $c$  is an equitable isolate in  $G$  as well as in  $\overline{G}$ . Therefore, every global equitable dominating set of  $G$  must contain  $c$ . Moreover, the vertices  $u_1, u_2, \dots, u_{n-1}$  being the equitable isolates in  $G$ , must belong to every global equitable dominating set of  $G$ . Now, the remaining vertices  $v_1, v_2, \dots, v_{n-1}$  in  $G$  induce a cycle  $C_{n-1}$  and by Proposition 2.2,

$$\gamma_g^e(C_n) = \begin{cases} 3 & \text{if } n = 3, 5 \\ \lceil \frac{n}{3} \rceil & \text{otherwise.} \end{cases}$$

Hence,

$$\begin{aligned} \gamma_g^e(Fl_n) &= \gamma_g^e(C_{n-1}) + (n - 1) + 1 \\ &= \gamma_g^e(C_{n-1}) + n \\ &= \begin{cases} n + 3 & \text{if } n = 4, 6 \\ \lceil \frac{n-1}{3} \rceil + n & \text{otherwise.} \end{cases} \end{aligned}$$

Thus,

$$\gamma_g^e(Fl_n) = \begin{cases} n + 3 & \text{if } n = 4, 6 \\ \lceil \frac{4n-1}{3} \rceil & \text{otherwise.} \end{cases}$$

□

**Definition 2.6** The sunflower graph  $Sf_n$  is the resultant graph obtained from the flower graph by attaching  $(n - 1)$  pendant edges to the apex vertex of wheel  $W_n$ .

**Theorem 2.7** For the sunflower graph,  $\gamma_g^e(Sf_n) = 3n - 2$ .

*Proof* Let  $c$  denotes the apex vertex of wheel  $W_n$  and let  $v_1, v_2, \dots, v_{n-1}$  be the rim vertices of  $W_n$ . Let  $u_1, u_2, \dots, u_{n-1}$  be the vertices of degree 2 in  $Sf_n$  and let  $x_1, x_2, \dots, x_{n-1}$  be the pendant vertices of  $Sf_n$ . Then,  $|V(Sf_n)| = 3n - 2$ .

Now,  $c$  is the equitable isolate in both  $Sf_n$  and  $\overline{Sf_n}$ . Moreover, the vertices  $u_1, u_2, \dots, u_{n-1}, x_1, x_2, \dots, x_{n-1}$  are equitable isolates in  $Sf_n$  while the vertices  $v_1, v_2, \dots, v_{n-1}$  are equitable isolates in  $\overline{Sf_n}$ . Since an equitable isolate must belong to every equitable dominating set, it follows that the vertex set  $V(Sf_n)$  is the only global equitable dominating set of  $Sf_n$ . Therefore,  $\gamma_g^e(Sf_n) = |V(Sf_n)| = 3n - 2$ .  $\square$

**Definition 2.8** The closed helm  $CH_n$  is the graph obtained from a helm by joining each pendant vertex to form a cycle.

**Theorem 2.9** For the closed helm  $CH_n$  ( $n > 5$ ),

$$\gamma_g^e(CH_n) = \begin{cases} \lfloor \frac{n+2}{2} \rfloor & \text{if } n \equiv 1 \pmod{4} \\ \lceil \frac{n+2}{2} \rceil & \text{if } n \equiv 0, 2 \text{ or } 3 \pmod{4}. \end{cases}$$

*Proof* Let  $v_1, v_2, \dots, v_{n-1}$  be the vertices of degree 4 and let  $u_1, u_2, \dots, u_{n-1}$  be the vertices of degree 3 in  $G = CH_n$ . Let  $c$  denotes the apex vertex of  $CH_n$ . Then the closed helm  $CH_n$  has  $2n - 1$  vertices. The vertex  $c$  is an equitable isolate in  $CH_n$  and the remaining vertices which are adjacent in  $CH_n$ , are also equitably adjacent in  $CH_n$ . The vertex  $c$  being an equitable isolate, must belong to every global equitable dominating set of  $CH_n$ . Hence, we construct a vertex set  $D \subset V(CH_n)$  as follows:

$$D = \{c, v_{4i+1}, u_{4j+3}\},$$

where  $0 \leq i \leq \lfloor \frac{n-2}{4} \rfloor$  and  $0 \leq j \leq \lfloor \frac{n}{4} \rfloor$  with

$$|D| = \begin{cases} \lfloor \frac{n+2}{2} \rfloor & \text{if } n \equiv 1 \pmod{4} \\ \lceil \frac{n+2}{2} \rceil & \text{if } n \equiv 0, 2 \text{ or } 3 \pmod{4}. \end{cases}$$

Now,  $d_G(v_{4i+1}) = 4$ ,  $d_G(u_{4j+3}) = 3$  and  $d_{\overline{G}}(u_{4j+3}) - d_{\overline{G}}(v_{4j+1}) = 1$ . Then for every  $v \in V(G) - D$ , there exists a vertex  $u \in D$  such that  $uv \in E(G)$  and  $|d_G(u) - d_G(v)| \leq 1$ . Moreover, for every  $v' \in V(G) - D$  there exists a vertex  $u' \in D$  such that  $u'v' \in E(\overline{G})$  and  $|d_{\overline{G}}(u') - d_{\overline{G}}(v')| \leq 1$ . Hence, the set  $D$  is an equitable dominating set of  $G$  as well as of  $\overline{G}$ . Therefore,  $D$  is a global equitable dominating set of  $G$ . Moreover, from the adjacency nature of the vertices of  $G$ , one can observe that the set  $D$  is of minimum cardinality.

Thus, the set  $D$  is a global equitable dominating set of  $G = CH_n$  ( $n > 5$ ) with minimum

cardinality implying that

$$\gamma_g^e(CH_n) = \begin{cases} \lfloor \frac{n+2}{2} \rfloor & \text{if } n \equiv 1 \pmod{4} \\ \lceil \frac{n+2}{2} \rceil & \text{if } n \equiv 0, 2 \text{ or } 3 \pmod{4}. \end{cases}$$

□

**Remark 2.10** For  $n = 4, 5$ , at least two vertices are required to equitably dominate all the vertices of  $CH_n$  as well as of  $\overline{CH_n}$ . Therefore,  $\gamma_g^e(CH_n) = 2$  for  $n = 4, 5$ .

**Definition 2.11** A web graph is the graph obtained by joining the pendant vertices of a helm to form a cycle and then adding a single pendant edge to each vertex of this outer cycle. We denote the web graph by  $Wb_n$ .

**Theorem 2.12** For the web graph  $Wb_n$  ( $n > 6$ ),

$$\gamma_g^e(Wb_n) = \begin{cases} \lfloor \frac{3n}{2} \rfloor & \text{if } n \equiv 1 \pmod{4} \\ \lceil \frac{3n}{2} \rceil & \text{if } n \equiv 0, 2 \text{ or } 3 \pmod{4}. \end{cases}$$

*Proof* Let  $c$  denotes the apex vertex of web graph  $Wb_n$ . Let  $v_1, v_2, \dots, v_{n-1}$  and  $u_1, u_2, \dots, u_{n-1}$  be the vertices of inner cycle and outer cycle of  $Wb_n$  respectively. Let  $x_1, x_2, \dots, x_{n-1}$  denote the pendant vertices of  $Wb_n$ .

Since the apex vertex  $c$  and the  $n - 1$  pendant vertices are the equitable isolates in  $Wb_n$  as well as in  $\overline{Wb_n}$  for  $n > 6$ , it follows that these vertices must belong to every global equitable dominating set of  $Wb_n$ . Moreover, the vertices except the pendant vertices induce the closed helm  $CH_n$  and by Theorem 2.9, we have

$$\gamma_g^e(CH_n) = \begin{cases} \lfloor \frac{n+2}{2} \rfloor & \text{if } n \equiv 1 \pmod{4} \\ \lceil \frac{n+2}{2} \rceil & \text{if } n \equiv 0, 2 \text{ or } 3 \pmod{4}. \end{cases}$$

Hence,

$$\begin{aligned} \gamma_g^e(Wb_n) &= \gamma_g^e(CH_n) + (n - 1) \\ &= \begin{cases} \lfloor \frac{n+2}{2} \rfloor + (n - 1) & \text{if } n \equiv 1 \pmod{4} \\ \lceil \frac{n+2}{2} \rceil + (n - 1) & \text{if } n \equiv 0, 2 \text{ or } 3 \pmod{4}. \end{cases} \end{aligned}$$

Thus, for  $n > 6$ ,

$$\gamma_g^e(Wb_n) = \begin{cases} \lfloor \frac{3n}{2} \rfloor & \text{if } n \equiv 1 \pmod{4} \\ \lceil \frac{3n}{2} \rceil & \text{if } n \equiv 0, 2 \text{ or } 3 \pmod{4}. \end{cases}$$

□

**Remark 2.13** (i) For  $n = 4, 5$ , the  $n - 1$  pendant vertices are the equitable isolates in  $Wb_n$

as well as in  $\overline{Wb_n}$  and the remaining vertices induce the closed helm  $CH_n$ . Thus,  $\gamma_g^e(Wb_n) = \gamma_g^e(CH_n) + (n-1)$  implying that  $\gamma_g^e(Wb_4) = 5$  and  $\gamma_g^e(Wb_5) = 6$ .

(ii) For  $n = 6$ , the apex vertex is not an equitable isolate in  $Wb_6$  as well as in  $\overline{Wb_6}$  while the pendant vertices are the equitable isolates in both  $Wb_6$  and  $\overline{Wb_6}$ . Hence,  $\gamma_g^e(Wb_6) = 8$ .

**Definition 2.14** A gear graph  $G_n$  is obtained from the wheel  $W_n$  by adding a vertex between every pair of adjacent vertices of the  $(n-1)$ -cycle of  $W_n$ .

**Theorem 2.15** For the gear graph,

$$\gamma_g^e(G_n) = \begin{cases} \lceil \frac{n}{2} \rceil & \text{if } n = 4, 5 \\ \lceil \frac{2n+1}{3} \rceil & \text{otherwise.} \end{cases}$$

*Proof* Let  $c$  denotes the apex vertex of wheel  $W_n$  and let  $v_1, v_2, \dots, v_{n-1}$  be the rim vertices of  $W_n$ . To obtain the gear graph  $G_n$ , subdivide each rim edge of wheel by the vertices  $u_1, u_2, \dots, u_{n-1}$  where each  $u_i$  is added between  $v_i$  and  $v_{i+1}$  for  $i = 1, 2, \dots, n-2$  and  $u_{n-1}$  is added between  $v_1$  and  $v_{n-1}$ . Then  $|V(G_n)| = 2n-1$  and  $|E(G_n)| = 3(n-1)$ . The graph  $G_n$  contains the outer cycle  $C_{2(n-1)}$ .

For  $n = 4, 5$ , the sets  $D = \{v_1, u_3\}$  and  $D = \{c, v_1, v_3\}$  are clearly the global equitable dominating sets of  $G_4$  and  $G_5$  respectively with minimum cardinality. Therefore,  $\gamma_g^e(G_n) = \lceil \frac{n}{2} \rceil$  for  $n = 4, 5$ .

For  $n > 5$ , since the vertex  $c$  is the equitable isolate in  $G_n$  as well as in  $\overline{G_n}$ , it must belong to every global equitable dominating set of  $G_n$ . Moreover, the vertices other than  $c$  induce a cycle  $C_{2(n-1)}$ . Furthermore,  $V(G_n) = V(C_{2(n-1)}) \cup \{c\}$  and by Proposition 2.2,  $\gamma_g^e(C_n) = \lceil \frac{n}{3} \rceil$  for  $n > 5$ . This implies that  $\gamma_g^e(G_n) = \gamma_g^e(C_{2(n-1)}) + 1 = \lceil \frac{2(n-1)}{3} \rceil + 1 = \lceil \frac{2n+1}{3} \rceil$ . Hence, we have proved that

$$\gamma_g^e(G_n) = \begin{cases} \lceil \frac{n}{2} \rceil & \text{if } n = 4, 6 \\ \lceil \frac{2n+1}{3} \rceil & \text{otherwise.} \end{cases}$$

□

**Definition 2.16** The splitting graph  $S'(G)$  of a graph  $G$  is obtained by adding a new vertex  $v'$  corresponding to each vertex  $v$  of  $G$  such that  $N(v) = N(v')$ .

**Theorem 2.17** For the splitting graph of wheel  $W_n$  ( $n > 7$ ),

$$\gamma_g^e(S'(W_n)) = \begin{cases} \lceil \frac{4n+3}{3} \rceil & \text{if } n \equiv 0 \text{ or } 2 \pmod{3} \\ \lfloor \frac{4n+3}{3} \rfloor & \text{if } n \equiv 1 \pmod{3}. \end{cases}$$

*Proof* Let  $v_1, v_2, \dots, v_{n-1}$  be the rim vertices of wheel  $W_n$  and let  $c$  denotes the apex vertex of  $W_n$ . Let  $c', v'_1, v'_2, \dots, v'_{n-1}$  be the added vertices corresponding to the vertices  $c, v_1, v_2, \dots, v_{n-1}$  of  $W_n$  to obtain  $G = S'(W_n)$ . Then  $|V(G)| = 2n$ .

For  $n = 8$ , the vertices  $c$  and  $c'$  are equitable isolates in  $\overline{G}$  and  $c, v'_1, v'_2, \dots, v'_7$  are equitable isolates in  $G$ . For  $n > 8$ , the vertices  $c$  and  $c'$  are equitable isolates in both  $G$  and  $\overline{G}$  while

the vertices  $v'_1, v'_2, \dots, v'_{n-1}$  are equitable isolates in  $G$ . Since an equitable isolate must belong to every equitable dominating set of  $G$ , the vertices  $c, c', v'_1, v'_2, \dots, v'_{n-1}$  being equitable isolates, must belong to every global equitable dominating set of  $G$ . Now, the remaining vertices  $v_1, v_2, \dots, v_{n-1}$  of  $G$  induce a cycle  $C_{n-1}$  and by Proposition 2.2,  $\gamma_g^e(C_n) = \lceil \frac{n}{3} \rceil$  for  $n > 5$ . This implies that  $\gamma_g^e(G) = \gamma_g^e(C_{n-1}) + n + 1 = \lceil \frac{n-1}{3} \rceil + n + 1$ . Thus, for  $n > 7$ ,

$$\gamma_g^e(S'(W_n)) = \begin{cases} \lceil \frac{4n+3}{3} \rceil & \text{if } n \equiv 0 \text{ or } 2 \pmod{3} \\ \lfloor \frac{4n+3}{3} \rfloor & \text{if } n \equiv 1 \pmod{3}. \end{cases}$$

□

**Remark 2.18** For  $4 \leq n \leq 7$ , the apex vertex and all the duplicated vertices are the equitable isolates either in  $S'(W_n)$  or in  $\overline{S'(W_n)}$  and by Proposition 2.2,

$$\gamma_g^e(C_n) = \begin{cases} 3 & \text{if } n = 3, 5 \\ \lceil \frac{n}{3} \rceil & \text{otherwise.} \end{cases}$$

Hence,

$$\gamma_g^e(S'(W_n)) = \begin{cases} 8 & \text{if } n = 4, 5 \\ 10 & \text{if } n = 6, 7. \end{cases}$$

### §3. Concluding Remarks

The concept of global equitable domination is a variant of global domination and equitable domination. We obtain the exact values of global equitable domination number of the helm  $H_n$ , the flower graph  $Fl_n$ , the sunflower graph  $Sf_n$ , the closed helm  $CH_n$ , the web graph  $Wb_n$ , the gear graph  $G_n$  and the splitting graph of wheel  $S'(W_n)$ .

### Acknowledgement

The authors are highly thankful to the anonymous referees for their kind comments and fruitful suggestions on the first draft of this paper.

### References

- [1] B.Basavanagoud, V. R.Kulli and V. V.Teli, Equitable total domination in graphs, *Journal of Computer and Mathematical Sciences*, Vol.5(2)(2014), 235-241.
- [2] B.Basavanagoud and V. V.Teli, Equitable global domination in graphs, *International Journal of Mathematical Archive*, Vol.6(3)(2015), 122-125.
- [3] D. B.Gangadharappa and A. R.Desai, On the dominating of a graph and its complement, *Journal of Mathematics and Computer Science*, Vol.2(2)(2011), 222-233.
- [4] T. W.Haynes, S. T.Hedetniemi and P. J.Slater, *Fundamentals of Domination in Graphs*,

- Monographs and Textbooks in Pure and Applied Mathematics, Marcel Dekker, New York, 1998.
- [5] T. W. Haynes, S. T. Hedetniemi and P. J. Slater, *Domination in Graphs - Advanced Topics*, Monographs and Textbooks in Pure and Applied Mathematics, Marcel Dekker, New York, 1998.
  - [6] V. R. Kulli and B. Janakiram, The total global domination number of a graph, *Indian Journal of Pure and Applied Mathematics*, Vol.27(6)(1996), 537-542.
  - [7] A. Nellai Murugan and G. Victor Emmanuel, Degree equitable domination number and independent domination number of a graph, *International Journal of Innovative Research in Science, Engineering and Technology*, Vol.2(11)(2013), 6419-6423.
  - [8] S. Revathi and C. V. R. Harinarayanan, Equitable domination in fuzzy graphs, *S. Revathi International Journal of Engineering Research and Applications*, Vol.4(2014), 80-83.
  - [9] E. Sampathkumar, The global domination number of a graph, *Journal of Mathematical and Physical Sciences*, Vol.23(5)(1989), 377-385.
  - [10] S. Sivakumar, N. D. Soner and A. Alwardi, Connected equitable domination in graphs, *Pure Mathematical Sciences*, Vol.1(3)(2012), 123-130.
  - [11] V. Swaminathan and K. Dharmalingam, Degree equitable domination on graphs, *Kragujevac Journal of Mathematics*, Vol.35(1)(2011), 191-197.
  - [12] S. K. Vaidya and R. M. Pandit, Some new perspectives on global domination in graphs, *ISRN Combinatorics*, Vol. 2013, Article ID 201654, 4 pages, 2013.
  - [13] S. K. Vaidya and R. M. Pandit, Some results on global dominating sets, *Proyecciones Journal of Mathematics*, Vol.32(3)(2013), 235-244.
  - [14] S. K. Vaidya and R. M. Pandit, The global equitable domination in graphs, (communicated).
  - [15] D. B. West, *Introduction to Graph Theory*, Prentice-Hall of India, New Delhi, 2003.