



Neutrosophic Overlap Function and Its Derived Neutrosophic Residual Implication

Xingna Zhang¹, Qingqing Hu² and Xiaohong Zhang^{3,*}

¹ School of Mathematics and Data Science, Shaanxi University of Science & Technology, Xi'an 710021, China; 221711032@sust.edu.cn

² School of Electrical and Control Engineering, Shaanxi University of Science & Technology, Xi'an 710021, China; bs1906001@sust.edu.cn

³ School of Mathematics and Data Science, Shaanxi University of Science & Technology, Xi'an 710021, China; zhangxiaohong@sust.edu.cn

* Correspondence: zhangxiaohong@sust.edu.cn; Tel.: 029-86168320

Abstract: A new concept of neutrosophic overlap function is given, furthermore a neutrosophic residual implication derived from it is also introduced. Firstly, we give new concept of neutrosophic overlap function and some classical examples which are introduced on the lattice. Secondly, the concept of representable neutrosophic overlap function and its pertinent examples are given, meanwhile the general method of constructing representable neutrosophic overlap function by using intuitionistic overlap function is given. Finally, neutrosophic residual implication induced by neutrosophic overlap function and its basic properties are studied.

Keywords: neutrosophic overlap function; lattice; representable neutrosophic overlap function; neutrosophic residual implication

1. Introduction

In 1998, smarandache added an independent membership degree of uncertainty to intuitionistic fuzzy set (IFS) [1], thus putting forward the neutrosophic set (NS) initially. NS is such a robust formal frame extending those concepts of typical set, fuzzy set, IFS and interval-valued IFS from philosophical viewpoint. Because IFS and interval-valued IFS that can solely address incomplete information, but can't address uncertainty and the lack of consistent information which exists in reality. Hence, NS is introduced. Its uncertainty can be explicitly quantified, and its true affiliation, uncertain affiliation and false affiliation are expressed independent. However, its application is hard to solve the actual problems, some scholars have brought forward that notion of single-valued NS [2], as one specific case of NS. And those relevant contents of using single-valued NS to address decision-making issues are as follows [3-8].

Since the triangular norm has a broad range of applications in solving pragmatic issues, it is also important to study the wide range of forms of the triangular norm in applications. The overlap function is a generalization of triangular norm that fulfils continuity [9]. Bustince et al. gave accurate definition of overlap(grouping) function in [10,11]. Over the past period of time, overlap function and grouping function evolved rapidly in theory and practice. See the following literature [12-16] for the rich achievements in the field of theoretical research about overlap function and grouping function. In decision problems, image processing and other fields of wide application see the following literature [17-20]. In an effort to better handle inconclusive information, some scholars extend the overlap function [12, 21] into the IFS, while introducing the method at [22].

Fuzzy implication and fuzzy residual implication play an integral part in traditional fuzzy logic. Fuzzy implication [23] generalizes classical implication into fuzzy logic via the consideration of truth values varying in $[0, 1]$ as opposed to $\{0, 1\}$. Fuzzy implication is one of the important components of fuzzy logic and acts as a very crucial part in some fields, such as image processing, fuzzy control, data mining sees the following literature [24-26], etc. Based on the wide application of fuzzy implication, it is necessary to research it from the theoretical viewpoint [19]. There are a few various models of fuzzy implication for example the R-implication induced by triangular norm [27], (S-N)-implication induced by triangular conorm and fuzzy negation [28], etc. Because overlap function is closely related to triangular norm, in view of the research of neutrosophic triangular norm on neutrosophic fuzzy residual implication, and referring to the research of neutrosophic triangular norm derived residual implication in Hu and Zhang [18], it is natural to consider the neutrosophic residual implication (NRI) induced by neutrosophic overlap function.

The second section mainly introduces the basic knowledge that needs to be used, such as overlap function, grouping function and NS etc. And in the third section, the new concept and related examples of neutrosophic overlap function are given. In addition, the notions and relevant examples of representable neutrosophic overlap function and non-representable neutrosophic overlap function are presented, respectively. Furthermore, the new concept of neutrosophic negation and De Morgan neutrosophic triple which can express the dual relationship between neutrosophic overlap function and neutrosophic grouping function is introduced. The general method of constructing representable neutrosophic overlap functions by intuitionistic overlap functions is given. The fourth section focuses on NRI induced by neutrosophic overlap function, and concludes that every NRI induced by neutrosophic overlap function must be a neutrosophic implication. The final section summarizes the research content.

2. Preliminaries

Definition 2.1 ([29]) O is referred to as an overlap function, if the binary map $O: [0, 1] \times [0, 1] \rightarrow [0, 1]$ fulfils prerequisites below, $\forall s, t, v \in [0, 1]$:

- (a) O fulfils exchangeability;
- (b) $O(s, t) = 0$ when and only when $st = 0$;
- (c) $O(s, t) = 1$ when and only when $st = 1$;
- (d) $O(s, t) \leq O(s, v)$ if $t \leq v$;
- (e) O fulfils continuity.

Example 2.1 The bivariate functions below are overlap functions, $\forall s, t \in [0, 1]$:

- (a) $O_{\text{mM}}(s, t) = \min(s, t) \max(s^2, t^2)$;
- (b) $O_p(s, t) = s^p t^p$, for $p > 0$ and $p \neq 1$;
- (c) $O_{\text{DB}}(s, t) = \begin{cases} \frac{2st}{s+t}, & \text{if } s+t \neq 0, \\ 0, & \text{if } s+t = 0. \end{cases}$;
- (d) $O_{\text{min}}(s, t) = \min\{\sqrt{s}, \sqrt{t}\}$.

Definition 2.2 ([30]) The bivariate function $G: [0, 1] \times [0, 1] \rightarrow [0, 1]$ is referred to as the grouping function, when it fulfils prerequisites below, $\forall s, t, v \in [0, 1]$:

- (a) G fulfils exchangeability;
- (b) $G(s, t) = 0$ when and only when $s = 0$ and $t = 0$;
- (c) $G(s, t) = 1$ when and only when $s = 1$ or $t = 1$;
- (d) $G(s, t) \leq_1 G(s, v)$ if $t \leq_1 v$;
- (e) G fulfils continuity.

Example 2.2 The bivariate functions below are grouping functions, $\forall s, t \in [0, 1]$:

- (a) $G_{\text{mM}}(s, t) = 1 - \min(1 - s, 1 - t) \max((1 - s)^2, (1 - t)^2)$;
- (b) $G_p(s, t) = 1 - (1 - s)^p (1 - t)^p$, for $p > 0$ and $p \neq 1$;
- (c) $G_{\text{DB}}(s, t) = \begin{cases} \frac{s + t - 2st}{2 - s - t}, & \text{if } s + t \neq 2, \\ 1, & \text{if } s + t = 2. \end{cases}$;
- (d) $G_{\text{min}}(s, t) = 1 - \min\{\sqrt{1 - s}, \sqrt{1 - t}\}$.

Definition 2.3 ([31]) An affiliation function $\mu_E(s)$ and a non-affiliation function $\nu_E(s)$ portray an IFS E in S . S is a set that is not empty. And the IFS E be denoted as

$$E = \{(s, \mu_E(s), \nu_E(s)) \mid s \in S\}.$$

In which $\mu_E(s), \nu_E(s) \in [0, 1]$ and satisfies the term of $0 \leq \mu_E(s) + \nu_E(s) \leq 1$.

Definition 2.4 ([2]) Truth-affiliation function $T_E(s)$, uncertainty-affiliation function $U_E(s)$ and falsity-affiliation function $F_E(s)$ portray the single-valued NS E in S . S is a set that is not empty. And the single-valued NS E is defined as

$$E = \{(s, T_E(s), U_E(s), F_E(s)) \mid s \in S\}.$$

In which $T_E(s), U_E(s), F_E(s) \in [0, 1]$ and satisfies the term of $0 \leq T_E(s) + U_E(s) + F_E(s) \leq 3$.

Definition 2.5 ([32]) The overlap function is a map $O: L^2 \rightarrow L$ on $(L; \leq_L)$ which fulfils monotonicity, commutative and continuity, while it fulfils $O(s, 0_L) = 0_L$, $O(0_L, t) = 0_L$ and $O(1_L, 1_L) = 1_L$, $\forall s, t \in L$; the grouping function is a map $G: L^2 \rightarrow L$ on $(L; \leq_L)$ which fulfils monotonicity, commutative and continuity, while it fulfils $G(s, 1_L) = 1_L$, $G(1_L, t) = 1_L$ and $G(0_L, 0_L) = 0_L$, $\forall s, t \in L$.

Definition 2.6 ([18]) Define the set D^* in the following way,

$$D^* = \{s = (s_1, s_2, s_3) \mid s_1, s_2, s_3 \in [0, 1]\}.$$

If $s \in D^*$, as above, then s has three components s_1, s_2 and s_3 .

$\forall s, t \in D^*$, where $s = (s_1, s_2, s_3)$, t is analogous to s . \leq_1 on D^* is defined as the order relation below,

$$s \leq_1 t \text{ iff } s_1 \leq t_1, s_2 \geq t_2, s_3 \geq t_3.$$

And the definition of the first type inclusion relation \subseteq_1 is analogous to the definition of \leq_1 .

Proposition 2.1 ([18]) $(D^*; \leq_1)$ is a complete lattice.

Definition 2.7 ([18]) The supplement of s is written as below, $\forall s \in D^*$,

$$s^c = (s_3, 1 - s_2, s_1).$$

In particular, $1_{D^*} = (1, 0, 0)$ and $0_{D^*} = (0, 1, 1)$ represent the maximum and minimum in $(D^*; \leq_1)$, respectively.

Proposition 2.2 ([18]) $s \wedge_1 t$ is defined as maximum lower bound of s, t , and expressed as $\inf(s, t)$; $s \vee_1 t$ is defined as minimum upper bound of s, t , and expressed as $\sup(s, t)$, $\forall s, t \in D^*$.

3. Neutrosophic overlap function

This section proposes new concept of neutrosophic overlap function and provides relevant examples, giving the concept and examples of representable and non-representable neutrosophic overlap function. Finally, a new method for constructing representable neutrosophic overlap function through intuitionistic fuzzy overlap function (IFO) is proposed.

Definition 3.1 A neutrosophic overlap function is a map $O: D^* \times D^* \rightarrow D^*$ which fulfils prerequisites below, $\forall s, t, v \in D^*$:

(NO1) O fulfils exchangeability;

(NO2) $O(s, t) \leq_1 O(s, v)$ if $t \leq_1 v$;

(NO3) $O(0_{D^*}, t) = 0_{D^*}$ or $O(s, 0_{D^*}) = 0_{D^*}$;

(NO4) $O(1_{D^*}, 1_{D^*}) = 1_{D^*}$;

(NO5) O fulfils continuity.

Definition 3.2 A neutrosophic grouping function is a map $G: D^* \times D^* \rightarrow D^*$ which fulfils prerequisites below, $\forall s, t, v \in D^*$:

(NG1) G fulfils exchangeability;

(NG2) $G(s, t) \leq_1 G(s, v)$ if $t \leq_1 v$;

(NG3) $G(1_{D^*}, t) = 1_{D^*}$ or $G(s, 1_{D^*}) = 1_{D^*}$;

(NG4) $G(0_{D^*}, 0_{D^*}) = 0_{D^*}$;

(NG5) G fulfils continuity.

Example 3.1 The following binary functions are neutrosophic overlap functions, $\forall s, t \in D^*$,

$$(1) \quad O_{\text{mM}}(s, t) = (O_{\text{mM}}(s_1, t_1), G_{\text{mM}}(s_2, t_2), G_{\text{mM}}(s_3, t_3)) \\ = (\min(s_1, t_1) \max(s_1^2, t_1^2), 1 - \min(1 - s_2, 1 - t_2) \max((1 - s_2)^2, (1 - t_2)^2), 1 - \min(1 - s_3, 1 - t_3) \max((1 - s_3)^2, (1 - t_3)^2));$$

$$(2) \quad O_p(s, t) = (O_p(s_1, t_1), G_p(s_2, t_2), G_p(s_3, t_3)) = (s_1^p t_1^p, 1 - (1 - s_2)^p (1 - t_2)^p, 1 - (1 - s_3)^p (1 - t_3)^p), \text{ for } p > 0 \text{ and } p \neq 1;$$

$$(3) \quad O_{\text{DB}}(s, t) = (O_{\text{DB}}(s_1, t_1), G_{\text{DB}}(s_2, t_2), G_{\text{DB}}(s_3, t_3)) = \left(\frac{2s_1 t_1}{s_1 + t_1}, \frac{s_2 + t_2 - 2s_2 t_2}{2 - s_2 - t_2}, \frac{s_3 + t_3 - 2s_3 t_3}{2 - s_3 - t_3} \right), \text{ for } s_1 \neq t_1 \neq 0, \\ s_2 \neq t_2 \neq 1 \text{ and } s_3 \neq t_3 \neq 1;$$

$$(4) \quad O_{\text{min}}(s, t) = (O_{\text{min}}(s_1, t_1), G_{\text{min}}(s_2, t_2), G_{\text{min}}(s_3, t_3)) \\ = (\min\{\sqrt{s_1}, \sqrt{t_1}\}, 1 - \min\{\sqrt{1 - s_2}, \sqrt{1 - t_2}\}, 1 - \min\{\sqrt{1 - s_3}, \sqrt{1 - t_3}\}).$$

Example 3.2 The following binary functions are neutrosophic grouping functions, $\forall s, t \in D^*$,

$$(1) \quad G_{\text{mM}}(s, t) = (G_{\text{mM}}(s_1, t_1), O_{\text{mM}}(s_2, t_2), O_{\text{mM}}(s_3, t_3)) \\ = (1 - \min(1 - s_1, 1 - t_1) \max((1 - s_1)^2, (1 - t_1)^2), \min(s_2, t_2) \max(s_2^2, t_2^2), \min(s_3, t_3) \max(s_3^2, t_3^2));$$

(2) $G_p(s, t) = (G_p(s_1, t_1), O_p(s_2, t_2), O_p(s_3, t_3)) = (1 - (1 - s_1)^p (1 - t_1)^p, s_2^p t_2^p, s_3^p t_3^p)$, for $p > 0$ and $p \neq 1$;

(3) $G_{DB}(s, t) = (G_{DB}(s_1, t_1), O_{DB}(s_2, t_2), O_{DB}(s_3, t_3)) = (\frac{s_1 + t_1 - 2s_1 t_1}{2 - s_1 - t_1}, \frac{2s_2 t_2}{s_2 + t_2}, \frac{2s_3 t_3}{s_3 + t_3})$, for $s_1 \neq t_1 \neq 1$, $s_2 \neq t_2 \neq 0$ and $s_3 \neq t_3 \neq 0$;

(4) $G_{min}(s, t) = (G_{min}(s_1, t_1), O_{min}(s_2, t_2), O_{min}(s_3, t_3))$
 $= (1 - \min\{\sqrt{1 - s_1}, \sqrt{1 - t_1}\}, \min\{\sqrt{s_2}, \sqrt{t_2}\}, \min\{\sqrt{s_3}, \sqrt{t_3}\})$.

Theorem 3.1 Let O is a bivariate operation on D^* , $\forall s, t \in D^*$,

$$O(s, t) = (O(s_1, t_1), G_1(s_2, t_2), G_2(s_3, t_3)).$$

Then O is a neutrosophic overlap function, where O is the overlap function, G_1 and G_2 are grouping functions on $[0, 1]$.

Proof. $\forall s, t, v \in D^*$ in which $s = (s_1, s_2, s_3)$, t and v are analogous to s .

(NO1) Since G_1 and G_2 are grouping functions, O is the overlap function, then $O(s_1, t_1) = O(t_1, s_1)$, $G_1(t_2, s_2) = G_1(s_2, t_2)$ and $G_2(t_3, s_3) = G_2(s_3, t_3)$, thus O fulfils exchangeability.

(NO2) Let $t \leq v$, then $O(s_1, t_1) \leq O(s_1, v_1)$, $G_1(s_2, t_2) \geq G_1(s_2, v_2)$, $G_2(s_3, t_3) \geq G_2(s_3, v_3)$. Therefore, $O(s, t) \leq O(s, v)$.

(NO3) $O(0_{D^*}, t) = (O(0, t_1), G_1(1, t_2), G_2(1, t_3)) = (0, 1, 1) = 0_{D^*}$, and $O(s, 0_{D^*}) = (O(s_1, 0), G_1(s_2, 1), G_2(s_3, 1)) = (0, 1, 1) = 0_{D^*}$.

(NO4) $O(1_{D^*}, 1_{D^*}) = (O(1, 1), G_1(0, 0), G_2(0, 0)) = (1, 0, 0) = 1_{D^*}$.

(NO5) Firstly, we prove left continuous. That is, prove that this equation $O(s, \bigvee_{i \in I} t_i) = \bigvee_{i \in I} O(s, t_i)$ holds. Because the overlap function and the grouping function are continuous, so $O(s_1, \bigvee_{i \in I} t_i) = \bigvee_{i \in I} O(s_1, t_i)$, $G_2(s_3, \bigvee_{i \in I} t_i) = \bigvee_{i \in I} G_2(s_3, t_i)$ and $G_1(s_2, \bigvee_{i \in I} t_i) = \bigvee_{i \in I} G_1(s_2, t_i)$ is valid.

So we can get

$$\begin{aligned} O(s, \bigvee_{i \in I} t_i) &= (O(s_1, \bigvee_{i \in I} t_i), G_1(s_2, \bigvee_{i \in I} t_i), G_2(s_3, \bigvee_{i \in I} t_i)) \\ &= (\bigvee_{i \in I} O(s_1, t_i), \bigvee_{i \in I} G_1(s_2, t_i), \bigvee_{i \in I} G_2(s_3, t_i)) \\ &= \bigvee_{i \in I} O(s, t_i) \end{aligned}$$

In this way, show that O is left continuous.

Likewise, $O(s, \bigwedge_{i \in I} t_i) = \bigwedge_{i \in I} O(s, t_i)$ is simple to prove, state clearly that O is right continuous. To sum up, O is shown to be a neutrosophic overlap function.

Theorem 3.2 G is a bivariate operation on D^* , $\forall s, t \in D^*$,

$$G(s, t) = (G(s_1, t_1), O_1(s_2, t_2), O_2(s_3, t_3)).$$

Then G can be called the neutrosophic grouping function, where G is the grouping function, O_1 and O_2 are overlap functions on $[0, 1]$.

Proof. The procedure for proving analogy **Theorem 3.1**.

Above **Theorem 3.1** supplies the measure for constructing neutrosophic overlap function using overlap function O and grouping functions G_1, G_2 which are defined on $[0, 1]$. But it requires a condition that $O = (O, G_1, G_2)$ holds. According to this condition, we bring in the concept of representable neutrosophic overlap function.

Definition 3.3 A neutrosophic overlap function O is referred to as representable, when and only when, there exists O which is an overlap function on $[0, 1]$ and G_1, G_2 which are grouping functions on $[0, 1]$ satisfying, $\forall s, t \in D^*$,

$$O(s, t) = (O(s_1, t_1), G_1(s_2, t_2), G_2(s_3, t_3)).$$

Example 3.3 The representable neutrosophic overlap function is shown below, $\forall s, t \in D^*$,

$$O(s, t) = (O_{DB}(s_1, t_1), G_p(s_2, t_2), G_{mM}(s_3, t_3)).$$

Proof. The first step verifies that O is a neutrosophic overlap function holds. $\forall s, t, v \in D^*$ in which $s = (s_1, s_2, s_3)$, t and v are analogous to s .

(NO1) Let $G_1 = G_p, G_2 = G_{mM}$ ($p=2$) are grouping functions, $O = O_{DB}$ is an overlap function on $[0, 1]$. Since $O_{DB}(s_1, t_1) = O_{DB}(t_1, s_1), G_p(s_2, t_2) = G_p(t_2, s_2), G_{mM}(s_3, t_3) = G_{mM}(s_3, t_3)$, thus O fulfils exchangeability.

(NO2) Let $t \leq v$, then $O_{DB}(s_1, t_1) \leq O_{DB}(s_1, v_1), G_p(s_2, t_2) \geq G_p(s_2, v_2), G_{mM}(s_3, t_3) \geq G_{mM}(s_3, v_3)$. Therefore, $O(s, t) \leq O(s, v)$.

(NO3) $O(0_{D^*}, t) = (O_{DB}(0, t_1), G_p(1, t_2), G_{mM}(1, t_3)) = (0, 1, 1) = 0_{D^*}$, and $O(s, 0_{D^*}) = (O_{DB}(s_1, 0), G_p(s_2, 1), G_{mM}(s_3, 1)) = (0, 1, 1) = 0_{D^*}$.

(NO4) $O(1_{D^*}, 1_{D^*}) = (O_{DB}(1, 1), G_p(0, 0), G_{mM}(0, 0)) = (1, 0, 0) = 1_{D^*}$.

(NO5) Firstly, we prove left continuous. That is, prove that this equation $O(s, \bigvee_{i \in I} t_i) = \bigvee_{i \in I} O(s, t_i)$ holds. Because of an overlap function O_{DB} and grouping functions G_p and G_{mM} are continuous, so $O_{DB}(s_1, \bigvee_{i \in I} t_i) = \bigvee_{i \in I} O_{DB}(s_1, t_i), G_{mM}(s_3, \bigvee_{i \in I} t_i) = \bigvee_{i \in I} G_{mM}(s_3, t_i)$ and $G_p(s_2, \bigvee_{i \in I} t_i) = \bigvee_{i \in I} G_p(s_2, t_i)$ is valid.

So we can get

$$\begin{aligned} O(s, \bigvee_{i \in I} t_i) &= (O_{DB}(s_1, \bigvee_{i \in I} t_i), G_p(s_2, \bigvee_{i \in I} t_i), G_{mM}(s_3, \bigvee_{i \in I} t_i)) \\ &= (\bigvee_{i \in I} O_{DB}(s_1, t_i), \bigvee_{i \in I} G_p(s_2, t_i), \bigvee_{i \in I} G_{mM}(s_3, t_i)) \\ &= \bigvee_{i \in I} O(s, t_i) \end{aligned}$$

In this way, show that O is left continuous.

Likewise, $O(s, \bigwedge_{i \in I} t_i) = \bigwedge_{i \in I} O(s, t_i)$ is simple to prove, state clearly that O is right continuous. To sum up, O is shown to be the neutrosophic overlap function.

Finally, it is simple to show that fulfils $O(s, t) = (O_{DB}(s_1, t_1), G_p(s_2, t_2), G_{mM}(s_3, t_3))$, so it must be the representable neutrosophic overlap function.

Definition 3.4 The neutrosophic overlap function O is known as standard representable, when and only when, there exists G which is a grouping function on $[0, 1]$ and O which is an overlap function on $[0, 1]$ satisfying, $\forall s, t \in D^*$,

$$O(s, t) = (O(s_1, t_1), G(s_2, t_2), G(s_3, t_3)).$$

Example 3.4 The standard representable neutrosophic overlap function is as follows, $\forall s, t \in D^*$,

$$O(s, t) = (O_{DB}(s_1, t_1), G_p(s_2, t_2), G_p(s_3, t_3)).$$

Proof. This procedure for proving analogy **Example 3.3**.

Definition 3.5 The N-dual representable neutrosophic overlap function O by the following being defined by, $\forall s, t \in D^*$,

$$O(s, t) = (O(s_1, t_1), G(s_2, t_2), G(s_3, t_3)).$$

O and G has dual relation as follows,

$$O(s_1, t_1) = 1 - G(1 - s_1, 1 - t_1).$$

Example 3.5 The N-dual representable neutrosophic overlap function is as follows, $\forall s, t \in D^*$,

$$O(s, t) = (O_p(s_1, t_1), G_p(s_2, t_2), G_p(s_3, t_3)).$$

Proof. This procedure for proving analogy **Example 3.3**.

Definition 3.6 G is referred to as representable neutrosophic grouping function, when and only when, there obtains the grouping function G and the overlap functions O_1, O_2 on $[0, 1]$ satisfying, $\forall s, t \in D^*$,

$$G(s, t) = (G(s_1, t_1), O_1(s_2, t_2), O_2(s_3, t_3)).$$

Other concepts can be derived from the analogy of the neutrosophic overlap function.

In recent years, there have been many extensions of overlap functions. However, due to the limitations of existing definitions in addressing practical issues by using intuitionistic fuzzy information, scholars have proposed IFO. In the preceding paragraphs, the representable neutrosophic overlap function is proposed and further the below propositions propose a method to construct new representable neutrosophic overlap function (grouping function) with IFO (intuitionistic fuzzy grouping function).

Proposition 3.1 Where $m = (m_1, m_3), n = (n_1, n_3), \forall m, n \in L$. O is an IFO while satisfying $O(m, n) = (O(m_1, n_1), G_2(m_3, n_3))$, with O being an overlap function on $[0, 1]$, G_2 being a grouping function on $[0, 1]$. Suppose G_1 is a grouping function on $[0, 1]$ satisfying

$$0 \leq O(s_1, t_1) + G_1(s_2, t_2) + G_2(s_3, t_3) \leq 3.$$

Then $O(s, t) = (O(s_1, t_1), G_1(s_2, t_2), G_2(s_3, t_3))$ is called the representable neutrosophic overlap function, $\forall s, t \in D^*$.

Proof. First, we can get $O(m, n) = (O(m_1, n_1), G_2(m_3, n_3))$ which is an IFO, and then we add another grouping function G_1 , satisfying $0 \leq O(s_1, t_1) + G_1(s_2, t_2) + G_2(s_3, t_3) \leq 3$.

$\forall s, t, v \in D^*$ in which $s = (s_1, s_2, s_3), t$ and v are analogous to s .

(NO1) Since $O(s_1, t_1) = O(t_1, s_1), G_2(s_3, t_3) = G_2(t_3, s_3), G_1(s_2, t_2) = G_1(t_2, s_2)$, then $O(s, t) = O(t, s)$, thus it shown that O fulfils exchangeability.

(NO2) Let $t \leq_1 v$, then $O(s_1, t_1) \leq O(s_1, v_1), G_2(s_3, t_3) \geq G_2(s_3, v_3), G_1(s_2, t_2) \geq G_1(s_2, v_2)$. Therefore, $O(s, t) \leq_1 O(s, v)$.

(NO3) $O(0_{D^*}, t) = (O(0, t_1), G_1(1, t_2), G_2(1, t_3)) = (0, 1, 1) = 0_{D^*}$, and $O(s, 0_{D^*}) = (O(s_1, 0), G_1(s_2, 1), G_2(s_3, 1)) = (0, 1, 1) = 0_{D^*}$.

(NO4) $O(1_{D^*}, 1_{D^*}) = (O(1, 1), G_1(0, 0), G_2(0, 0)) = (1, 0, 0) = 1_{D^*}$.

(NO5) Firstly, we prove left continuous. That is, prove that this equation $O(s, \bigvee_{i \in I} t_i) = \bigvee_{i \in I} O(s, t_i)$ holds. Because the overlap function O and the grouping functions G_2, G_1 are continuous, $O(s_1, \bigvee_{i \in I} t_i) = \bigvee_{i \in I} O(s_1, t_i), G_2(s_3, \bigvee_{i \in I} t_i) = \bigvee_{i \in I} G_2(s_3, t_i)$ and $G_1(s_2, \bigvee_{i \in I} t_i) = \bigvee_{i \in I} G_1(s_2, t_i)$ is holding.

So we can get

$$\begin{aligned} O(s, \bigvee_{i \in I} t_i) &= (O(s_1, \bigvee_{i \in I} t_i), G_1(s_2, \bigvee_{i \in I} t_i), G_2(s_3, \bigvee_{i \in I} t_i)) \\ &= (\bigvee_{i \in I} O(s_1, t_i), \bigvee_{i \in I} G_1(s_2, t_i), \bigvee_{i \in I} G_2(s_3, t_i)) \\ &= \bigvee_{i \in I} O(s, t_i) \end{aligned}$$

In this way, show that O is left continuous.

Likewise, $O(s, \bigwedge_{i \in I} t_i) = \bigwedge_{i \in I} O(s, t_i)$ is simple to prove, state clearly that O is right continuous. To sum up, O is shown to be the neutrosophic overlap function.

It is simple to show that satisfies $O(s, t) = (O(s_1, t_1), G_1(s_2, t_2), G_2(s_3, t_3))$, so O is a representable neutrosophic overlap function.

Proposition 3.2 Where $m = (m_1, m_3), n = (n_1, n_3), \forall m, n \in L$. G is an intuitionistic fuzzy grouping function, while satisfying the fact that $G(m, n) = (G(m_1, n_1), O_2(m_3, n_3))$, with G being a grouping function on $[0, 1]$, O_2 being an overlap function on $[0, 1]$. Suppose O_1 is an overlap function on $[0, 1]$ satisfying,

$$0 \leq G(s_1, t_1) + O_1(s_2, t_2) + O_2(s_3, t_3) \leq 3.$$

Then $G(s, t) = (G(s_1, t_1), O_1(s_2, t_2), O_2(s_3, t_3))$ is a representable neutrosophic grouping function, $\forall s, t \in D^*$.

Proof. This procedure for proving analogy **Proposition 3.1**.

The dual relation between triangular norm and triangular conorm in relation to fuzzy negation can be characterized by De Morgan triple, which is a proper expression for the relationship between triangular norm, triangular conorm and fuzzy negation [18]. There are also corresponding studies on NS. Based on the close connection between triangular norm and overlap function, one can naturally consider De Morgan neutrosophic triple about neutrosophic overlap function, neutrosophic grouping function and neutrosophic negation. First, neutrosophic negation as an extension of fuzzy negation can be denoted by the method below.

Definition 3.7 ([18]) A neutrosophic negaton is a map $N: D^* \rightarrow D^*$ that fulfils prerequisites below:

- (a) $N(t) \geq_1 N(s), \forall s, t \in D^*$ such as $t \leq_1 s$;
- (b) $N(0_{D^*}) = 1_{D^*}$;
- (c) $N(1_{D^*}) = 0_{D^*}$.

N is referred to as the involutive neutrosophic negaton when and only when that fulfils $N(N(s)) = s, \forall s \in D^*$.

An involutive neutrosophic negaton $Ng: D^* \rightarrow D^*$ satisfies the following, where $s = (s_1, s_2, s_3), \forall s \in D^*$,

$$Ng(s_1, s_2, s_3) = (s_3, 1-s_2, s_1).$$

Further, we define such Ng as the standard neutrosophic negaton.

Definition 3.8 O, G and N are a neutrosophic overlap function, a neutrosophic grouping function, and a neutrosophic negation, respectively.

For this triple (O, N, G) if the conditions below holding true, $\forall s, t \in D^*$,

$$N(O(s, t)) = G(N(s), N(t));$$

$$N(G(s, t)) = O(N(s), N(t)).$$

Then such the triple is referred to as De Morgan neutrosophic triple. In addition, O and G have a dual relationship in relation to N .

Theorem 3.3 Suppose neutrosophic negaton N is involutory, that it fulfils $N(N(s)) = s, \forall s \in D^*$.

(a) Assume G is the neutrosophic grouping function, O is expressed in the following form, $\forall s, t \in D^*$,

$$O(s, t) = N(G(N(s), N(t))).$$

Then O is the neutrosophic overlap function. Moreover, (O, N, G) is De Morgan neutrosophic triple.

(b) Assume O is the neutrosophic overlap function, G is expressed in the following form, $\forall s, t \in D^*$,

$$G(s, t) = N(O(N(s), N(t))).$$

Then G is the neutrosophic grouping function. Moreover, (O, N, G) is De Morgan neutrosophic triple.

Proof. (a) Suppose N, G are the involutory neutrosophic negaton and the neutrosophic grouping function, respectively. $\forall s, t, v \in D^*$ in which $s = (s_1, s_2, s_3), t$ and v are analogous to s .

(NO1) It is pretty simple to justify that $O(s, t) = N(G(N(s), N(t))) = N(G(N(t), N(s))) = O(t, s), O$ fulfils exchangeability.

(NO2) Let $t \leq_1 v, O(s, t) = N(G(N(s), N(t))), O(s, v) = N(G(N(s), N(v))),$ because N is non-increasing, then $N(t) \geq_1 N(v)$. Moreover $G(s, t) \leq_1 G(s, v)$ and when $t \leq_1 v$, then $G(N(s), N(t)) \geq_1 G(N(s), N(v))$. Hence $N(G(N(s), N(t))) \leq_1 N(G(N(s), N(v)))$, then $O(s, t) \leq_1 O(s, v)$.

(NO3) $O(0_{D^*}, t) = N(G(N(0_{D^*}), N(t))) = N(G(1_{D^*}, N(t))) = N(1_{D^*}) = 0_{D^*}$, similarly $O(s, 0_{D^*}) = N(G(N(s), N(0_{D^*}))) = 0_{D^*}$.

(NO4) $O(1_{D^*}, 1_{D^*}) = N(G(N(1_{D^*}), N(1_{D^*}))) = N(G(0_{D^*}, 0_{D^*})) = N(0_{D^*}) = 1_{D^*}$.

(NO5) Firstly, we prove left continuous. That is, prove that this equation $O(s, \bigvee_{i \in I} t_i) = \bigvee_{i \in I} O(s, t_i)$ holds. As a result of $O(s, \bigvee_{i \in I} t_i) = N(G(N(s), N(\bigvee_{i \in I} t_i))) = N(G(N(s), \bigwedge_{i \in I} N(t_i))) = N(\bigwedge_{i \in I} G(N(s), N(t_i))) =$

$\forall_{i \in I} N(G(N(s), N(t))) = \forall_{i \in I} O(s, t_i)$. Then we could get $O(s, \forall_{i \in I} t_i) = \forall_{i \in I} O(s, t_i)$. In this way, show that O is left continuous.

Likewise, $O(s, \wedge_{i \in I} t_i) = \wedge_{i \in I} O(s, t_i)$ is simple to prove, state clearly that O is right continuous. To sum up, $O(s, t)$ is shown to be the neutrosophic overlap function.

Moreover, (O, N, G) is the De Morgan neutrosophic triple.

(b) Likewise, suppose O is the neutrosophic overlap function and that G can be shown to be the neutrosophic grouping function, (O, N, G) would be the De Morgan neutrosophic triple.

Example 3.6 The following functions are the neutrosophic overlap(grouping) functions, which are dual in relation to Ng , $\forall s, t \in D^*$,

(1) $O_{mM}(s, t) = (O_{mM}(s_1, t_1), G_{mM}(s_2, t_2), G_{mM}(s_3, t_3))$ and $G_{mM}(s, t) = (G_{mM}(s_1, t_1), O_{mM}(s_2, t_2), O_{mM}(s_3, t_3))$;

In fact, $O_{mM}(N(s), N(t)) = O_{mM}((s_3, 1-s_2, s_1), (t_3, 1-t_2, t_1)) = (O_{mM}(s_3, t_3), G_{mM}(1-s_2, 1-t_2), G_{mM}(s_1, t_1))$, then $N(O_{mM}(N(s), N(t))) = N(O_{mM}(s_3, t_3), G_{mM}(1-s_2, 1-t_2), G_{mM}(s_1, t_1)) = (G_{mM}(s_1, t_1), 1-G_{mM}(1-s_2, 1-t_2), O_{mM}(s_3, t_3)) = (G_{mM}(s_1, t_1), O_{mM}(s_2, t_2), O_{mM}(s_3, t_3)) = G_{mM}(s, t)$. Thus, O_{mM} and G_{mM} are dual with respect to Ng .

(2) $O_p(s, t) = (O_p(s_1, t_1), G_p(s_2, t_2), G_p(s_3, t_3))$ and $G_p(s, t) = (G_p(s_1, t_1), O_p(s_2, t_2), O_p(s_3, t_3))$;

(3) $O_{DB}(s, t) = (O_{DB}(s_1, t_1), G_{DB}(s_2, t_2), G_{DB}(s_3, t_3))$ and $G_{DB}(s, t) = (G_{DB}(s_1, t_1), O_{DB}(s_2, t_2), O_{DB}(s_3, t_3))$;

(4) $O_{min}(s, t) = (O_{min}(s_1, t_1), G_{min}(s_2, t_2), G_{min}(s_3, t_3))$ and $G_{min}(s, t) = (G_{min}(s_1, t_1), O_{min}(s_2, t_2), O_{min}(s_3, t_3))$;

(5) $O(s, t) = (O_{DB}(s_1, t_1), G_p(s_2, t_2), G_p(s_3, t_3))$ and $G(s, t) = (G_p(s_1, t_1), O_p(s_2, t_2), O_{DB}(s_3, t_3))$.

We give the following theorem for non-representable neutrosophic overlap function.

Theorem 3.4 Let O be a map on D^* below, $\forall s, t \in D^*$,

$$O(s, t) = \begin{cases} 0_{D^*}, & \text{if } s = 0_{D^*} \text{ or } t = 0_{D^*}, \\ 1_{D^*}, & \text{if } s = t = 1_{D^*}, \\ (s_1 t_1, s_3 t_3, s_3 t_3), & \text{otherwise.} \end{cases}$$

Then O is a non-representable neutrosophic overlap function.

Proof. The first step is to verify that O is the neutrosophic overlap function. $\forall s, t, v, u \in D^*$ in which $s = (s_1, s_2, s_3)$, t, u and v are analogous to s .

(NO1) The proof that O fulfils exchangeability is very straightforward.

(NO2) Let $t \leq_1 v$. The obvious one is $O(s, t) \leq_1 O(s, v)$.

(NO3) $O(0_{D^*}, t) = O(s, 0_{D^*}) = (0, 1, 1) = 0_{D^*}$.

(NO4) $O(1_{D^*}, 1_{D^*}) = (1, 0, 0) = 1_{D^*}$.

(NO5) Firstly, we prove left continuous. That is, prove that this equation $O(s, \forall_{i \in I} t_i) = \forall_{i \in I} O(s, t_i)$ holds. As a result of $O(s, \forall_{i \in I} t_i) = (s_1^* \max(t_i), s_3^* \max(t_i), s_3^* \max(t_i))$; $\forall_{i \in I} O(s, t_i) = (s_1^* t_i, s_3^* t_i, s_3^* t_i) \vee (s_1^* t_2, s_3^* t_3, s_3^* t_3) \vee (s_1^* t_3, s_3^* t_3, s_3^* t_3) = (s_1^* \max(t_i), s_3^* \max(t_i), s_3^* \max(t_i))$. We can get $O(s, \forall_{i \in I} t_i) = \forall_{i \in I} O(s, t_i)$. Therefore, it is show that O fulfils left continuity.

Likewise, $O(s, \wedge_{i \in I} t_i) = \wedge_{i \in I} O(s, t_i)$ is simple to prove, state clearly that O is right continuous. To sum up, $O(s, t)$ is shown to be the neutrosophic overlap function.

And then, verify that for the representable neutrosophic overlap function O whether there has the overlap function O and grouping functions G_1, G_2 on $[0, 1]$ fulfilling the form $O = (O, G_1, G_2)$.

Have $s = (0.3, 0.5, 0.6)$, $u = (0.3, 0.5, 0.2)$ and $t = (0.4, 0.5, 0.8)$ respectively. From $O(s, t) = (0.12, 0.48, 0.48)$ and $O(u, t) = (0.12, 0.16, 0.16)$. We get $G_1(s_2, t_2) = 0.48$ and $G_1(u_2, t_2) = 0.16$, so $G_1(u_2, t_2) \neq G_1(s_2, t_2)$. Thus, G_1 is not independent from s_3 , which suggests that O is non-representable.

In addition, the neutrosophic grouping function G is the dual of O in relation to the standard neutrosophic negaton Ng , defined as below, $\forall s, t \in D^*$,

$$G(s, t) = \begin{cases} 0_{D^*}, & \text{if } s = t = 0_{D^*}, \\ 1_{D^*}, & \text{if } s = 1_{D^*} \text{ or } t = 1_{D^*}, \\ (1 - (1 - s_1)(1 - t_1), 1 - (1 - s_3)(1 - t_3), 1 - (1 - s_3)(1 - t_3)), & \text{otherwise.} \end{cases}$$

Then G is a non-representable neutrosophic grouping function.

4. NRI derived from neutrosophic overlap function

This section would bring in the concept of NRI on D^* and research fundamental properties of NRI. First, the notion of neutrosophic implication is introduced on D^* .

Definition 4.1 ([18]) The map $I: (D^*)^2 \rightarrow D^*$ is known as the neutrosophic implication when it fulfils the prerequisites below, $\forall s, u, t, v \in D^*$:

- (a) I is non-increasing for the first variable component (in relation to the order relation \leq_1), which means that when $s \leq_1 u$, there is $I(s, t) \geq_1 I(u, t)$;
- (b) I is non-decreasing for the second variable component (in relation to the order relation \leq_1), which means that when $t \leq_1 v$, there is $I(s, t) \leq_1 I(s, v)$;
- (c) $I(0_{D^*}, 0_{D^*}) = 1_{D^*}$;
- (d) $I(1_{D^*}, 1_{D^*}) = 1_{D^*}$;
- (e) $I(1_{D^*}, 0_{D^*}) = 0_{D^*}$.

Definition 4.2 Suppose $I: (D^*)^2 \rightarrow D^*$ is a binary map. A neutrosophic overlap function O exists which enables the following condition to hold, $\forall s, t, h \in D^*$,

$$I(s, t) = \sup\{h \mid h \in D^*, O(s, h) \leq_1 t\}.$$

Thus such $I: (D^*)^2 \rightarrow D^*$ is referred to as the NRI.

When I is a NRI derived from a neutrosophic overlap function O , it is written as I_o .

Additionally, a neutrosophic overlap function O fulfils the residual principle, $\forall s, t, h \in D^*$:

$$h \leq_1 I_o(s, t) \text{ iff } O(s, h) \leq_1 t.$$

Example 4.1 The functions below are NRIs derived from neutrosophic overlap functions in **Example 3.1**, $\forall s, t \in D^*$,

(1)

$$I_{O_{\text{min}}}(s, t) = \begin{cases} (1, 0, 0), & \text{if } s_1 \leq t_1, s_2 \geq t_2 \text{ and } s_3 \geq t_3, \\ (1, \max\{1 - \sqrt{\frac{1-t_2}{1-s_2}}, 1 - \frac{1-t_2}{(1-s_2)^2}\}, \max\{1 - \sqrt{\frac{1-t_3}{1-s_3}}, 1 - \frac{1-t_3}{(1-s_3)^2}\}), & \text{if } s_1 \leq t_1, s_2 < t_2 \text{ and } s_3 < t_3, \\ (1, 0, \max\{1 - \sqrt{\frac{1-t_3}{1-s_3}}, 1 - \frac{1-t_3}{(1-s_3)^2}\}), & \text{if } s_1 \leq t_1, s_2 \geq t_2 \text{ and } s_3 < t_3, \\ (1, \max\{1 - \sqrt{\frac{1-t_2}{1-s_2}}, 1 - \frac{1-t_2}{(1-s_2)^2}\}, 0), & \text{if } s_1 \leq t_1, s_2 < t_2 \text{ and } s_3 \geq t_3, \\ (\min\{\sqrt{\frac{t_1}{s_1}}, \frac{t_1}{s_1^2}\}, \max\{1 - \sqrt{\frac{1-t_2}{1-s_2}}, 1 - \frac{1-t_2}{(1-s_2)^2}\}, \max\{1 - \sqrt{\frac{1-t_3}{1-s_3}}, 1 - \frac{1-t_3}{(1-s_3)^2}\}), & \text{if } s_1 > t_1, s_2 < t_2 \text{ and } s_3 < t_3, \\ (\min\{\sqrt{\frac{t_1}{s_1}}, \frac{t_1}{s_1^2}\}, 0, 0), & \text{if } s_1 > t_1, s_2 \geq t_2 \text{ and } s_3 \geq t_3, \\ (\min\{\sqrt{\frac{t_1}{s_1}}, \frac{t_1}{s_1^2}\}, \max\{1 - \sqrt{\frac{1-t_2}{1-s_2}}, 1 - \frac{1-t_2}{(1-s_2)^2}\}, 0), & \text{if } s_1 > t_1, s_2 < t_2 \text{ and } s_3 \geq t_3, \\ (\min\{\sqrt{\frac{t_1}{s_1}}, \frac{t_1}{s_1^2}\}, 0, \max\{1 - \sqrt{\frac{1-t_3}{1-s_3}}, 1 - \frac{1-t_3}{(1-s_3)^2}\}), & \text{if } s_1 > t_1, s_2 \geq t_2 \text{ and } s_3 < t_3. \end{cases}$$

(2)

$$I_{O_{ps2}}(s, t) = \begin{cases} \left(\frac{\sqrt{t_1}}{s_1}, 1 - \frac{\sqrt{1-t_2}}{1-s_2}, 1 - \frac{\sqrt{1-t_3}}{1-s_3}\right), & \text{if } s_1 > \sqrt{t_1}, s_2 < 1 - \sqrt{1-t_2} \text{ and } s_3 < 1 - \sqrt{1-t_3}, \\ \left(\frac{\sqrt{t_1}}{s_1}, 0, 0\right), & \text{if } s_1 > \sqrt{t_1}, s_2 \geq 1 - \sqrt{1-t_2} \text{ and } s_3 \geq 1 - \sqrt{1-t_3}, \\ \left(\frac{\sqrt{t_1}}{s_1}, 1 - \frac{\sqrt{1-t_2}}{1-s_2}, 0\right), & \text{if } s_1 > \sqrt{t_1}, s_2 < 1 - \sqrt{1-t_2} \text{ and } s_3 \geq 1 - \sqrt{1-t_3}, \\ \left(\frac{\sqrt{t_1}}{s_1}, 0, 1 - \frac{\sqrt{1-t_3}}{1-s_3}\right), & \text{if } s_1 > \sqrt{t_1}, s_2 \geq 1 - \sqrt{1-t_2} \text{ and } s_3 < 1 - \sqrt{1-t_3}, \\ \left(1, 1 - \frac{\sqrt{1-t_2}}{1-s_2}, 1 - \frac{\sqrt{1-t_3}}{1-s_3}\right), & \text{if } s_1 \leq \sqrt{t_1}, s_2 < 1 - \sqrt{1-t_2} \text{ and } s_3 < 1 - \sqrt{1-t_3}, \\ (1, 0, 0), & \text{if } s_1 \leq \sqrt{t_1}, s_2 \geq 1 - \sqrt{1-t_2} \text{ and } s_3 \geq 1 - \sqrt{1-t_3}, \\ \left(1, 0, 1 - \frac{\sqrt{1-t_3}}{1-s_3}\right), & \text{if } s_1 \leq \sqrt{t_1}, s_2 \geq 1 - \sqrt{1-t_2} \text{ and } s_3 < 1 - \sqrt{1-t_3}, \\ \left(1, 1 - \frac{\sqrt{1-t_2}}{1-s_2}, 0\right), & \text{if } s_1 \leq \sqrt{t_1}, s_2 < 1 - \sqrt{1-t_2} \text{ and } s_3 \geq 1 - \sqrt{1-t_3}. \end{cases}$$

(3)

$$I_{O_{ob}}(s, t) = \begin{cases} \left(\min\left\{1, \frac{s_1 t_1}{2s_1 - t_1}\right\}, \frac{2t_2 - s_2 t_2 - s_2}{1 - 2s_2 + t_2}, \frac{2t_3 - s_3 t_3 - s_3}{1 - 2s_3 + t_3}\right), & \text{if } s_1 > \frac{1}{2}t_1, s_2 < \frac{1}{2} + \frac{1}{2}t_2 \text{ and } s_3 < \frac{1}{2} + \frac{1}{2}t_3, \\ \left(\min\left\{1, \frac{s_1 t_1}{2s_1 - t_1}\right\}, 0, \frac{2t_3 - s_3 t_3 - s_3}{1 - 2s_3 + t_3}\right), & \text{if } s_1 > \frac{1}{2}t_1, 1 > s_2 \geq \frac{1}{2} + \frac{1}{2}t_2 \text{ and } s_3 < \frac{1}{2} + \frac{1}{2}t_3, \\ \left(\min\left\{1, \frac{s_1 t_1}{2s_1 - t_1}\right\}, \frac{2t_2 - s_2 t_2 - s_2}{1 - 2s_2 + t_2}, 0\right), & \text{if } s_1 > \frac{1}{2}t_1, s_2 < \frac{1}{2} + \frac{1}{2}t_2 \text{ and } 1 > s_3 \geq \frac{1}{2} + \frac{1}{2}t_3, \\ \left(\min\left\{1, \frac{s_1 t_1}{2s_1 - t_1}\right\}, 0, 0\right), & \text{if } s_1 > \frac{1}{2}t_1, 1 > s_2 \geq \frac{1}{2} + \frac{1}{2}t_2 \text{ and } 1 > s_3 \geq \frac{1}{2} + \frac{1}{2}t_3, \\ \left(1, \frac{2t_2 - s_2 t_2 - s_2}{1 - 2s_2 + t_2}, \frac{2t_3 - s_3 t_3 - s_3}{1 - 2s_3 + t_3}\right), & \text{if } 0 < s_1 \leq \frac{1}{2}t_1, s_2 < \frac{1}{2} + \frac{1}{2}t_2 \text{ and } s_3 < \frac{1}{2} + \frac{1}{2}t_3, \\ \left(1, 0, \frac{2t_3 - s_3 t_3 - s_3}{1 - 2s_3 + t_3}\right), & \text{if } 0 < s_1 \leq \frac{1}{2}t_1, 1 > s_2 \geq \frac{1}{2} + \frac{1}{2}t_2 \text{ and } s_3 < \frac{1}{2} + \frac{1}{2}t_3, \\ \left(1, \frac{2t_2 - s_2 t_2 - s_2}{1 - 2s_2 + t_2}, 0\right), & \text{if } 0 < s_1 \leq \frac{1}{2}t_1, s_2 < \frac{1}{2} + \frac{1}{2}t_2 \text{ and } 1 > s_3 \geq \frac{1}{2} + \frac{1}{2}t_3, \\ (1, 0, 0), & \text{if } 0 < s_1 \leq \frac{1}{2}t_1, 1 > s_2 \geq \frac{1}{2} + \frac{1}{2}t_2 \text{ and } 1 > s_3 \geq \frac{1}{2} + \frac{1}{2}t_3, \\ (0, 1, 1) & \text{if } s_1 = t_1 = 0, s_2 = t_2 = 1 \text{ and } s_3 = t_3 = 1. \end{cases}$$

(4)

$$I_{O_{min}}(s, t) = \begin{cases} (1, 0, 0), & \text{if } s_1 \leq t_1^2, s_2 \geq 1 - (1 - t_2)^2 \text{ and } s_3 \geq 1 - (1 - t_3)^2, \\ (1, 1 - (1 - t_2)^2, 0), & \text{if } s_1 \leq t_1^2, s_2 < 1 - (1 - t_2)^2 \text{ and } s_3 \geq 1 - (1 - t_3)^2, \\ (1, 0, 1 - (1 - t_3)^2), & \text{if } s_1 \leq t_1^2, s_2 \geq 1 - (1 - t_2)^2 \text{ and } s_3 < 1 - (1 - t_3)^2, \\ (1, 1 - (1 - t_2)^2, 1 - (1 - t_3)^2), & \text{if } s_1 \leq t_1^2, s_2 < 1 - (1 - t_2)^2 \text{ and } s_3 < 1 - (1 - t_3)^2, \\ (t_1^2, 0, 0), & \text{if } s_1 > t_1^2, s_2 \geq 1 - (1 - t_2)^2 \text{ and } s_3 \geq 1 - (1 - t_3)^2, \\ (t_1^2, 1 - (1 - t_2)^2, 0), & \text{if } s_1 > t_1^2, s_2 < 1 - (1 - t_2)^2 \text{ and } s_3 \geq 1 - (1 - t_3)^2, \\ (t_1^2, 0, 1 - (1 - t_3)^2), & \text{if } s_1 > t_1^2, s_2 \geq 1 - (1 - t_2)^2 \text{ and } s_3 < 1 - (1 - t_3)^2, \\ (t_1^2, 1 - (1 - t_2)^2, 1 - (1 - t_3)^2), & \text{if } s_1 > t_1^2, s_2 < 1 - (1 - t_2)^2 \text{ and } s_3 < 1 - (1 - t_3)^2. \end{cases}$$

Theorem 4.1 Suppose O is the neutrosophic overlap function on D^* , $\forall s, t, h \in D^*$:

$$I_O(s, t) = \sup\{h \mid h \in D^*, O(s, h) \leq_1 t\}.$$

Thus, I_O is a neutrosophic implication.

Proof. $\forall s, u, t, h, v \in D^*$ with $s = (s_1, s_2, s_3)$, t, h, u and v are analogous to s .

(a) Let $s \leq_1 u$ and since O is non-decreasing, $\{h \mid h \in D^*, O(s, h) \leq_1 t\} \supseteq \{h \mid h \in D^*, O(u, h) \leq_1 t\}$ then $\sup\{h \mid h \in D^*, h \leq_1 I_O(s, t)\} \geq_1 \sup\{h \mid h \in D^*, h \leq_1 I_O(u, t)\}$. Thus $I_O(s, t) \geq_1 I_O(u, t)$. In other words, the first variable of I_O regarding \leq_1 is non-increasing.

(b) Let $t \leq_1 v$ and since O is non-decreasing, $\{h \mid h \in D^*, O(s, h) \leq_1 t\} \subseteq \{h \mid h \in D^*, O(s, h) \leq_1 v\}$ then $\sup\{h \mid h \in D^*, h \leq_1 I_O(s, t)\} \leq_1 \sup\{h \mid h \in D^*, h \leq_1 I_O(s, v)\}$. Thus $I_O(s, t) \leq_1 I_O(s, v)$. In other words, the second variable of I_O regarding \leq_1 is non-decreasing.

(c) $I_O(0_{D^*}, 0_{D^*}) = \sup\{h \mid h \in D^*, O(0_{D^*}, h) \leq_1 0_{D^*}\} = 1_{D^*};$

(d) $I_O(1_{D^*}, 1_{D^*}) = \sup\{h \mid h \in D^*, O(1_{D^*}, h) \leq_1 1_{D^*}\} = \sup\{h \mid h \in D^*, h \leq_1 1_{D^*}\} = 1_{D^*};$

(e) $I_O(1_{D^*}, 0_{D^*}) = \sup\{h \mid h \in D^*, O(1_{D^*}, h) \leq_1 0_{D^*}\} = \sup\{h \mid h \in D^*, h \leq_1 0_{D^*}\} = 0_{D^*}.$

The NRI has the following important properties.

Theorem 4.2 Assume that I_O is NRI, O is neutrosophic overlap function on D^* . $\forall s, t, h \in D^*$, the follows properties are valid,

- (1) $I_O(0_{D^*}, t) = 1_{D^*};$
- (2) $I_O(s, 1_{D^*}) = 1_{D^*};$
- (3) $I_O(s, s) = 1_{D^*};$
- (4) $I_O(1_{D^*}, t) = t;$
- (5) $I_O(s, t) \geq_1 t;$
- (6) $I_O(s, t) = 1_{D^*}$ iff $s \leq_1 t;$
- (7) $s \leq_1 I_O(t, h)$ iff $t \leq_1 I_O(s, h);$
- (8) $s \leq_1 I_O(t, I_O(s, t)).$

Proof. $\forall s, t, h \in D^*$ in which $s = (s_1, s_2, s_3)$, t and h are analogous to s .

(1) $I_O(0_{D^*}, t) = 1_{D^*}$ is the same thing as $I_O(0_{D^*}, t) = \sup\{h \mid h \in D^*, O(0_{D^*}, h) \leq_1 t\} = 1_{D^*}$, then $h = 1_{D^*}$ for $O(0_{D^*}, h) \leq_1 t$. Then this formula is proved.

The proofs of (2)–(4) is similar to that of (1).

(5) I is non-increasing for the first variable component (in relation to the order relation \leq_1), then $I_O(s, t) \geq_1 I_O(1_{D^*}, t) = t$.

(6) $I_O(s, t) = 1_{D^*}$ iff $s \leq_1 t$. Let $s \leq_1 t$, $O(1_{D^*}, s) \leq_1 t$, then $I_O(s, t) = 1_{D^*}$. In contrast, let $I_O(s, t) = 1_{D^*}$, thus $O(1_{D^*}, s) \leq_1 t$, hence $s \leq_1 t$.

(7) $s \leq_1 I_O(t, h)$ iff $t \leq_1 I_O(s, h)$. Since $s \leq_1 I_O(t, h)$, $O(t, s) \leq_1 h$. Thus, $t \leq_1 I_O(s, h)$. Likewise, $s \leq_1 I_O(t, h)$ can be proved from $t \leq_1 I_O(s, h)$.

(8) $s \leq_1 I_O(t, I_O(s, t))$. Since $O(t, s) \leq_1 O(s, t)$, then $s \leq_1 I_O(t, I_O(s, t))$.

Example 4.2 These concrete cases about NRI deduced from neutrosophic overlap function as shown in **Example 4.1** are given. And it is readily proved that NRI deduced from neutrosophic overlap functions satisfy the properties characterised by **Theorem 4.2**.

Furthermore, for the non-representable neutrosophic overlap function, using the neutrosophic overlap function got from **Theorem 3.4** as an example, $\forall s, t \in D^*$,

$$I_O(s, t) = \begin{cases} 1_{D^*}, & \text{if } s = 0_{D^*} \text{ or } t = 0_{D^*}; \text{ or } s = t = 1_{D^*}, \\ (1, [0, 1], \max\{\frac{t_3}{s_2}, \frac{t_3}{s_3}\}), & \text{if } s_1 \leq t_1, \\ (\frac{t_1}{s_1}, [0, 1], \max\{\frac{t_3}{s_2}, \frac{t_3}{s_3}\}), & \text{if } s_1 > t_1. \end{cases}$$

Then it follows that $Io(s, t)$ is a neutrosophic implication, while the properties from **Theorem 4.2** is satisfied.

5. Conclusions

As an important part of NS theory, neutrosophic logic plays a significant part in it. Neutrosophic overlap function, neutrosophic grouping function and neutrosophic implication which are crucial neutrosophic logic operators. For the first kind of inclusion relationship, the definitions of neutrosophic overlap function (neutrosophic grouping function) on $(D^*; \leq_1)$ are defined and related examples are given. At the same time, new definitions of representable and non-representable neutrosophic overlap function are proposed. In the next place, based on the close relationship between overlap function and triangular norm, a new description of neutrosophic negation is offered through analogy research, then the dual relationship between neutrosophic overlap function and neutrosophic grouping function on neutrosophic negation is described. Moreover, we show that definition of neutrosophic implication is given based on $(D^*; \leq_1)$ and the basic properties of NRI are studied. Finally, the result that NRI induced by neutrosophic overlap function must be neutrosophic implication is proved. Based on these results and some new results [33-44], we consider applying them to generalized neutrosophic overlap function and neutrosophic inference systems of the future.

References

1. Smarandache, F. *Neutrosophy: Neutrosophic Probability, Set, and Logic: Analytic Synthesis and Synthetic Analysis*; American Research Press: Santa Fe, NM, USA, 1998; pp. 25-38.
2. Wang, H.B.; Smarandache, F.; Zhang, Y.Q.; Sunderraman, R. Single valued neutrosophic sets. *Multispace Multistruct. Neutrosophic Transdiscipl.* **2010**, *4*, 410-413.
3. Tan, R.P.; Zhang, W.D. Decision-making method based on new entropy and refined single-valued neutrosophic sets and its application in typhoon disaster assessment. *Applied Intelligence*, **2021**, *51*, 283-307.
4. Stanujkic, D.; Karabasevic, D.; Popovic, G.; et al. Multiple-criteria decision-making based on the use of single-valued neutrosophic sets and similarity measures. *ECON COMPUT ECON CYB.* **2021**, *55*, 5-22.
5. Liu, P.D. The aggregation operators based on archimedean t-conorm and t-norm for single-valued neutrosophic numbers and their application to decision making. *Int. J. Fuzzy Syst.* **2016**, *18*, 849-863.
6. Ye, J. Multicriteria decision-making method using the correlation coefficient under single-valued neutrosophic environment. *Int. J. Gen. Syst.* **2013**, *42*, 386-394.
7. Hu, Q.Q.; Zhang, X.H. New similarity measures of single-valued neutrosophic multisets based on the decomposition theorem and its application in medical diagnosis. *Symmetry* **2018**, *10*, 466. DOI: 10.3390/sym10100466.
8. Wang, J.Q.; Zhang, X.H. Two types of single-valued neutrosophic covering rough sets and an application to decision making. *Symmetry* **2018**, *10*, 710. DOI: 10.3390/sym10120710.
9. Wen, X.F.; Zhang, X.H.; Lei, T. Intuitionistic Fuzzy (IF) Overlap Functions and IF-Rough Sets with Applications. *Symmetry* **2021**, *13*, 1494. DOI: 10.3390/SYM13081494.
10. Bustince, H.; Fernandez, J.; Mesiar, R.; Montero, J.; Orduna, R. Overlap functions. *Nonlinear Anal.* **2010**, *72*, 1488-1499.
11. Bustince, H.; Pagola, M.; Mesiar, R.; et al. Grouping, overlap, and generalized bientropic functions for fuzzy modeling of pairwise comparisons. *IEEE Transactions on Fuzzy Systems.* **2011**, *20*, 405-415.
12. Dimuro, G.P.; Bedregal, B.; Bustince, H.; et al. On additive generators of overlap functions. *Fuzzy Sets and Systems.* **2016**, *287*, 76-96.
13. Bedregal, B.; Dimuro, G.P.; Bustince, H.; et al. New results on overlap and grouping functions. *Information Sciences.* **2013**, *249*: 148-170.
14. Qiao, J.S. On binary relations induced from overlap and grouping functions. *Int. J. Approx. Reason.* **2019**, *106*, 155-171.
15. Qiao, J.S. On (IO, O)-fuzzy rough sets based on overlap functions. *Int. J. Approx. Reason.* **2021**, *132*, 26-48.
16. Qiao, J.S. Overlap and grouping functions on complete lattices. *Inf. Sci.* **2021**, *542*, 406-424.

17. Bustince, H.; Barrenechea, E.; Pagola, M. Image thresholding using restricted equivalence functions and maximizing the measures of similarity. *Fuzzy Sets and Systems*. **2007**, *158*, 496-516.
18. Hu, Q.Q.; Zhang, X.H. Neutrosophic triangular norms and their derived residuated lattices. *Symmetry* **2019**, *11*, 817. DOI: 10.3390/sym11060817.
19. Driankov, D.; Hellendoorn, H. Chaining of fuzzy if-then rules in Mamdani-controllers. *Proceedings of 1995 IEEE International Conference on Fuzzy Systems*. **1995**, *1*, 103-108.
20. Baczyński, M.; Jayaram, B. (S, N)-and R-implications: a state-of-the-art survey. *Fuzzy Sets and Systems*. **2008**, *159*, 1836-1859.
21. Dimuro, G.P.; Bedregal, B. On residual implications derived from overlap functions. *Information Sciences*. **2015**, *312*, 78-88.
22. De Miguel, L.; Gómez, D.; Rodríguez, J.T.; et al. General overlap functions. *Fuzzy Sets and Systems*, **2019**, *372*, 81-96.
23. Baczyński, M.; Jayaram, B. *Fuzzy implications*. Studies in Fuzziness and Soft Computing, Springer, Berlin Heidelberg, 2008; pp. 151-160.
24. Zhang, F.X.; Liu, H.W. On a new class of implications: (g, u)-implications and the distributive equations. *International journal of approximate reasoning*. **2013**, *54*, 1049-1065.
25. Klement, E.P.; Mesiar, R.; Pap, E. Triangular norms. Position paper I: Basic analytical and algebraic properties. *Fuzzy Sets Syst.* **2004**, *143*, 5-26.
26. Vemuri, N.R. Mutually exchangeable fuzzy implications. *Information Sciences*. **2015**, *317*, 1-24.
27. Baczyński, M.; Jayaram, B. (S, N)-and R-implications: a state-of-the-art survey. *Fuzzy Sets and Systems*. **2008**, *159*, 1836-1859.
28. Baczyński, M.; Jayaram, B. On the characterizations of (S, N)-implications. *Fuzzy sets and systems*. **2007**, *158*, 1713-1727.
29. Bustince, H.; Fernández, J.; Mesiar, R.; et al. Overlap index, overlap functions and migrativity. Proceedings of the Joint 2009 International Fuzzy Systems Association World Congress and 2009 European Society of Fuzzy Logic and Technology Conference, Lisbon, Portugal, July 20-24, 2009.
30. Bustince, H.; Pagola, M.; Mesiar, R.; et al. Grouping, overlap, and generalized bientropic functions for fuzzy modeling of pairwise comparisons. *IEEE Transactions on Fuzzy Systems*, **2011**, *20*, 405-415.
31. Atanassov, K. Intuitionistic fuzzy sets. *Fuzzy Sets Syst.* **1986**, *20*, 87-96.
32. Deschrijver, G. Implication functions in interval-valued fuzzy set theory. *Stud. Fuzziness Soft Comput.* **2013**, *300*, 73-99.
33. Liang, R.; Zhang, X.H. Interval-valued pseudo overlap functions and application. *Axioms* **2022**, *11*, 216. DOI: 10.3390/axioms11050216.
34. Jing, M.; Zhang, X.H. Pseudo-quasi overlap functions and related fuzzy inference methods. *Axioms* **2023**, *12*, 217. DOI: 10.3390/axioms12020217.
35. Liang, R.; Zhang, X.H. Pseudo general overlap functions and weak inflationary pseudo BL-algebras. *Mathematics*. **2022**, *10*, 1396. DOI: 10.3390/math10091396.
36. Zhang, X.H.; Liang, R.; Bustince, H.; Bedregal, B.; Fernandez, J.; Li, M.Y.; Ou, Q.Q. Pseudo overlap functions, fuzzy implications and pseudo grouping functions with applications. *Axioms* **2022**, *11*, 593. DOI: 10.3390/axioms11110593.
37. Zhang, X.H.; Liang, R. Interval-valued general residuated lattice-ordered groupoids and expanded triangle algebras. *Axioms* **2023**, *12*, 42. DOI: 10.3390/axioms12010042.
38. Zhang, X.H.; Sheng, N.; Borzooei, R.A. Partial residuated implications induced by partial triangular norms and partial residuated lattices. *Axioms* **2023**, *12*, 63. DOI: 10.3390/axioms12010063.
39. Zhang, X.H.; Shang, J.Y.; Wang, J.Q. Multi-granulation fuzzy rough sets based on overlap functions with a new approach to MAGDM. *Information Sciences*. **2023**, *622*, 536-559.
40. Wang, J.Q.; Zhang, X.H.; Hu, Q.Q. Three-way fuzzy sets and their applications (II). *Axioms* **2022**, *11*, 532. DOI: 10.3390/axioms11100532.
41. Hu, Q.Q.; Zhang, X.H. Three-way fuzzy sets and their applications (III). *Axioms* **2023**, *12*, 57. DOI: 10.3390/axioms12010057.
42. Zhang, X.H.; Liang, R.; Bedregal, B. Weak inflationary BL-algebras and filters of inflationary (pseudo) general residuated lattices. *Mathematics*. **2022**, *10*, 3394. DOI: 10.3390/math10183394.

43. Zhang, X.H.; Li, M.Y.; Liu, H. Overlap functions-based fuzzy mathematical morphological operators and their applications in image edge extraction. *Fractal Fract.* 2023, 7, 465. DOI: 10.3390/fractalfract7060465.
44. Zhang, X.H.; Jiang, H.J.; Wang, J.Q. New classifier ensemble and fuzzy community detection methods using POP Choquet-like integrals. *Fractal Fract.* 2023, 7, 588. DOI: 10.3390/fractalfract7080588.

Received: March 19, 2023. Accepted: July 20, 2023