



## Original Article

Smarandache curves in Euclidean 4- space  $E^4$ 

Mervat Elzawy

Mathematics Department, Faculty of Science, Tanta University, Tanta, Egypt



## ARTICLE INFO

## Article history:

Received 1 January 2017

Revised 15 February 2017

Accepted 5 March 2017

Available online 17 March 2017

## 2010 MSC:

14H45

14H50

53A04

## Keywords:

Euclidean 4- space

Smarandache curves

Frenet frame

Parallel transport frame

## ABSTRACT

The purpose of this paper is to study Smarandache curves in the 4-dimensional Euclidean space  $E^4$ , and to obtain the Frenet–Serret and Bishop invariants for the Smarandache curves in  $E^4$ . The first, the second and the third curvatures of Smarandache curves are calculated. These values depending upon the first, the second and the third curvature of the given curve.

© 2017 Egyptian Mathematical Society. Production and hosting by Elsevier B.V.

This is an open access article under the CC BY-NC-ND license.

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

## 1. Introduction

It is well known that, if a curve is differentiable in an open interval, at each point, a set of mutually orthogonal unit vectors can be constructed, and these vectors are called Frenet frame or moving frame vectors. The rates of these frame vectors along the curve define curvatures of the curve. The set, whose elements are frame vectors and curvatures of a curve is called Frenet apparatus of the curves. In recent years, the theory of degenerate submanifolds has been treated by researchers and some classical differential geometry topics have been extended to Minkowski space [1–3] and Galilean space [4]. For instance in [1,2] the authors extended and studied Smarandache curves in Minkowski space-time. A regular curve in Euclidean space  $E^4$ , whose position vectors is composed by Frenet frame vectors on another regular curve is called Smarandache curve. Special Smarandache curves in three dimensional Euclidean space studied in [5].

The Bishop frame [6] or parallel transport frame is an alternative approach to defining a moving frame that is well defined even when the curve has vanishing second derivative. We can parallel transport an orthonormal frame along a curve simply by parallel transporting each component of the frame in Euclidean 4-space. The parallel transport frame is based on the observation that while  $T(s)$  for the given curve model is unique, we may choose any convenient arbitrary basis which consists of relatively paral-

lel vector fields  $\{M_1(s), M_2(s), M_3(s)\}$  of the frame, such that they are perpendicular to  $T(s)$  at each point [7,8]. The parallel transport frame in four dimensional Euclidean space is studied in [9]. Smarandache curves were studied from deferent researchers in three dimensional Euclidean space [10–14]. Smarandache curves in 4-dimensional Galilean space are presented in [15]. In this paper we study Smarandache curves in 4-dimensional space according to the Frenet frame and parallel transport frame.

## 2. Preliminaries

Let  $\alpha: R \rightarrow E^4$  be an arbitrary curve in the Euclidean space  $E^4$ . Let  $\vec{a} = (a_1, a_2, a_3, a_4)$ ,  $\vec{b} = (b_1, b_2, b_3, b_4)$  and  $\vec{c} = (c_1, c_2, c_3, c_4)$  be three vectors in  $E^4$ , equipped with the standard inner product given by  $\langle \vec{a}, \vec{b} \rangle = a_1b_1 + a_2b_2 + a_3b_3 + a_4b_4$ . The norm of a vector  $a \in E^4$  is given by  $\|\vec{a}\| = \sqrt{\langle \vec{a}, \vec{a} \rangle}$ . The curve  $\alpha$  is said to be of a unit speed or parametrized by arc length function  $s$  if  $\langle \alpha', \alpha' \rangle = 1$ . The vector product of  $\vec{a}$ ,  $\vec{b}$ , and  $\vec{c}$  is defined by the determinant

$$\vec{a} \times \vec{b} \times \vec{c} = \begin{vmatrix} e_1 & e_2 & e_3 & e_4 \\ a_1 & a_2 & a_3 & a_4 \\ b_1 & b_2 & b_3 & b_4 \\ c_1 & c_2 & c_3 & c_4 \end{vmatrix}$$

where  $e_1 \times e_2 \times e_3 = e_4, e_2 \times e_3 \times e_4 = e_1, e_3 \times e_4 \times e_1 = e_2, e_4 \times e_1 \times e_2 = e_3, e_3 \times e_2 \times e_1 = e_4$ .

E-mail address: [mervatelzawy@science.tanta.edu.eg](mailto:mervatelzawy@science.tanta.edu.eg)

The vectors  $t(s), n(s), b_1(s), b_2(s)$  are the moving Frenet frame along the unit speed curve  $\alpha$ . Then the Frenet formulas are given by

$$\begin{bmatrix} t' \\ n' \\ b_1' \\ b_2' \end{bmatrix} = \begin{bmatrix} 0 & k_1 & 0 & 0 \\ -k_1 & 0 & k_2 & 0 \\ 0 & -k_2 & 0 & k_3 \\ 0 & 0 & -k_3 & 0 \end{bmatrix} \begin{bmatrix} t \\ n \\ b_1 \\ b_2 \end{bmatrix}$$

$t, n, b_1$ , and  $b_2$  are called, respectively, the tangent, the principal normal, the first binormal and the second binormal vector fields of the curves. The functions  $k_1(s), k_2(s)$  and  $k_3(s)$  are called respectively, the first, the second and the third curvature of the curve  $\alpha$ . The curve is called  $W$ -curve if it has constant curvatures  $k_1, k_2$  and  $k_3$ .

Let  $\alpha = \alpha(t)$  be an arbitrary curve in  $E^4$ . The Frenet apparatus of the curve  $\alpha$  can be calculated by the following equations.

$$t = \frac{\alpha'}{\|\alpha'\|}$$

$$n = \frac{\|\alpha'\|^2 \alpha'' - \langle \alpha', \alpha'' \rangle \alpha'}{\|\|\alpha'\|^2 \alpha'' - \langle \alpha', \alpha'' \rangle \alpha'\|}$$

$$b_1 = \eta b_2 \times t \times n$$

$$b_2 = \eta \frac{t \times n \times \alpha'''}{\|t \times n \times \alpha'''\|}$$

$$k_1 = \frac{\|\|\alpha'\|^2 \alpha'' - \langle \alpha', \alpha'' \rangle \alpha'\|}{\|\alpha'\|^4}$$

$$k_2 = \frac{\|t \times n \times \alpha'''\| \|\alpha'\|}{\|\|\alpha'\|^2 \alpha'' - \langle \alpha', \alpha'' \rangle \alpha'\|}$$

$$k_3 = \frac{\langle \alpha^{(IV)}, b_2 \rangle}{\|t \times n \times \alpha'''\| \|\alpha'\|}$$

where  $\eta$  is taken  $\pm 1$  such that determinant of matrix  $[t, n, b_1, b_2]$  is equal to one.

The Bishop frame or parallel transport frame is an alternative approach to define a moving frame that is well defined even when the curve has vanishing second derivative [9]. One can express parallel transport of an orthonormal frame along a curve simply by parallel transporting each component of the frame. The tangent vector and the convenient arbitrary basis for the remainder of the frame are used. The parallel transport equations in  $E^4$  can be expressed as

$$\begin{bmatrix} T' \\ M_1' \\ M_2' \\ M_3' \end{bmatrix} = \begin{bmatrix} 0 & K_1 & K_2 & K_3 \\ -K_1 & 0 & 0 & 0 \\ -K_2 & 0 & 0 & 0 \\ -K_3 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} T \\ M_1 \\ M_2 \\ M_3 \end{bmatrix}$$

where  $K_1, K_2$ , and  $K_3$  are the principal curvature functions according to parallel transport frame of the curve  $\alpha$ . The set  $\{T, M_1, M_2, M_3\}$  is called the parallel transport frame of  $\alpha$  [6,14].

The expressions of the principal curvatures are given as follows:

$$K_1 = k_1 \cos \theta \cos \psi,$$

$$K_2 = k_1 (-\cos \phi \sin \psi + \sin \phi \sin \theta \cos \psi),$$

$$K_3 = k_1 (\sin \phi \sin \psi + \cos \phi \sin \theta \cos \psi) \text{ and}$$

$$k_1 = \sqrt{K_1^2 + K_2^2 + K_3^2}, \quad k_2 = -\psi' + \phi' \sin \theta, \quad k_3 = \frac{\theta'}{\sin \psi},$$

$$\phi' \cos \theta + \theta' \cot \psi = 0$$

where

$$\theta' = \frac{k_3}{\sqrt{k_1^2 + k_2^2}}, \quad \psi' = -(k_2 + k_3 \frac{\sqrt{k_3^2 - (\theta')^2}}{\sqrt{k_1^2 + k_2^2}}), \quad \phi' = \frac{\sqrt{k_3^2 - (\theta')^2}}{\cos \theta}$$

Note that  $k_1, k_2, k_3$  are the principal curvature functions according to Frenet frame and  $K_1, K_2, K_3$  are the principal curvature functions according to the parallel transport frame of the curve  $\alpha$ .

### 3. Main results

#### 3.1. $tb_1$ Smarandache curves in $E^4$ according to the Frenet frame.

In this subsection we define  $tb_1$  Smarandache curves and obtain their Frenet apparatus.

**Definition 1.** A regular curve in  $E^4$ , whose position vector is obtained by Frenet frame vectors on another regular curve, is called Smarandache curve.

**Definition 2.** Let  $\alpha = \alpha(s)$  be a unit-speed curve with constant and nonzero curvatures  $k_1, k_2, k_3$  and  $\{t, n, b_1, b_2\}$  be moving frame on it,  $tb_1$  Smarandache curves are defined by  $\beta(s_\beta) = \frac{1}{\sqrt{2}}(t(s) + b_1(s))$ .

**Theorem 1.** Let  $\alpha(s)$  be a unit speed curve with constant non zero curvatures  $k_1, k_2, k_3$  and  $\beta(s_\beta)$  be  $tb_1$  Smarandache curves in  $E^4$  defined by the frame vectors of  $\alpha(s)$ . Then the Frenet apparatus of  $\beta(\{t_\beta, n_\beta, b_{1\beta}, b_{2\beta}, k_{1\beta}, k_{2\beta}, k_{3\beta}\})$  can be formed by Frenet apparatus of  $\alpha(\{t, n, b_1, b_2, k_1, k_2, k_3\})$ .

**Proof.** Let  $\beta = \beta(s_\beta)$  be  $tb_1$  Smarandache curve of the curve  $\alpha$ . Then

$$\text{From Definition (2) we have } \beta(s_\beta) = \frac{1}{\sqrt{2}}(t(s) + b_1(s))$$

By differentiating  $\beta(s_\beta)$  with respect to  $s$  we obtain

$$\frac{d\beta(s_\beta)}{ds} = \frac{d\beta(s_\beta)}{ds_\beta} \frac{ds_\beta}{ds} = \frac{1}{\sqrt{2}}((k_1 - k_2)n + k_3 b_2)$$

The tangent vector of the curve  $\beta$  is given by

$$t_\beta = A_1 n + A_2 b_2 \tag{3.1}$$

where  $\frac{ds_\beta}{ds} = \frac{\sqrt{(k_1 - k_2)^2 + k_3^2}}{\sqrt{2}}, A_1 = \frac{(k_1 - k_2)}{\sqrt{(k_1 - k_2)^2 + k_3^2}}$  and  $A_2 = \frac{k_3}{\sqrt{(k_1 - k_2)^2 + k_3^2}}$

Again differentiating the tangent vector of  $\beta$  with respect to  $s_\beta$  we can obtain  $\beta''$  as follows

$$\beta'' = \frac{\sqrt{2}[-k_1(k_1 - k_2)t + (k_1 k_2 - k_2^2 - k_3^2)b_1]}{(k_1 - k_2)^2 + k_3^2}$$

The principal normal of the curve  $\beta$  is

$$n_\beta = A_3 t + A_4 b_1 \tag{3.2}$$

where  $A_3 = \frac{-k_1(k_1 - k_2)}{\sqrt{k_1^2(k_1 - k_2)^2 + (k_1 k_2 - k_2^2 - k_3^2)^2}}$  and  $A_4 = \frac{k_1 k_2 - k_2^2 - k_3^2}{\sqrt{k_1^2(k_1 - k_2)^2 + (k_1 k_2 - k_2^2 - k_3^2)^2}}$

$$\beta''' = A_5 n + A_6 b_2$$

where  $A_5 = \frac{2(-k_1^2(k_1 - k_2) - k_2(k_1 k_2 - k_2^2 - k_3^2))}{((k_1 - k_2)^2 + k_3^2)^{\frac{3}{2}}}$  and  $A_6 = \frac{k_3(k_1 k_2 - k_2^2 - k_3^2)}{((k_1 - k_2)^2 + k_3^2)^{\frac{3}{2}}}$

The second binormal vector of the curve  $\beta$  is given easily as follows

$$b_{2\beta} = \frac{(k_1 k_2 - k_2^2 - k_3^2)t + k_1(k_1 - k_2)b_1}{\sqrt{(k_1 k_2 - k_2^2 - k_3^2)^2 + k_1^2(k_1 - k_2)^2}} \tag{3.3}$$

The first binormal vector of the curve  $\beta$  is

$$b_{1\beta} = \frac{-k_3 n + (k_1 - k_2)b_2}{\sqrt{k_3^2 + (k_1 - k_2)^2}} \tag{3.4}$$

The first, second and third curvature of the of the curve  $\beta$  are

$$k_{1\beta} = \frac{2[(k_1k_2 - k_2^2 - k_3^2)^2 + k_1^2(k_1 - k_2)^2]}{(k_3^2 + (k_1 - k_2)^2)^2} \tag{3.5}$$

$$k_{2\beta} = \frac{\sqrt{2}k_3[k_1(k_1k_2 - k_2^2 - k_3^2) + k_1^2(k_1 - k_2)]}{(k_3^2 + (k_1 - k_2)^2)\sqrt{(k_1k_2 - k_2^2 - k_3^2)^2 + k_1^2(k_1 - k_2)^2}} \tag{3.6}$$

$$k_{3\beta} = \frac{\sqrt{2}(-k_1A_4A_5 - k_2A_3A_5 + k_3A_3A_6)}{\sqrt{k_3^2 + (k_1 - k_2)^2}\sqrt{(k_1k_2 - k_2^2 - k_3^2)^2 + k_1^2(k_1 - k_2)^2}} \tag{3.7}$$

This completes the proof.  $\square$

3.2.  $TM_1$  Smarandache curves in  $E^4$  according to the parallel transport frame.

In this subsection we define  $TM_1$  Smarandache curves and obtain their parallel transport frame and the principal curvatures.

**Definition 3.** Let  $\alpha = \alpha(s)$  be a unit speed curve in  $E^4$  and  $\{T_\alpha, M_{1\alpha}, M_{2\alpha}, M_{3\alpha}\}$  be its moving parallel transport frame.  $TM_1$  Smarandache curves is defined by  $\beta(s_\beta) = \frac{1}{\sqrt{2}}(T_\alpha + M_{1\alpha})$ .

**Theorem 2.** Let  $\alpha = \alpha(s)$  be a unit speed curve with constant principal curvatures  $K_{1\alpha}, K_{2\alpha}, K_{3\alpha}$  and  $\beta(s_\beta)$  be  $TM_1$  Smarandache curves in  $E^4$  defined by the parallel transport frame vectors of  $\alpha = \alpha(s)$ . Then the parallel transport frame of  $\beta$  can be formed by the parallel transport frame of  $\alpha$  and the principle curvatures of  $\beta$  ( $K_{1\beta}, K_{2\beta}, K_{3\beta}$ ) can be obtained by the principal curvatures of  $\alpha$ .

**Proof.** To investigate the parallel transport frame of  $TM_1$  Smarandache curve according to  $\alpha(s)$  differentiating  $\beta(s_\beta) = \frac{1}{\sqrt{2}}(T_\alpha + M_{1\alpha})$  with respect to  $s$

$$T_\beta \frac{ds_\beta}{ds} = \frac{1}{\sqrt{2}}(-K_{1\alpha}T_\alpha + K_{1\alpha}M_{1\alpha} + K_{2\alpha}M_{2\alpha} + K_{3\alpha}M_{3\alpha})$$

The tangent vector of the curve  $\beta$  can be written as follows

$$T_\beta = \frac{-K_{1\alpha}T_\alpha + K_{1\alpha}M_{1\alpha} + K_{2\alpha}M_{2\alpha} + K_{3\alpha}M_{3\alpha}}{\sqrt{2K_{1\alpha}^2 + K_{2\alpha}^2 + K_{3\alpha}^2}} \tag{3.8}$$

where  $\frac{ds_\beta}{ds} = \frac{1}{\sqrt{2}}\sqrt{2K_{1\alpha}^2 + K_{2\alpha}^2 + K_{3\alpha}^2}$

Differentiating (3.8) with respect to  $s$

$$T'_\beta = \frac{dT_\beta}{ds_\beta} = \lambda_0T_\alpha + \lambda_1M_{1\alpha} + \lambda_2M_{2\alpha} + \lambda_3M_{3\alpha}$$

where  $\lambda_0 = \frac{-\sqrt{2}(K_{1\alpha}^2 + K_{2\alpha}^2 + K_{3\alpha}^2)}{(2K_{1\alpha}^2 + K_{2\alpha}^2 + K_{3\alpha}^2)}$ ,  $\lambda_1 = \frac{-\sqrt{2}K_{1\alpha}^2}{(2K_{1\alpha}^2 + K_{2\alpha}^2 + K_{3\alpha}^2)}$

$\lambda_2 = \frac{-\sqrt{2}K_{1\alpha}K_{2\alpha}}{(2K_{1\alpha}^2 + K_{2\alpha}^2 + K_{3\alpha}^2)}$ ,  $\lambda_3 = \frac{-\sqrt{2}K_{1\alpha}K_{3\alpha}}{(2K_{1\alpha}^2 + K_{2\alpha}^2 + K_{3\alpha}^2)}$

The first curvature of the curve  $\beta$  according to Frenet frame is

$$k_{1\beta} = \sqrt{\lambda_0^2 + \lambda_1^2 + \lambda_2^2 + \lambda_3^2} = \frac{\sqrt{2}\sqrt{K_{1\alpha}^2 + K_{2\alpha}^2 + K_{3\alpha}^2}}{\sqrt{2K_{1\alpha}^2 + K_{2\alpha}^2 + K_{3\alpha}^2}} \tag{3.9}$$

The principal normal of the curve  $\beta$  is given by the following formula

$$n_\beta = \frac{\lambda_0T_\alpha + \lambda_1M_{1\alpha} + \lambda_2M_{2\alpha} + \lambda_3M_{3\alpha}}{\sqrt{\lambda_0^2 + \lambda_1^2 + \lambda_2^2 + \lambda_3^2}} \tag{3.10}$$

The third derivative of the curve  $\beta$  reads

$$\beta''' = (\lambda_0T'_\alpha + \lambda_1M'_{1\alpha} + \lambda_2M'_{2\alpha} + \lambda_3M'_{3\alpha}) \frac{\sqrt{2}}{\sqrt{2K_{1\alpha}^2 + K_{2\alpha}^2 + K_{3\alpha}^2}}$$

$$\beta''' = \frac{\sqrt{2}[(-\lambda_1K_{1\alpha} - \lambda_2K_{2\alpha} - \lambda_3K_{3\alpha})T_\alpha + \lambda_0K_{1\alpha}M_{1\alpha} + \lambda_0K_{2\alpha}M_{2\alpha} + \lambda_0K_{3\alpha}M_{3\alpha}]}{\sqrt{2K_{1\alpha}^2 + K_{2\alpha}^2 + K_{3\alpha}^2}} \tag{3.11}$$

$T_\beta \times n_\beta \times \beta''' = C_1M_{1\alpha} + C_2M_{2\alpha} + C_3M_{3\alpha}$  where

$$C_1 = \frac{\sqrt{2}[\lambda_0\lambda_3K_{1\alpha}K_{2\alpha} - \lambda_0\lambda_2K_{1\alpha}K_{3\alpha} - (\lambda_1K_{1\alpha} + \lambda_2K_{2\alpha} + \lambda_3K_{3\alpha})(\lambda_3K_{2\alpha} - \lambda_2K_{3\alpha})]}{\sqrt{\lambda_0^2 + \lambda_1^2 + \lambda_2^2 + \lambda_3^2}(2K_{1\alpha}^2 + K_{2\alpha}^2 + K_{3\alpha}^2)}$$

$$C_2 = \frac{\sqrt{2}[\lambda_0\lambda_3K_{1\alpha}^2 - \lambda_0\lambda_1K_{1\alpha}K_{3\alpha} - (\lambda_1K_{1\alpha} + \lambda_2K_{2\alpha} + \lambda_3K_{3\alpha})(\lambda_3K_{1\alpha} - \lambda_1K_{3\alpha})]}{\sqrt{\lambda_0^2 + \lambda_1^2 + \lambda_2^2 + \lambda_3^2}(2K_{1\alpha}^2 + K_{2\alpha}^2 + K_{3\alpha}^2)}$$

$$C_3 = \frac{\sqrt{2}[\lambda_0\lambda_2K_{1\alpha}^2 - \lambda_0\lambda_1K_{1\alpha}K_{2\alpha} - (\lambda_1K_{1\alpha} + \lambda_2K_{2\alpha} + \lambda_3K_{3\alpha})(\lambda_1K_{2\alpha} - \lambda_2K_{2\alpha})]}{\sqrt{\lambda_0^2 + \lambda_1^2 + \lambda_2^2 + \lambda_3^2}(2K_{1\alpha}^2 + K_{2\alpha}^2 + K_{3\alpha}^2)}$$

The second binormal of the curve  $\beta$  is given by the following formula

$$b_{2\beta} = \frac{C_1M_{1\alpha} + C_2M_{2\alpha} + C_3M_{3\alpha}}{\sqrt{C_1^2 + C_2^2 + C_3^2}} \tag{3.12}$$

The first binormal of the curve  $\beta$  is given by the following formula

$$b_{1\beta} = b_{2\beta} \times T_\beta \times n_\beta = \gamma_0T_\alpha + \gamma_1M_{1\alpha} + \gamma_2M_{2\alpha} + \gamma_3M_{3\alpha} \tag{3.13}$$

where the constants are given by

$$\gamma_0 = \frac{C_1\lambda_3K_{2\alpha} - C_1\lambda_2K_{3\alpha} + C_2\lambda_1K_{3\alpha} - C_2\lambda_3K_{1\alpha} + C_3\lambda_2K_{1\alpha} - C_3\lambda_1K_{2\alpha}}{\sqrt{C_1^2 + C_2^2 + C_3^2}\sqrt{\lambda_0^2 + \lambda_1^2 + \lambda_2^2 + \lambda_3^2}\sqrt{2K_{1\alpha}^2 + K_{2\alpha}^2 + K_{3\alpha}^2}}$$

$$\gamma_1 = \frac{C_2\lambda_3K_{1\alpha} + C_2\lambda_0K_{3\alpha} - C_3\lambda_2K_{1\alpha} - C_3\lambda_0K_{1\alpha}}{\sqrt{C_1^2 + C_2^2 + C_3^2}\sqrt{\lambda_0^2 + \lambda_1^2 + \lambda_2^2 + \lambda_3^2}\sqrt{2K_{1\alpha}^2 + K_{2\alpha}^2 + K_{3\alpha}^2}}$$

$$\gamma_2 = \frac{C_1\lambda_3K_{1\alpha} + C_1\lambda_0K_{3\alpha} - C_3\lambda_1K_{1\alpha} - C_3\lambda_0K_{1\alpha}}{\sqrt{C_1^2 + C_2^2 + C_3^2}\sqrt{\lambda_0^2 + \lambda_1^2 + \lambda_2^2 + \lambda_3^2}\sqrt{2K_{1\alpha}^2 + K_{2\alpha}^2 + K_{3\alpha}^2}}$$

$$\gamma_3 = \frac{C_1\lambda_2K_{1\alpha} + C_1\lambda_0K_{2\alpha} - C_2\lambda_1K_{1\alpha} - C_2\lambda_0K_{1\alpha}}{\sqrt{C_1^2 + C_2^2 + C_3^2}\sqrt{\lambda_0^2 + \lambda_1^2 + \lambda_2^2 + \lambda_3^2}\sqrt{2K_{1\alpha}^2 + K_{2\alpha}^2 + K_{3\alpha}^2}}$$

The parallel transport frame for the curve  $\beta$  has the form

$$M_{1\beta} = \left( \frac{\lambda_0 \cos \theta_\beta \cos \psi_\beta}{\sqrt{\lambda_0^2 + \lambda_1^2 + \lambda_2^2 + \lambda_3^2}} + \gamma_0 \cos \theta_\beta \sin \psi_\beta \right) T_\alpha$$

$$+ \left( \frac{\lambda_1 \cos \theta_\beta \cos \psi_\beta}{\sqrt{\lambda_0^2 + \lambda_1^2 + \lambda_2^2 + \lambda_3^2}} + \gamma_1 \cos \theta_\beta \sin \psi_\beta \right. \\ \left. - \frac{C_1 \sin \theta_\beta}{\sqrt{C_1^2 + C_2^2 + C_3^2}} \right) M_{1\alpha} + \left( \frac{\lambda_2 \cos \theta_\beta \cos \psi_\beta}{\sqrt{\lambda_0^2 + \lambda_1^2 + \lambda_2^2 + \lambda_3^2}} \right. \\ \left. + \gamma_2 \cos \theta_\beta \sin \psi_\beta - \frac{C_2 \sin \theta_\beta}{\sqrt{C_1^2 + C_2^2 + C_3^2}} \right) M_{2\alpha}$$

$$+ \left( \frac{\lambda_3 \cos \theta_\beta \cos \psi_\beta}{\sqrt{\lambda_0^2 + \lambda_1^2 + \lambda_2^2 + \lambda_3^2}} + \gamma_3 \cos \theta_\beta \sin \psi_\beta \right. \\ \left. - \frac{C_3 \sin \theta_\beta}{\sqrt{C_1^2 + C_2^2 + C_3^2}} \right) M_{3\alpha} \tag{3.14}$$

$$\begin{aligned}
 M_{2_\beta} = & \left[ \frac{\lambda_0(-\cos\phi_\beta \sin\psi_\beta + \sin\phi_\beta \sin\theta_\beta \cos\psi_\beta)}{\sqrt{\lambda_0^2 + \lambda_1^2 + \lambda_2^2 + \lambda_3^2}} \right. \\
 & + \gamma_0(\cos\phi_\beta \cos\psi_\beta + \sin\phi_\beta \sin\theta_\beta \sin\psi_\beta) \left. \right] T_\alpha \\
 & + \left[ \frac{\lambda_1(-\cos\phi_\beta \sin\psi_\beta + \sin\phi_\beta \sin\theta_\beta \cos\psi_\beta)}{\sqrt{\lambda_0^2 + \lambda_1^2 + \lambda_2^2 + \lambda_3^2}} \right. \\
 & + \gamma_1(\cos\phi_\beta \cos\psi_\beta + \sin\phi_\beta \sin\theta_\beta \sin\psi_\beta) \\
 & + \left. \left( \frac{C_1 \sin\phi_\beta \cos\theta_\beta}{\sqrt{C_1^2 + C_2^2 + C_3^2}} \right) \right] M_{1_\alpha} \\
 & + \left[ \frac{\lambda_2(-\cos\phi_\beta \sin\psi_\beta + \sin\phi_\beta \sin\theta_\beta \cos\psi_\beta)}{\sqrt{\lambda_0^2 + \lambda_1^2 + \lambda_2^2 + \lambda_3^2}} \right. \\
 & + \gamma_2(\cos\phi_\beta \cos\psi_\beta + \sin\phi_\beta \sin\theta_\beta \sin\psi_\beta) \\
 & + \left. \left( \frac{C_2 \sin\phi_\beta \cos\theta_\beta}{\sqrt{C_1^2 + C_2^2 + C_3^2}} \right) \right] M_{2_\alpha} \\
 & + \left[ \frac{\lambda_3(-\cos\phi_\beta \sin\psi_\beta + \sin\phi_\beta \sin\theta_\beta \cos\psi_\beta)}{\sqrt{\lambda_0^2 + \lambda_1^2 + \lambda_2^2 + \lambda_3^2}} \right. \\
 & + \gamma_3(\cos\phi_\beta \cos\psi_\beta + \sin\phi_\beta \sin\theta_\beta \sin\psi_\beta) \\
 & + \left. \left( \frac{C_3 \sin\phi_\beta \cos\theta_\beta}{\sqrt{C_1^2 + C_2^2 + C_3^2}} \right) \right] M_{3_\alpha} \tag{3.15}
 \end{aligned}$$

$$\begin{aligned}
 M_{3_\beta} = & \left[ \frac{\lambda_0(\sin\phi_\beta \sin\psi_\beta + \cos\phi_\beta \sin\theta_\beta \sin\psi_\beta)}{\sqrt{\lambda_0^2 + \lambda_1^2 + \lambda_2^2 + \lambda_3^2}} \right. \\
 & + \gamma_0(-\sin\phi_\beta \cos\psi_\beta + \cos\phi_\beta \sin\theta_\beta \sin\psi_\beta) \left. \right] T_\alpha \\
 & + \left[ \frac{\lambda_1(\sin\phi_\beta \sin\psi_\beta + \cos\phi_\beta \sin\theta_\beta \sin\psi_\beta)}{\sqrt{\lambda_0^2 + \lambda_1^2 + \lambda_2^2 + \lambda_3^2}} \right. \\
 & + \gamma_1(-\sin\phi_\beta \cos\psi_\beta + \cos\phi_\beta \sin\theta_\beta \sin\psi_\beta) \\
 & + \left. \frac{C_1 \cos\phi_\beta \cos\theta_\beta}{\sqrt{C_1^2 + C_2^2 + C_3^2}} \right] M_{1_\alpha} \\
 & + \left[ \frac{\lambda_2(\sin\phi_\beta \sin\psi_\beta + \cos\phi_\beta \sin\theta_\beta \sin\psi_\beta)}{\sqrt{\lambda_0^2 + \lambda_1^2 + \lambda_2^2 + \lambda_3^2}} \right. \\
 & + \gamma_2(-\sin\phi_\beta \cos\psi_\beta + \cos\phi_\beta \sin\theta_\beta \sin\psi_\beta) \\
 & + \left. \frac{C_2 \cos\phi_\beta \cos\theta_\beta}{\sqrt{C_1^2 + C_2^2 + C_3^2}} \right] M_{2_\alpha} \\
 & + \left[ \frac{\lambda_3(\sin\phi_\beta \sin\psi_\beta + \cos\phi_\beta \sin\theta_\beta \sin\psi_\beta)}{\sqrt{\lambda_0^2 + \lambda_1^2 + \lambda_2^2 + \lambda_3^2}} \right. \\
 & + \gamma_3(-\sin\phi_\beta \cos\psi_\beta + \cos\phi_\beta \sin\theta_\beta \sin\psi_\beta) \\
 & + \left. \frac{C_3 \cos\phi_\beta \cos\theta_\beta}{\sqrt{C_1^2 + C_2^2 + C_3^2}} \right] M_{3_\alpha} \tag{3.16}
 \end{aligned}$$

Note that

$$\theta_\beta = \int \frac{k_{3_\beta}}{\sqrt{k_{1_\beta}^2 + k_{2_\beta}^2}} ds_\beta, \quad \psi_\beta = - \int \left[ k_{2_\beta} + k_{3_\beta} \frac{\sqrt{k_{3_\beta}^2 - \theta_\beta'^2}}{\sqrt{k_{1_\beta}^2 + k_{2_\beta}^2}} \right] ds_\beta,$$

$$\text{and } \phi_\beta = - \int \frac{\sqrt{k_{3_\beta}^2 - \theta_\beta'^2}}{\cos\theta_\beta} ds_\beta$$

The second curvature of the curve  $\beta$  according to Frenet frame is given by

$$k_{2_\beta} = \sqrt{\frac{C_1^2 + C_2^2 + C_3^2}{\sqrt{\lambda_0^2 + \lambda_1^2 + \lambda_2^2 + \lambda_3^2}}} \tag{3.17}$$

The third curvature of the curve  $\beta$  according to Frenet frame is given by

$$k_{3_\beta} = \frac{-(C_1 K_{1_\alpha} + C_2 K_{2_\alpha} + C_3 K_{3_\alpha})(\lambda_1 K_{1_\alpha} + \lambda_2 K_{2_\alpha} + \lambda_3 K_{3_\alpha})}{\sqrt{C_1^2 + C_2^2 + C_3^2}} \tag{3.18}$$

The first curvature of the curve  $\beta$  according to parallel transport frame reads

$$K_{1_\beta} = \sqrt{\lambda_0^2 + \lambda_1^2 + \lambda_2^2 + \lambda_3^2} \cos\theta_\beta \cos\psi_\beta \tag{3.19}$$

The second curvature of the curve  $\beta$  according to parallel transport frame reads

$$K_{2_\beta} = \sqrt{\lambda_0^2 + \lambda_1^2 + \lambda_2^2 + \lambda_3^2} [-\cos\phi_\beta \sin\psi_\beta + \sin\phi_\beta \sin\theta_\beta \cos\psi_\beta] \tag{3.20}$$

The third curvature of the curve  $\beta$  according to parallel transport frame reads

$$K_{3_\beta} = \sqrt{\lambda_0^2 + \lambda_1^2 + \lambda_2^2 + \lambda_3^2} [\sin\phi_\beta \sin\psi_\beta + \cos\phi_\beta \sin\theta_\beta \cos\psi_\beta] \tag{3.21}$$

The proof is complete.  $\square$

**Acknowledgement**

I am grateful to the referee for his/her valuable comments and suggestions.

**References**

- [1] M. Turgut, Smarandache breadth pseudo null curves in Minkowski space-time, *Int. J. Math. Combin.* Vol. 1 (2009) 46–49.
- [2] M. Turgut, S. Yilmaz, Smarandache curves in Minkowski space-time, *Int. J. Math. Combin.* Vol. 3 (2008) 51–55.
- [3] U. Ozturkab, E.B.K. Ozturkab, K. İlarıslanc, E. Neřovićđ, On Smarandache curves lying in lightcone in Minkowski 3-space, *J. Dyn. Syst. Geom. Theor.* Vol. 12 (Number 1) (2014) 81–91.
- [4] A. Magden, S. Yilmaz, On the curves of constant breadth in four dimensional Galilean space, *Int. Math. Forum* Vol. 9 (no. 25) (2014) 1229–1236.
- [5] A.T. Ali, Special Smarandache curves in the Euclidean space, *Int. J. Math. Combin.* Vol. 2 (2010) 30–36.
- [6] R. Bishop, There is more than one way to frame a curve, in: *The American Mathematical Monthly* March, 1975, pp. 246–252.
- [7] S. Buyukkutuk, G. Ozturk, Constant ratio curves according to parallel transport frame in Euclidean 4-space  $E^4$ , *New Trends Math. Sci. NTMSCI* 3 (No. 4) (2015) 171–178.
- [8] T. Krpınar, E. Turhan, Biharmonic curves according to parallel transport frame in  $E^4$ , *Bol. Soc. Paran. Mat.* (3s.) v. 31 (2) (2013) 213–217.
- [9] F. celik, Z. Bozkurt, I. Gok, F.N. Ekmekci, Y. Yayl, Parallel transport frame in 4-dimensional euclidean space  $E^4$ , *Caspian J. Math. Sci. (CJMS)* 3 (1) (2014) 91–103.
- [10] A.T. Yakut, M. SavaG, T. Tamirci, The Smarandache curves on  $S_2^2$  and its duality on  $H_0^2$ , *J. Appl. Math.* Volume (2014) 12. Hindawi Publishing Corporation, Article ID 193586.
- [11] M. Cetin, H. Kocayigıt, On the quaternionic Smarandache curves in Euclidean 3-space, *Int. J. Contemp. Math. Sciences* Vol. 8 (no. 3) (2013) 139–150.
- [12] M. Cetin, Y. Tuncer, M.K. Karacan, Smarandache curves according to Bishop frame in Euclidean 3-space, *Gen. Math. Notes* Vol. 20 (No. 2) (February 2014) 50–66.
- [13] S. SENYURT, A. CALISKAN, N C- Smarandache curves of Mannheim curve couple according to Frenet frame, *Int. J. Math. Combin.* Vol.1 (2015) 1–13.
- [14] S. Senyurt, A. Calskan, An application according to spatial quaternionic Smarandache curve, *Appl. Math. Sci.* Vol. 9 (no. 5) (2015) 219–228.
- [15] M. Elzawy, S. Mosa, Smarandache curves in the Galilean 4-space  $G_4$ , *J. Egypt. Math. Soc.* 25 (2017) 53–56.