

# On an inequality for the Smarandache function

J. Sándor

Babeş-Bolyai University, 3400 Cluj-Napoca, Romania

1. In paper [2] the author proved among others the inequality  $S(ab) \leq aS(b)$  for all  $a, b$  positive integers. This was refined to

$$S(ab) \leq S(a) + S(b) \quad (1)$$

in [1]. Our aim is to show that certain results from our recent paper [3] can be obtained in a simpler way from a generalization of relation (1). On the other hand, by the method of Le [1] we can deduce similar, more complicated inequalities of type (1).

2. By mathematical induction we have from (1) immediately:

$$S(a_1 a_2 \dots a_n) \leq S(a_1) + S(a_2) + \dots + S(a_n) \quad (2)$$

for all integers  $a_i \geq 1$  ( $i = 1, \dots, n$ ). When  $a_1 = \dots = a_n = n$  we obtain

$$S(a^n) \leq nS(a). \quad (3)$$

For three applications of this inequality, remark that

$$S((m!)^n) \leq nS(m!) = nm \quad (4)$$

since  $S(m!) = m$ . This is inequality 3) part 1. from [3]. By the same way,  $S((n!)^{(n-1)!}) \leq (n-1)!S(n!) = (n-1)!n = n!$ , i.e.

$$S((n!)^{(n-1)!}) \leq n! \quad (5)$$

Inequality (5) has been obtained in [3] by other arguments (see 4) part 1.).

Finally, by  $S(n^2) \leq 2S(n) \leq n$  for  $n$  even (see [3], inequality 1),  $n > 4$ , we have obtained a refinement of  $S(n^2) \leq n$ :

$$S(n^2) \leq 2S(n) \leq n \quad (6)$$

for  $n > 4$ , even.

3. Let  $m$  be a divisor of  $n$ , i.e.  $n = km$ . Then (1) gives  $S(n) = S(km) \leq S(m) + S(k)$ , so we obtain:

If  $m|n$ , then

$$S(n) - S(m) \leq S\left(\frac{n}{m}\right). \quad (7)$$

As an application of (7), let  $d(n)$  be the number of divisors of  $n$ . Since  $\prod_{k|n} k = n^{d(n)/2}$ , and  $\prod_{k \leq n} k = n!$  (see [3]), and by  $\prod_{k|n} k | \prod_{k \leq n} k$ , from (7) we can deduce that

$$S(n^{d(n)/2}) + S(n!/n^{d(n)/2}) \geq n. \quad (8)$$

This improves our relation (10) from [3].

4. Let  $S(a) = u$ ,  $S(b) = v$ . Then  $b|v!$  and  $u!|x(x-1)\dots(x-u+1)$  for all integers  $x \geq u$ . But from  $a|u!$  we have  $a|x(x-1)\dots(x-u+1)$  for all  $x \geq u$ . Let  $x = u+v+k$  ( $k \geq 1$ ). Then, clearly  $ab(v+1)\dots(v+k)|(u+v+k)!$ , so we have  $S[ab(v+1)\dots(v+k)] \leq u+v+k$ . Here  $v = S(b)$ , so we have obtained that

$$S[ab(S(b)+1)\dots(S(b)+k)] \leq S(a) + S(b) + k. \quad (9)$$

For example, for  $k = 1$  one has

$$S[ab(S(b)+1)] \leq S(a) + S(b) + 1. \quad (10)$$

This is not a consequence of (2) for  $n = 3$ , since  $S[S(b)+1]$  may be much larger than 1.

## References

- [1] M. Le, *An inequality concerning the Smarandache function*, Smarandache Notions J., vol. 9(1998), 124-125.
- [2] J. Sándor, *On certain inequalities involving the Smarandache function*, Smarandache Notions J., vol. 7(1996), 3-6.
- [3] J. Sándor, *On certain new inequalities and limits for the Smarandache function*, Smarandache Notions J., vol. 9(1998), 63-69.