

# SMARANDACHE MAXIMUM RECIPROCAL REPRESENTATION FUNCTION

(Amarnath Murthy ,S.E. (E &T), Well Logging Services, Oil And Natural  
Gas Corporation Ltd. ,Sabarmati, Ahmedbad, India- 380005.)

**ABSTRACT:** Smarandache Maximum Reciprocal Representation

(SMRR) Function  $f_{SMRR}(n)$  is defined as follows

$f_{SMRR}(n) = t$  if

$$\sum_{r=1}^t 1/r \leq n \leq \sum_{r=1}^{t+1} 1/r$$

## SMARANDACHE MAXIMUM RECIPROCAL REPRESENTATION SEQUENCE

SMRRS is defined as  $T_n = f_{SMRR}(n)$

$$f_{SMRR}(1) = 1$$

$$f_{SMRR}(2) = 3, \quad (1 + 1/2 + 1/3 < 2 < 1 + 1/2 + 1/3 + 1/4)$$

$$f_{SMRR}(3) = 10$$

$$\sum_{r=1}^{10} 1/r \leq 3 \leq \sum_{r=1}^{11} 1/r$$

SMRRS is

**1, 3, 10, . . .**

**A note on The SMRR Function:**

The harmonic series  $\sum 1/n$  satisfies the following inequality

$$\log (n+1) < \sum 1/n < \log n + 1 \quad \text{-----(1)}$$

This inequality can be derived as follows

We have  $e^x > 1 + x > 1, x > 0$

and  $(1 + 1/n)^{(1 + 1/n)} > 1, n > 0$

which gives

$$1/(r+1) < \log(1 + 1/r) < 1/r$$

summing up for  $r = 1$  to  $n+1$  and with some algebraic jugglery we get (1). With the help of (1) we get the following result on the **SMRR function**.

**If  $SMRR(n) = m$  then  $[\log(m)] \approx n - 1$**

Where  $[\log(m)]$  stands for the integer value of  $\log(m)$ .

### **SOME CONJECTURES:**

(1.1). Every positive integer can be expressed as the sum of the reciprocal of a finite number of distinct natural numbers. ( in infinitely many ways.).

Let us define a function  $R_m(n)$  as the minimum number of natural numbers required for such an expression.

(1.2). Every natural number can be expressed as the sum of the reciprocals of a set of natural numbers which are in Arithmetic Progression.

(1.3). Let

$$\sum 1/r \leq n \leq \sum 1/(r+1)$$

where  $\sum 1/r$  stands for the sum of the reciprocals of first  $r$

natural numbers and let  $S_1 = \sum 1/r$

let  $S_2 = S_1 + 1/(r+k_1)$  such that  $S_2 + 1/(r+k_1+1) > n \geq S_2$

let  $S_3 = S_2 + 1/(r+k_2)$  such that  $S_3 + 1/(r+k_2+1) > n \geq S_3$

and so on , then there exists a finite  $m$  such that

$$S_{m+1} + 1/(r+k_m) = n$$

**Remarks :** The veracity of conjecture (1.1) is deducible from conjecture (1.3) .

(1.4). (a) There are infinitely many disjoint sets of natural numbers sum of whose reciprocals is unity.

(b) Among the sets mentioned in (a) , there are sets which can be organised in an order such that the largest element of any set is smaller than the smallest element of the next set.

#### REFERENCES:

- [1] 'Smarandache Notion Journal' Vol. 10 ,No. 1-2-3, Spring 1999. Number Theory Association of the UNIVERSITY OF CRAIOVA.
- [2] "Amarnath Murthy" , 'Smarandache Reciprocal Partition Of Unity Sets And Sequences', SNJ, Vol. 11, No. 1-2-3, 2000.
- [3] " The Florentine Smarandache " Special Collection, Archives of American Mathematics, Centre for American History, University of Texax at Austin, USA.