

## SMARANDACHE PASCAL DERIVED SEQUENCES

(Amarnath Murthy, S.E.(E&T), WLS, Oil and Natural Gas Corporation Ltd.,  
Sabarmati, Ahmedabad,-380005 INDIA. )

Given a sequence say  $S_b$ . We call it the base sequence. We define a **Smarandache Pascal derived sequence  $S_d$**  as follows:

$$T_{n+1} = \sum_{k=0}^n {}^n C_k \cdot t_{k+1}, \text{ where } t_k \text{ is the } k^{\text{th}} \text{ term of the base sequence.}$$

Let the terms of the the base sequence  $S_b$  be

$$b_1, b_2, b_3, b_4, \dots$$

Then the Smarandache Pascal derived Sequence  $S_d$

$d_1, d_2, d_3, d_4, \dots$  is defined as follows:

$$d_1 = b_1$$

$$d_2 = b_1 + b_2$$

$$d_3 = b_1 + 2b_2 + b_3$$

$$d_4 = b_1 + 3b_2 + 3b_3 + b_4$$

...

$$d_{n+1} = \sum_{k=0}^n {}^n C_k \cdot b_{k+1}$$

These derived sequences exhibit interesting properties for some base sequences.

Examples:

{1}  $S_b \rightarrow 1, 2, 3, 4, \dots$  ( natural numbers)

$S_d \rightarrow 1, 3, 8, 20, 48, 112, 256, \dots$  ( Smarandache Pascal derived natural number sequence)

The same can be rewritten as

$$2 \times 2^{-1}, 3 \times 2^0, 4 \times 2^1, 5 \times 2^2, 6 \times 2^3, \dots$$

It can be verified and then proved easily that  $T_n = 4(T_{n-1} - T_{n-2})$  for  $n > 2$ .

And also that  $T_n = (n+1) \cdot 2^{n-2}$

{2}  $S_b \rightarrow 1, 3, 5, 7, \dots$  ( odd numbers)

$S_d \rightarrow 1, 4, 12, 32, 80, \dots$

The first difference  $1, 3, 8, 20, 48, \dots$  is the same as the  $S_d$  for natural numbers.

The sequence  $S_d$  can be rewritten as

$$1 \cdot 2^0, 2 \cdot 2^1, 3 \cdot 2^2, 4 \cdot 2^3, 5 \cdot 2^4, \dots$$

Again we have  $T_n = 4(T_{n-1} - T_{n-2})$  for  $n > 2$ . Also  $T_n = n \cdot 2^{n-1}$ .

{3} **Smarandache Pascal Derived Bell Sequence:**

Consider the Smarandache Factor Partitions (SFP) sequence for the square free numbers:

( The same as the **Bell number** sequence.)

$S_b \rightarrow 1, 1, 2, 5, 15, 52, 203, 877, 4140, \dots$

We get the derived sequence as follows

$S_d \rightarrow 1, 2, 5, 15, 52, 203, 877, 4140, \dots$

The **Smarandache Pascal Derived Bell Sequence** comes out to be the same. We

call it **Pascal Self Derived Sequence**. This has been established in ref. [1]  
 In what follows, we shall see that this Transformation applied to Fibonacci  
 Numbers gives beautiful results.

**\*\*{4} Smarandache Pascal derived Fibonacci Sequence:**

Consider the Fibonacci Sequence as the Base Sequence:

$$S_b \rightarrow 1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, 144, 233, \dots$$

We get the following derived sequence

$$S_d \rightarrow 1, 2, 5, 13, 34, 89, 233, \dots \quad (\text{A})$$

It can be noticed that the above sequence is made of the alternate (even numbered  
 terms of the sequence ) Fibonacci numbers.

This gives us the following result on the Fibonacci numbers.

$$F_{2n} = \sum_{k=0}^n {}^n C_k \cdot F_k, \text{ where } F_k \text{ is the } k^{\text{th}} \text{ term of the base Fibonacci sequence.}$$

Some more interesting properties are given below.

If we take (A) as the base sequence we get the following derived sequence  $S_{dd}$

$$S_{dd} \rightarrow 1, 3, 10, 35, 125, 450, 1625, 5875, 21250, \dots$$

An interesting observation is ,the first two terms are divisible by  $5^0$ , the next two  
 terms by  $5^1$ , the next two by  $5^2$ , the next two by  $5^3$  and so on.

$$T_{2n} \equiv T_{2n-1} \equiv 0 \pmod{5^n}$$

On carrying out this division we get the following sequence i.e.

$$1, 3, 2, 7, 5, 18, 13, 47, 34, 123, 89, \dots \quad (\text{B})$$

The sequence formed by the odd numbered terms is

$$1, 2, 5, 13, 34, 89, \dots$$

which is again nothing but  $S_d$  ( the base sequence itself).

Another interesting observation is every even numbered term of (B) is the sum of  
 the two adjacent odd numbered terms. (  $3 = 1+2$ ,  $7 = 2 +5$ ,  $18 = 5 + 13$  etc.)

**CONJECTURE:** Thus we have the possibility of another beautiful result on the  
 Fibonacci numbers which of-course is yet to be established.

$$F_{2m+1} = (1/5^m) \sum_{r=0}^{2m+1} C_r \left( \sum_{k=0}^r C_k F_k \right)$$

**Note:** It can be verified that all the above properties hold good for the Lucas  
 sequence ( 1 , 3 , 4 , 7 , 11 , ... ) as well.

**Pascalisation of Fibonacci sequence with index in arithmetic progression:**

Consider the following sequence formed by the Fibonacci numbers whose indexes  
 are in A. P.

$F_1, F_{d+1}, F_{2d+1}, F_{3d+1}, \dots$  on pascalisation gives the following sequence

$$1, d.F_2, d^2.F_4, d^3.F_6, d^4.F_8, \dots, d^n.F_{2n}, \dots$$

for  $d = 2$  and  $d = 3$ .

For  $d = 5$  we get the following

Base sequence :  $F_1, F_6, F_{11}, F_{16}, \dots$

$$1, 13, 233, 4181, 46368, \dots$$

Derived sequence: 1, 14, 260, 4920, 93200, . . . in which we notice that  
 $260 = 20 \cdot (14 - 1)$ ,  $4920 = 20 \cdot (260 - 14)$ ,  $93200 = 4920 - 260$  ) etc . which suggests  
the possibility of

**Conjecture: The terms of the pascal derived sequence for  $d = 5$  are given by**  
 $T_n = 20 \cdot (T_{n-1} - T_{n-2})$  ( $n > 2$ )

For  $d = 8$  we get

Base sequence :  $F_1, F_9, F_{17}, F_{25}, \dots$

$S_b \rightarrow 1, 34, 1597, 75025, \dots$

$S_d \rightarrow 1, 35, 1666, 79919, \dots$

$= 1, 35, (35-1) \cdot 7^2, (1666 - 35) \cdot 7^2, \dots$  etc. which suggests the possibility of

**Conjecture: The terms of the pascal derived sequence for  $d = 8$  are given by**  
 $T_n = 49 \cdot (T_{n-1} - T_{n-2})$ , ( $n > 2$ )

Similarly we have Conjectures:

For  $d = 10$ ,  $T_n = 90 \cdot (T_{n-1} - T_{n-2})$ , ( $n > 2$ )

For  $d = 12$ ,  $T_n = 18^2 \cdot (T_{n-1} - T_{n-2})$ , ( $n > 2$ )

**Note:** There seems to be a direct relation between  $d$  and the coefficient of  $(T_{n-1} - T_{n-2})$  (or the common factor) of each term which is to be explored.

**{5} Smarandache Pascal derived square sequence:**

$S_b \rightarrow 1, 4, 9, 16, 25, \dots$

$S_d \rightarrow 1, 5, 18, 56, 160, 432, \dots$

Or  $1, 5 \times 1, 6 \times 3, 7 \times 8, 8 \times 20, 9 \times 48, \dots$ , ( $T_n = (n+3)t_{n-1}$ ), where  $t_r$  is the  $r^{\text{th}}$  term of Pascal derived natural number sequence.

Also one can derive  $T_n = 2^{n-2} \cdot (n+3)(n)/2$ .

**{6} Smarandache Pascal derived cube sequence:**

$S_b \rightarrow 1, 8, 27, 64, 125$

$S_d \rightarrow 1, 9, 44, 170, 576, 1792, \dots$

We have  $T_n \equiv 0 \pmod{(n+1)}$ .

Similarly we have derived sequences for higher powers which can be analyzed for patterns.

**{7} Smarandache Pascal derived Triangular number sequence:**

$S_b \rightarrow 1, 3, 6, 10, 15, 21, \dots$

$S_d \rightarrow 1, 4, 13, 38, 104, 272, \dots$

**{8} Smarandache Pascal derived Factorial sequence:**

$S_b \rightarrow 1, 2, 6, 24, 120, 720, 5040, \dots$

$S_d \rightarrow 1, 3, 11, 49, 261, 1631, \dots$

We can verify that  $T_n = n \cdot T_{n-1} + \sum T_{n-2} + 1$ .

**Problem: Are there infinitely many primes in the above sequence?**

**Smarandache Pascal derived sequence of the  $k^{\text{th}}$  order.**

Consider the natural number sequence again:

$S_b \rightarrow 1, 2, 3, 4, 5, \dots$  The corresponding derived sequence is

$S_d \rightarrow 2 \times 2^1, 3 \times 2^0, 4 \times 2^1, 5 \times 2^2, 6 \times 2^3, \dots$  With this as the base sequence we get the derived sequence denoted by  $S_{d2}$  as

$S_{d2}$  or  $S_{d2} \rightarrow 1, 4, 15, 54, 189, 648, \dots$  which can be rewritten as

$1, 4 \times 3^0, 5 \times 3^1, 6 \times 3^2, 7 \times 3^3 \dots$

similarly we get  $S_{d3}$  as  $1, 5 \times 4^0, 6 \times 4^1, 7 \times 4^2, 8 \times 4^3, \dots$  which suggests the

possibility of the terms of  $S_{dk}$ , the  $k^{\text{th}}$  order Smarandache Pascal derived natural

number sequence being given by

$1, (k+2) \cdot (k+1)^0, (k+3) \cdot (k+1)^1, (k+4) \cdot (k+1)^2, \dots, (k+r) \cdot (k+1)^{r-2}$  etc. This can be proved by induction.

**We can take an arithmetic progression with the first term 'a' and the common difference 'b' as the base sequence and get the derived  $k^{\text{th}}$  order sequences to generalize the above results.**

**Reference:[1] Amarnath Murthy, 'Generalization of Partition Function,. Introducing Smarandache Factor Partitions' SNJ, Vol. 11, No. 1-2-3,2000.**