

Interpretation of Solution of Radial Biquaternion Klein-Gordon Equation and Comparison with EQPET/TSC Model

V. Christianto* and F. Smarandache**

Abstract

In a previous publication,¹ we argued that the biquaternionic extension of the Klein-Gordon equation has numerical solution with sinusoidal form, which differs appreciably from conventional Yukawa potential. In the present article we interpret and compare this result from the viewpoint of the EQPET/TSC (Electronic Quasi-Particle Expansion Theory/Tetrahedral Symmetric Condensate) model described by Takahashi.² Further observation is of course recommended in order to refute or verify this proposition.

Introduction

In the preceding article¹ we argued that biquaternionic extension of the radial Klein-Gordon equation (radialBQKGE) has numerical solution with sinusoidal form, which differs appreciably from conventional Yukawa potential. We also argued that this biquaternionic extension of KGE may be useful in particular to explore new effects in the context of low-energy nuclear reaction (LENR).³

Interestingly, Takahashi² has discussed key experimental results in condensed matter nuclear effects in light of EQPET/TSC. We argue here that the potential model used in his paper, STTBA (Sudden Tall Thin Barrier Approximation), may be comparable to our derived sinusoidal potential from radial biquaternion KGE.¹ While we can't yet offer numerical prediction, our qualitative comparison may be useful in verifying further experiments.

Solution of Radial Biquaternionic KGE (radial BQKGE)

In our previous paper,¹ we argued that it is possible to write the biquaternionic extension of the Klein-Gordon equation as follows:

$$(\diamond\bar{\diamond} + m^2) \varphi(x, t) = 0 \quad (1)$$

Provided we use this definition:^{1,3}

$$\begin{aligned} \diamond = \nabla^q + i\nabla^q = & \left(-i \frac{\partial}{\partial t} + e_1 \frac{\partial}{\partial X} + e_2 \frac{\partial}{\partial Y} + e_3 \frac{\partial}{\partial Z} \right) \\ & + i \left(-i \frac{\partial}{\partial T} + e_1 \frac{\partial}{\partial X} + e_2 \frac{\partial}{\partial Y} + e_3 \frac{\partial}{\partial Z} \right) \end{aligned} \quad (2)$$

where e_1, e_2, e_3 are *quaternion imaginary units* obeying (with ordinary quaternion symbols $e_1=i, e_2=j, e_3=k$):^{3,4}

$$\begin{aligned} i^2 = j^2 = k^2 = & -1, \quad ij = -ji = k, \\ jk = -kj = & i, \quad ki = -ik = j. \end{aligned} \quad (3)$$

And quaternion Nabla operator is defined as:²

$$\nabla^q = -i \frac{\partial}{\partial t} + e_1 \frac{\partial}{\partial X} + e_2 \frac{\partial}{\partial Y} + e_3 \frac{\partial}{\partial Z} \quad (4)$$

By using polar coordinates transformation,⁵ we get this for the one-dimensional situation:

$$\left(\frac{\partial}{\partial r} \left(\frac{\partial}{\partial r} \right) - i \frac{\partial}{\partial r} \left(\frac{\partial}{\partial r} \right) + m^2 \right) \varphi(x, t) = 0 \quad (5)$$

The solution is given by:¹

$$y = k_1 \cdot \sin \left(\frac{|m|r}{\sqrt{-i-1}} \right) + k_2 \cdot \cos \left(\frac{|m|r}{\sqrt{-i-1}} \right) \quad (6)$$

Therefore, we may conclude that numerical solution of radial biquaternionic extension of Klein-Gordon equation yields different potential compared to the well-known Yukawa potential.¹

$$u(r) = -\frac{g^2}{r} e^{-mr} \quad (7)$$

In the next section we will discuss an interpretation of this new potential (6) compared to the findings discussed by Takahashi² from condensed matter nuclear experiments.

Comparison with Takahashi's EQPET/TSC/STTBA model

Takahashi² reported some findings from condensed matter nuclear experiments, including intense production of helium-4 (⁴He) atoms by electrolysis and laser irradiation experiments.

Takahashi analyzed those experimental results using EQPET formation of TSC were modelled with numerical estimations by STTBA. This STTBA model includes strong interaction with negative potential near the center (where $r \rightarrow 0$). See Figure 1.

Takahashi described that Gamow integral of STTBA is given by:

$$\Gamma_n = 0.218 (\mu^{1/2}) \int_{r_0}^b (V_b - E_d)^{1/2} dr \quad (8)$$

Using $b=5.6$ fm and $r_0=5$ fm, he obtained:

$$P_{Ad} = 0.77 \quad (9)$$

and

$$V_B = 0.257 \text{ MeV} \quad (10)$$

While his EQPET model gave significant underestimation for 4D fusion rate when rigid constraint of motion in 3D space was attained, introducing different values of λ_{4d} can improve the result.² Therefore we may conclude that STTBA can offer good approximation of condensed matter nuclear reactions.⁵




Figure 1. Potential for Coulomb barrier reversal for STTBA, from Takahashi.²

Interestingly, the STTBA lacks sufficient theoretical basis, therefore one can expect that a sinusoidal form (or combined sinusoidal waves such as in Fourier method) may offer better result which agrees with experiments. This will be pursued in a later paper.

Nonetheless, we recommend further observation in order to refute or verify this proposition of a new type of potential derived from the biquaternion radial Klein-Gordon equation.

Acknowledgment

V. Christianto would like to thank Prof. C. Castro for numerous discussions.

References

1. Christianto, V. and Smarandache, F. 2008. "Numerical Solution of Radial Biquaternion Klein-Gordon Equation," *Progress in Physics*, 4, 1, January.

2. Takahashi, A. 2005. "A Theoretical Summary of Condensed Matter Nuclear Effects," Siena Workshop on Anomalies in Metal-D/H Systems, Siena, May (<http://lenr-canr.org>).

3. Yefremov, A., Smarandache, F., and Christianto, V. 2007. "Yang-Mills Field from Quaternion Space Geometry, and Its Klein-Gordon Representation," *Progress in Physics*, 3, 3, www.ptep-online.com.

4. Christianto, V. 2006. "A New Wave Quantum Relativistic Equation from Quaternionic Representation of Maxwell-Dirac Equation as an Alternative to Barut-Dirac Equation," *Electronic Journal of Theoretical Physics*, 3, 12, www.ejtp.com.

5. Storms, E. <http://lenr-canr.org>.

6. Nishikawa, M. 2004. "A Derivation of Electroweak Unified and Quantum Gravity Theory without Assuming Higgs Particle," arXiv:hep-th/0407057, p.15.

7. Maxima from <http://maxima.sourceforge.net>. (Using Lisp GNU Common Lisp).

About the Authors

Dr. Christianto _____

Dr. Smarandache was born in Romania and received a Ph.D. in mathematics from the State University of Kishinev in 1997. After emigration to the U.S., he continued postdoctoral studies at various universities. Smarandache worked as a software engineer for Honeywell (1990-1995) and was an adjunct professor at Pima Community College (1995-1997). In 1997 he became an assistant professor at the University of New Mexico (Gallup) and was promoted to associate professor of mathematics in 2003. Smarandache is the author of many books and has published in many journals and proceedings.

*Email: admin@sciprint.org ; website: www.sciprint.org

**Department of Mathematics, University of New Mexico, Gallup, NM ; Email: fsmarandache@yahoo.com