

# A Neutrosophic Extension of the MULTIMOORA Method

Dragisa STANUJKIC<sup>1\*</sup>, Edmundas Kazimieras ZAVADSKAS<sup>2</sup>,  
Florentin SMARANDACHE<sup>3</sup>, Willem K.M. BRAUERS<sup>4</sup>,  
Darjan KARABASEVIC<sup>5</sup>

<sup>1</sup>*Faculty of Management in Zajecar, John Naisbitt University, Belgrade, Serbia*

<sup>2</sup>*Research Institute of Smart Building Technologies*

*Civil Engineering Faculty, Vilnius Gediminas Technical University, Vilnius, Lithuania*

<sup>3</sup>*Department of Mathematics, University of New Mexico  
705 Gurley Avenue, Gallup, NM 87301, USA*

<sup>4</sup>*Faculty of Applied Economics, University of Antwerp, Antwerp, Belgium*

<sup>5</sup>*Faculty of Applied Management, Economics and Finance, Business Academy University  
Jevrejska 24, Belgrade, Serbia*

*e-mail: dragisa.stanujkic@fmz.edu.rs, edmundas.zavadskas@vgtu.lt, fsmarandache@gmail.com,  
willem.brauers@uantwerpen.be, darjankarabasevic@gmail.com*

Received: October 2016; accepted: February 2017

**Abstract.** The aim of this manuscript is to propose a new extension of the MULTIMOORA method adapted for usage with a neutrosophic set. By using single valued neutrosophic sets, the MULTIMOORA method can be more efficient for solving complex problems whose solving requires assessment and prediction, i.e. those problems associated with inaccurate and unreliable data. The suitability of the proposed approach is presented through an example.

**Key words:** neutrosophic set, single valued neutrosophic set, MULTIMOORA, MCDM.

## 1. Introduction

The MULTIMOORA (Multi-Objective Optimization by a Ratio Analysis plus the Full Multiplicative Form) was proposed by Brauers and Zavadskas (2010).

The ordinary MULTIMOORA method has been proposed for usage with crisp numbers. In order to enable its use in solving a larger number of complex decision-making problems, several extensions have been proposed, out of which only the most prominent are mentioned: Brauers *et al.* (2011) proposed a fuzzy extension of the MULTIMOORA method; Balezentis and Zeng (2013) proposed an interval-valued fuzzy extension; Balezentis *et al.* (2014) proposed an intuitionistic fuzzy extension and Zavadskas *et al.* (2015) proposed an interval-valued intuitionistic extension of the MULTIMOORA method.

---

\* Corresponding author.

The MULTIMOORA method has been applied for the purpose of solving a wide range of problems.

As some of the most cited, the studies that consider different problems in economics (Brauers and Zavadskas, 2010, 2011; Brauers, 2010), personnel selection (Balezentis *et al.*, 2012a, 2012b), construction (Kracka *et al.*, 2015), risk management (Liu *et al.*, 2014a) and waste treatment (Liu *et al.*, 2014b) can be mentioned.

As some of the newest studies in which the MULTIMOORA method is used for solving various decision-making problems, the following ones can be mentioned: material selection (Hafezalkotob and Hafezalkotob, 2016; Hafezalkotob *et al.*, 2016) and the CNC machine tool evaluation (Sahu *et al.*, 2016).

A significant approach in solving complex decision-making problems was formed by adapting the multiple criteria decision-making methods for the purpose of using fuzzy numbers, proposed by Zadeh in the fuzzy set theory (Zadeh, 1965).

Based on the fuzzy set theory, some extensions are also proposed, such as: interval-valued fuzzy sets (Turksen, 1986), intuitionistic fuzzy sets (Atanassov, 1986) and interval-valued intuitionistic fuzzy sets (Atanassov and Gargov, 1989).

In addition to the membership function proposed in fuzzy sets, Atanassov (1986) introduced the non-membership function that expresses non-membership to a set, thus having created the basis for solving a much larger number of decision-making problems.

The intuitionistic fuzzy set is composed of membership (the so-called truth-membership)  $T_A(x)$  and non-membership (the so-called falsity-membership)  $F_A(x)$ , which satisfies the conditions  $T_A(x), F_A(x) \in [0, 1]$  and  $0 \leq T_A(x) + F_A(x) \leq 1$ . Therefore, intuitionistic fuzzy sets are capable of operating with incomplete pieces of information, but do not include intermediate and inconsistent information (Li *et al.*, 2016).

In intuitionistic fuzzy sets, indeterminacy  $\pi_A(x)$  is  $1 - T_A(x) - F_A(x)$  by default. Smarandache (1998, 1999) further extended intuitionistic fuzzy sets by proposing Neutrosophic, also introducing independent indeterminacy-membership.

Such a proposed neutrosophic set is composed of three independent membership functions named the truth-membership  $T_A(x)$ , the falsity-membership  $F_A(x)$  and the indeterminacy-membership  $I_A(x)$  functions.

Smarandache (1999) and Wang *et al.* (2010) further proposed a single valued neutrosophic set, by modifying the conditions  $T_A(x), I_A(x)$  and  $F_A(x) \in [0, 1]$  and  $0 \leq T_A(x) + I_A(x) + F_A(x) \leq 3$ , which are more suitable for solving scientific and engineering problems (Li *et al.*, 2016).

Compared with the fuzzy set and its extensions, the single valued neutrosophic set can be identified as more flexible, for which reason an extension of the MULTIMOORA method adapted for the purpose of using the single valued neutrosophic set is proposed in this approach.

Therefore, the rest of this paper is organized as follows: in Section 2, some basic definitions related to the single valued neutrosophic set are given. In Section 3, the ordinary MULTIMOORA method is presented, whereas in Section 4, the Single Valued Neutrosophic Extension of the MULTIMOORA method is proposed. In Section 5, an example is considered with the aim to explain in detail the proposed methodology. The conclusions are presented in the final section.

**2. The Single Valued Neutrosophic Set**

DEFINITION 1. (See Smarandache, 1999.) Let  $X$  be the universe of discourse, with a generic element in  $X$  denoted by  $x$ . Then, the Neutrosophic Set (NS)  $A$  in  $X$  is as follows:

$$A = \{x \langle T_A(x), I_A(x), F_A(x) \rangle \mid x \in X\}, \tag{1}$$

where  $T_A(x)$ ,  $I_A(x)$  and  $F_A(x)$  are the truth-membership function, the indeterminacy-membership function and the falsity-membership function, respectively,

$$T_A, I_A, F_A : X \rightarrow ]^{-}0, 1^{+}[ \quad \text{and} \quad ]^{-}0 \leq T_A(x) + I_A(x) + F_A(x) \leq 3^{+}.$$

DEFINITION 2. (See Smarandache, 1999; Wang *et al.*, 2010.) Let  $X$  be the universe of discourse. The Single valued neutrosophic set (SVNS)  $A$  over  $X$  is an object having the following form:

$$A = \{x \langle T_A(x), I_A(x), F_A(x) \rangle \mid x \in X\}, \tag{2}$$

where  $T_A(x)$ ,  $I_A(x)$  and  $F_A(x)$  are the truth-membership function, the intermediacy-membership function and the falsity-membership function, respectively,

$$T_A, I_A, F_A : X \rightarrow [0, 1] \quad \text{and} \quad 0 \leq T_A(x) + I_A(x) + F_A(x) \leq 3.$$

DEFINITION 3. (See Smarandache, 1999.) For an SVNS  $A$  in  $X$ , the triple  $\langle t_A, i_A, f_A \rangle$  is called the single valued neutrosophic number (SVNN).

DEFINITION 4. Let  $x_1 = \langle t_1, i_1, f_1 \rangle$  and  $x_2 = \langle t_2, i_2, f_2 \rangle$  be two SVNNs and  $\lambda > 0$ ; then the basic operations are defined as follows:

$$x_1 + x_2 = \langle t_1 + t_2 - t_1 t_2, i_1 i_2, f_1 f_2 \rangle, \tag{3}$$

$$x_1 \cdot x_2 = \langle t_1 t_2, i_1 + i_2 - i_1 i_2, f_1 + f_2 - f_1 f_2 \rangle, \tag{4}$$

$$\lambda x_1 = \langle 1 - (1 - t_1)^\lambda, i_1^\lambda, f_1^\lambda \rangle, \tag{5}$$

$$x_1^\lambda = \langle t_1^\lambda, 1 - (1 - i_1)^\lambda, 1 - (1 - f_1)^\lambda \rangle. \tag{6}$$

DEFINITION 5. (See Sahin, 2014.) Let  $x = \langle t_x, i_x, f_x \rangle$  be an SVNN; then the score function  $s_x$  of  $x$  can be as follows:

$$s_x = (1 + t_x - 2i_x - f_x)/2, \tag{7}$$

where  $s_x \in [-1, 1]$ .

DEFINITION 6. Let  $x_1 = \langle t_1, i_1, f_1 \rangle$  and  $x_2 = \langle t_2, i_2, f_2 \rangle$  be two SVNNS. Then the maximum distance between  $x_1$  and  $x_2$  is as follows:

$$d_{\max}(x_1, x_2) = \begin{cases} |t_1 - t_2|, & x_1, x_2 \in \Omega_{\max}, \\ |f_1 - f_2|, & x_1, x_2 \in \Omega_{\min}. \end{cases} \quad (8)$$

DEFINITION 7. (See Sahin, 2014.) Let  $A_j = \langle t_j, i_j, f_j \rangle$  be a collection of SVNNS and  $W = (w_1, w_2, \dots, w_n)^T$  be an associated weighting vector. Then the Single Valued Neutrosophic Weighted Average (SVNWA) operator of  $A_j$  is as follows:

$$\begin{aligned} & SVNWA(A_1, A_2, \dots, A_n) \\ &= \sum_{j=1}^n w_j A_j = \left( 1 - \prod_{j=1}^n (1 - t_j)^{w_j}, \prod_{j=1}^n (i_j)^{w_j}, \prod_{j=1}^n (f_j)^{w_j} \right). \end{aligned} \quad (9)$$

where:  $w_j$  is the element  $j$  of the weighting vector,  $w_j \in [0, 1]$  and  $\sum_{j=1}^n w_j = 1$ .

DEFINITION 8. (See Sahin, 2014.) Let  $A_j = \langle t_j, i_j, f_j \rangle$  be a collection of SVNNS and  $W = (w_1, w_2, \dots, w_n)^T$  be an associated weighting vector. Then the Single Valued Neutrosophic Weighted Geometric (SVNWG) operator of  $A_j$  is as follows:

$$\begin{aligned} & SVNWG(A_1, A_2, \dots, A_n) \\ &= \prod_{j=1}^n (A_j)^{w_j} = \left( \prod_{j=1}^n (t_j)^{w_j}, 1 - \prod_{j=1}^n (1 - i_j)^{w_j}, 1 - \prod_{j=1}^n (1 - f_j)^{w_j} \right). \end{aligned} \quad (10)$$

where:  $w_j$  is the element  $j$  of the weighting vector,  $w_j \in [0, 1]$  and  $\sum_{j=1}^n w_j = 1$ .

### 3. The MULTIMOORA Method

The MULTIMOORA method consists of three approaches named as follows: the Ratio System (RS) Approach, the Reference Point (RP) Approach and the Full Multiplicative Form (FMF).

The considered alternatives are ranked based on all three approaches and the final ranking order and the final decision is made based on the theory of dominance. In other words, the alternative with the highest number of appearances in the first positions on all ranking lists is the best-ranked alternative.

**The ratio system approach.** In this approach, the overall importance of the alternative  $i$  can be calculated as follows:

$$y_i = y_i^+ - y_i^-, \quad (11)$$

with:

$$y_i^+ = \sum_{j \in \Omega_{\max}} w_j r_{ij}, \quad \text{and} \quad (12)$$

$$y_i^- = \sum_{j \in \Omega_{\min}} w_j r_{ij}, \quad (13)$$

$$r_{ij} = \frac{x_{ij}}{\sqrt{\sum_{i=1}^n x_{ij}^2}}, \quad (14)$$

where:  $y_i$  denotes the overall importance of the alternative  $i$ , obtained on the basis of all the criteria;  $y_i^+$  and  $y_i^-$  denote the overall importance of the alternative  $i$ , obtained on the basis of the benefit and cost criteria, respectively;  $r_{ij}$  denotes the normalized performance of the alternative  $i$  with respect to the criterion  $j$ ;  $x_{ij}$  denotes the performance of the alternative  $i$  to the criterion  $j$ ;  $\Omega_{\max}$  and  $\Omega_{\min}$  denote the sets of the benefit cost criteria, respectively;  $i = 1, 2, \dots, m$ ;  $m$  is the number of the alternatives,  $j = 1, 2, \dots, n$ ;  $n$  is the number of the criteria.

In this approach, the compared alternatives are ranked based on  $y_i$  in descending order and the alternative with the highest value of  $y_i$  is considered to be the best-ranked.

**The reference point approach.** The optimization based on this approach can be shown as follows:

$$d_i^{\max} = \max_j (w_j |r_j^* - r_{ij}|), \quad (15)$$

where:  $d_i^{\max}$  denotes the maximum distance of the alternative  $i$  to the reference point and  $r_j^*$  denotes the coordinate  $j$  of the reference point as follows:

$$r_j^* = \begin{cases} \max_i r_{ij}, & j \in \Omega_{\max}, \\ \min_i r_{ij}, & j \in \Omega_{\min}. \end{cases} \quad (16)$$

In this approach, the compared alternatives are ranked based on  $d_i^{\max}$  in ascending order and the alternative with the lowest value of  $d_i^{\max}$  is considered the best-ranked.

**The full multiplicative form.** In the FMF, the overall utility of the alternative  $i$  can be determined in the following manner:

$$u_i = \frac{a_i}{b_i}, \quad (17)$$

with:

$$a_i = \prod_{j \in \Omega_{\max}} w_j r_{ij}, \quad (18)$$

$$b_i = \prod_{j \in \Omega_{\min}} w_j r_{ij}, \quad (19)$$

where:  $u_i$  denotes the overall utility of the alternative  $i$ ,  $a_i$  denotes the product of the weighted performance ratings of the benefit criteria and  $b_i$  denotes the product of the weighted performance ratings of the cost criteria of the alternative  $i$ .

As in the RSA, the compared alternatives are ranked based on their  $u_i$  in descending order and the alternative with the highest value of  $u_i$  is considered the best-ranked.

**The final ranking of alternatives based on the MULTIMOORA method.** As a result of evaluation by applying the MULTIMOORA method, three ranking lists of the considered alternatives are obtained. Based on Brauers and Zavadskas (2011), the final ranking order of the alternatives is determined based on the theory of dominance.

#### 4. An Extension of the MULTIMOORA Method Based on Single Valued Neutrosophic Numbers

For an MCDM problem involving  $m$  alternatives and  $n$  criteria, whereby the performances of the alternatives are expressed by using SVNS, the calculation procedure of the extended MULTIMOORA method can be expressed as follows:

##### Step 1. Determine the ranking order of the alternatives based on the RS approach.

The ranking of the alternatives and the selection of the best one based on this approach in the proposed extension of the MULTIMOORA method can be expressed through the following sub steps:

**Step 1.1.** Calculate  $Y_i^+$  and  $Y_i^-$  by using the SVNWA operator, as follows:

$$Y_i^+ = \left( 1 - \prod_{j \in \Omega_{\max}} (1 - t_j)^{w_j}, \prod_{j \in \Omega_{\max}} (i_j)^{w_j}, \prod_{j \in \Omega_{\max}} (f_j)^{w_j} \right), \quad (20)$$

$$Y_i^- = \left( 1 - \prod_{j \in \Omega_{\min}} (1 - t_j)^{w_j}, \prod_{j \in \Omega_{\min}} (i_j)^{w_j}, \prod_{j \in \Omega_{\min}} (f_j)^{w_j} \right), \quad (21)$$

where:  $Y_i^+$  and  $Y_i^-$  denote the importance of the alternative  $i$  obtained based on the benefit and cost criteria, respectively;  $Y_i^+$  and  $Y_i^-$  are SVNNS.

**Step 1.2.** Calculate  $y_i^+$  and  $y_i^-$  by using the Score Function, as follows:

$$y_i^+ = s(Y_i^+), \quad (22)$$

$$y_i^- = s(Y_i^-). \quad (23)$$

**Step 1.3.** Calculate the overall importance for each alternative, as follows:

$$y_i = y_i^+ - y_i^-. \quad (24)$$

**Step 1.4.** Rank the alternatives and select the best one. The ranking of the alternatives can be performed in the same way as in the RS approach of the ordinary MULTIMOORA method.

**Step 2. Determine the ranking order of the alternatives based on the RP approach.**

The ranking of the alternatives and the selection of the best one, based on the RP approach, can be expressed through the following substeps:

**Step 2.1. Determine the reference point.** In this approach, each coordinate of the reference point  $r^* = \{r_1^*, r_2^*, \dots, r_n^*\}$  is an SVNN,  $r_j^* = \langle t_j^*, i_j^*, f_j^* \rangle$ , whose values are determined as follows:

$$r_j^* = \begin{cases} \langle \max_i t_{ij}, \min_i i_{ij}, \min_i f_{ij} \rangle, & j \in \Omega_{\max}, \\ \langle \min_i t_{ij}, \min_i i_{ij}, \max_i f_{ij} \rangle, & j \in \Omega_{\min}, \end{cases} \quad (25)$$

where:  $r_j^*$  denotes the coordinate  $j$  of the reference point.

For the sake of simplicity,  $r_j^*$  could be determined as follows:

$$r_j^* = \begin{cases} \langle 1, 0, 0 \rangle, & j \in \Omega_{\max}, \\ \langle 0, 0, 1 \rangle, & j \in \Omega_{\min}. \end{cases} \quad (25a)$$

**Step 2.2.** Determine the maximum distance from each alternative to all the coordinates of the reference point as follows:

$$d_{ij}^{\max} = d_{\max}(r_{ij}, r_j^*)w_j, \quad (26)$$

where  $d_{ij}^{\max}$  denotes the maximum distance of the alternative  $i$  obtained based on the criterion  $j$  determined by Eq. (8).

**Step 2.3.** Determine the maximum distance of each alternative, as follows:

$$d_i^{\max} = \max_j d_{ij}^{\max}. \quad (27)$$

**Step 2.4.** Rank the alternatives and select the best one. At this step, the ranking of the alternatives can be done in the same way as in the RPA of the ordinary MULTIMOORA method.

**Step 3. Determine the ranking order of the alternatives and select the best one based on the FMF.** The ranking of the alternatives and the selection of the best one can be expressed through the following sub steps:

**Step 3.1.** Calculate  $A_i$  and  $B_i$  as follows:

$$A_i = \left( \prod_{j \in \Omega_{\max}} (t_j)^{w_j}, 1 - \prod_{j \in \Omega_{\max}} (1 - i_j)^{w_j}, 1 - \prod_{j \in \Omega_{\max}} (1 - f_j)^{w_j} \right), \quad (28)$$

$$B_i = \left( \prod_{j \in \Omega_{\min}} (t_j)^{w_j}, 1 - \prod_{j \in \Omega_{\min}} (1 - i_j)^{w_j}, 1 - \prod_{j \in \Omega_{\min}} (1 - f_j)^{w_j} \right), \quad (29)$$

where:  $A_i = \langle t_{Ai}, i_{Ai}, f_{Ai} \rangle$  and  $B_i = \langle t_{Bi}, i_{Bi}, f_{Bi} \rangle$  are SVNNs.

**Step 3.2.** Calculate  $a_i$  and  $b_i$  by using the Score Function as follows:

$$a_i = s(A_i), \quad (30)$$

$$b_i = s(B_i). \quad (31)$$

**Step 3.3.** Determine the overall utility for each alternative as follows:

$$u_i = \frac{a_i}{b_i}. \quad (32)$$

**Step 3.4.** Rank the alternatives and select the best one. The ranking of the alternatives can be performed in the same way as in the FMF of the ordinary MULTIMOORA method.

**Step 4. Determine the final ranking order of the alternatives.** The final ranking order of the alternatives can be determined as in the case of the ordinary MULTIMOORA method, i.e. based on the dominance theory.

## 5. A Numerical Example

In order to demonstrate the applicability and efficiency of the proposed approach, an example has been adopted from Stanujkic *et al.* (2015). In order to briefly demonstrate the advantages of the proposed methodology, this example has been slightly modified.

Suppose that a mining and smelting company has to build a new flotation plant, for which reason an expert has been engaged to evaluate the three Comminution Circuit Designs (CCDs) listed below:

- $A_1$ , the CCDs based on the combined use of rod mills and ball mills;
- $A_2$ , the CCDs based on the use of ball mills; and
- $A_3$ , the CCDs based on the use of semi-autogenous mills.

For the purpose of conducting an evaluation, the following criteria have been chosen:

- $C_1$ , Grinding efficiency;
- $C_2$ , Economic efficiency;
- $C_3$ , Technological reliability;
- $C_4$ , Capital investment costs; and
- $C_5$ , Environmental impact.

The ratings obtained from the expert are shown in Table 1.

**The ranking based on the RS approach.** The ranking results and the ranking order of the alternatives obtained based on the RS approach, i.e. by applying Eqs. (19) to (23), are accounted for in Table 2.

**The ranking based on the RPA.** The ranking of the alternatives based on the RP approach begins by determining the reference point, as it is shown in Table 3.



Table 1  
The ratings of the three generic CCDs obtained from an expert.

	$C_1$	$C_2$	$C_3$	$C_4$	$C_5$
$A_1$	$\langle 0.9, 0.1, 0.2 \rangle$	$\langle 0.7, 0.2, 0.3 \rangle$	$\langle 0.9, 0.1, 0.2 \rangle$	$\langle 0.9, 0.1, 0.2 \rangle$	$\langle 0.9, 0.1, 0.2 \rangle$
$A_2$	$\langle 0.8, 0.1, 0.3 \rangle$	$\langle 0.8, 0.1, 0.3 \rangle$	$\langle 0.8, 0.1, 0.3 \rangle$	$\langle 0.9, 0.1, 0.2 \rangle$	$\langle 0.8, 0.1, 0.3 \rangle$
$A_3$	$\langle 1.0, 0.1, 0.3 \rangle$	$\langle 0.9, 0.1, 0.2 \rangle$	$\langle 0.9, 0.1, 0.2 \rangle$	$\langle 0.7, 0.2, 0.5 \rangle$	$\langle 0.7, 0.2, 0.3 \rangle$

Table 2  
The ranking orders of the alternatives obtained on the basis of the RS approach.

	$Y_i^+$	$Y_i^-$	$y_i^+$	$y_i^-$	$y_i$	Rank
$A_1$	$\langle 0.73, 0.25, 0.38 \rangle$	$\langle 0.55, 0.45, 0.57 \rangle$	0.425	0.045	0.380	2
$A_2$	$\langle 0.65, 0.22, 0.46 \rangle$	$\langle 0.51, 0.45, 0.60 \rangle$	0.372	0.006	0.366	3
$A_3$	$\langle 1.0, 0.22, 0.39 \rangle$	$\langle 0.34, 0.57, 0.73 \rangle$	0.583	-0.263	0.845	1

Table 3  
The reference point.

	$C_1$	$C_2$	$C_3$	$C_4$	$C_5$
$r_j^*$	$\langle 1.0, 0.1, 0.3 \rangle$	$\langle 0.9, 0.2, 0.3 \rangle$	$\langle 0.9, 0.1, 0.3 \rangle$	$\langle 0.7, 0.1, 0.2 \rangle$	$\langle 0.7, 0.1, 0.2 \rangle$

Table 4  
The ranking order of the alternatives obtained based on the RP approach.

I	II	III	IV	V	VI	VII	VI
	$r_1^*$	$r_2^*$	$r_3^*$	$r_4^*$	$r_5^*$	$d_i^{\max}$	Rank
$A_1$	0.02	0.03	0.00	0.00	0.00	0.034	1
$A_2$	0.05	0.02	0.02	0.00	0.01	0.048	2
$A_3$	0.00	0.00	0.00	0.06	0.01	0.063	3

Table 5  
The ranking order of the alternatives obtained based on the FMF.

	$A_i$	$B_i$	$a_i$	$b_i$	$u_i$	Rank
$A_1$	$\langle 0.89, 0.25, 0.15 \rangle$	$\langle 0.96, 0.45, 0.08 \rangle$	0.618	0.498	1.242	3
$A_2$	$\langle 0.86, 0.22, 0.21 \rangle$	$\langle 0.95, 0.45, 0.09 \rangle$	0.605	0.481	1.258	2
$A_3$	$\langle 0.96, 0.22, 0.16 \rangle$	$\langle 0.88, 0.57, 0.18 \rangle$	0.674	0.283	2.379	1

The maximum distances from each alternative to the coordinate  $j$  of the reference point obtained by using Eq. (25) and the maximum distance of each alternative obtained by using Eq. (26) are presented in Table 4. The ranking order of the alternatives is also presented in Table 4.

**The ranking based on the FMF.** The ranking results and the ranking order of the alternatives obtained on the basis of the FMF approach, i.e. by applying Eqs. (27) to (31), are demonstrated in Table 5.

Table 6  
The final ranking order of the alternatives according to the MULTIMOORA method.

	RS	RP	FMF	Rank
$A_1$	2	1	3	3
$A_2$	3	2	2	2
$A_3$	1	3	1	1

The final ranking order of the alternatives which summarizes the three different ranks provided by the respective parts of the MULTIMOORA method is shown in Table 6.

As it can be seen from Table 6, all three approaches, integrated in the MULTIMOORA, have resulted in different ranking orders, for which reason the final ranking order is determined based on the dominance theory.

## 6. Conclusion

The MULTIMOORA method has been proven in solving different decision-making problems. In order to enable its application in the solving of a larger number of complex decision-making problems, numerous extensions have been proposed for the MULTIMOORA method.

Compared to crisp, fuzzy, interval-valued and intuitionistic fuzzy numbers, the neutrosophic set provides significantly greater flexibility, which can be conducive to solving decision-making problems associated with uncertainty, estimations and predictions.

Therefore, an extension of the MULTIMOORA method enabling the use of single valued neutrosophic numbers is proposed in this paper.

The usability and efficiency of the proposed extension is presented in the example of the comminution circuit design selection.

Finally, it should be noted that the proposed extension of the MULTIMOORA method can be used for solving a much larger number of complex decision-making problems. A number of real-world decision making problems which have to be solved is based on the data acquired from respondents can be identified as one of the areas where the proposed extension of the MULTIMOORA method can reach its advantages.

## References

- Atanassov, K.T. (1986). Intuitionistic fuzzy sets. *Fuzzy Sets and Systems*, 20(1), 87–96.
- Atanassov, K., Gargov, G. (1989). Interval valued intuitionistic fuzzy sets. *Fuzzy Sets and Systems*, 31(3), 343–349.
- Balezentis, T., Zeng, S. (2013). Group multi-criteria decision making based upon interval-valued fuzzy numbers: an extension of the MULTIMOORA method. *Expert Systems with Applications*, 40(2), 543–550.
- Balezentis, A., Balezentis, T., Brauers, W.K.M. (2012a). MULTIMOORA-FG: a multi-objective decision making method for linguistic reasoning with an application to personnel selection. *Informatica*, 23(2), 173–190.
- Balezentis, A., Balezentis, T., Brauers, W.K.M. (2012b). Personnel selection based on computing with words and fuzzy MULTIMOORA. *Expert Systems with applications*, 39(9), 7961–7967.

- Balezentis, T., Zeng, S., Balezentis, A. (2014). MULTIMOORA-IFN: a MCDM method based on intuitionistic fuzzy number for performance management. *Economic Computation & Economic Cybernetics Studies & Research*, 48(4), 85–102.
- Brauers, W.K.M. (2010). The economy of the Belgian regions tested with MULTIMOORA. *Journal of Business Economics and Management*, 11(2), 173–209.
- Brauers, W.K.M., Zavadskas, E.K. (2010). Project management by MULTIMOORA as an instrument for transition economies. *Technological and Economic Development of Economy*, 16(1), 5–24.
- Brauers, W.K.M., Zavadskas, E.K. (2011). MULTIMOORA optimization used to decide on a bank loan to buy property. *Technological and Economic Development of Economy*, 17(1), 174–188.
- Brauers, W.K., Balezentis, A., Balezentis, T. (2011). MULTIMOORA for the EU Member States updated with fuzzy number theory. *Technological and Economic Development of Economy*, 17(2), 259–290.
- Hafezalkotob, A., Hafezalkotob, A. (2016). Extended MULTIMOORA method based on Shannon entropy weight for materials selection. *Journal of Industrial Engineering International*, 12(1), 1–13.
- Hafezalkotob, A., Hafezalkotob, A., Sayadi, M.K. (2016). Extension of MULTIMOORA method with interval numbers: an application in materials selection. *Applied Mathematical Modelling*, 40(2), 1372–1386.
- Kracka, M., Brauers, W.K.M., Zavadskas, E.K. (2015). Ranking heating losses in a building by applying the MULTIMOORA. *Engineering Economics*, 21(4).
- Li, Y., Liu, P., Chen, Y. (2016). Some single valued neutrosophic number heronian mean operators and their application in multiple attribute group decision making. *Informatica*, 27(1), 85–110.
- Liu, H.C., Fan, X.J., Li, P., Chen, Y.Z. (2014a). Evaluating the risk of failure modes with extended MULTIMOORA method under fuzzy environment. *Engineering Applications of Artificial Intelligence*, 34, 168–177.
- Liu, H.C., You, J.X., Lu, C., Shan, M.M. (2014b). Application of interval 2-tuple linguistic MULTIMOORA method for health-care waste treatment technology evaluation and selection. *Waste Management*, 34(11), 2355–2364.
- Sahin, R. (2014). Multi-criteria neutrosophic decision making method based on score and accuracy functions under neutrosophic environment. arXiv:1412.5202.
- Sahu, A.K., Sahu, N.K., Sahu, A.K. (2016). Application of modified MULTI-MOORA for CNC machine tool evaluation in IVGTFNS environment: an empirical study. *International Journal of Computer Aided Engineering and Technology*, 8(3), 234–259.
- Smarandache, F. (1998). *Neutrosophy Probability Set and Logic*. American Research Press, Rehoboth.
- Smarandache, F. (1999). *A Unifying Field in Logics. Neutrosophy: Neutrosophic Probability, Set and Logic*. American Research Press, Rehoboth.
- Stanujkic, D., Zavadskas, E.K., Brauers, W.K.M., Karabasevic, D. (2015). An extension of the MULTIMOORA method for solving complex decision-making problems based on the use of interval-valued triangular fuzzy numbers. *Transformations in Business and Economics* 14(2B(35B)), 355–377.
- Turksen, I.B. (1986). Interval valued fuzzy sets based on normal forms. *Fuzzy Sets and Systems*, 20(2), 191–210.
- Wang, H., Smarandache F., Zhang, Y.Q., Sunderraman, R. (2010). Single valued neutrosophic sets. *Multispace and Multistructure*, 4, 410–413.
- Zadeh, L.A. (1965). Fuzzy sets. *Information and Control*, 8(3), 338–353.
- Zavadskas, E.K., Antucheviciene, J., Razavi Hajiagha, S.H., Hashemi, S.S. (2015). The interval-valued intuitionistic fuzzy MULTIMOORA method for group decision making in engineering. *Mathematical Problems in Engineering*, 1–13, Article ID 560690.

**D. Stanujkic** is a full-time professor of Information Technology and Decision Sciences at the Faculty of Management in Zajecar, John Naisbitt University, Belgrade. He has received his MSc degree in Information Science and PhD in Organizational Sciences from the Faculty of Organizational Sciences, University of Belgrade. His current research is focused on decision-making theory, expert systems and intelligent decision support systems.

**E.K. Zavadskas** is a professor, head of the Department of Construction Technology and Management at Vilnius Gediminas Technical University, Vilnius, Lithuania, and senior researcher at Research Institute of Internet and Intelligent Technologies. He has a PhD in building structures (1973) and DrSc (1987) in building technology and management. He is a member of the Lithuanian and several foreign Academies of Sciences. He is doctor honoris causa at Poznan, Saint Petersburg, and Kiev universities. He is the editor in chief and a member of editorial boards of a number of research journals. He is an author and coauthor of more than 400 papers and a number of monographs. Research interests are: building technology and management, decision-making theory, automation in design and decision support systems.

**F. Smarandache** is a professor of mathematics at the University of New Mexico, USA. He has published many papers and books on neutrosophic set and logic and their applications and has presented to many international conferences. He got his MSc in Mathematics and Computer Science from the University of Craiova, Romania, PhD from the State University of Kishinev, and Post-Doctoral in Applied Mathematics from Okayama University of Sciences, Japan.

**W.K.M. Brauers**, doctor honoris causa Vilnius Gediminas Technical University, was graduated as: PhD in economics (un. of Leuven), master of arts (in economics) of Columbia University (New York), master in economics, master in management and financial sciences, master in political and diplomatic sciences and bachelor in philosophy (all in the University of Leuven). He is professor ordinarius at the Faculty of Applied Economics of the University of Antwerp, honorary professor at the University of Leuven, the Belgian War College, the School of Military Administrators and the Antwerp Business School. His scientific publications consist of eighteen books and several hundreds of articles and reports in English, Dutch and French.

**D. Karabasevic** is an assistant professor at the Faculty of Applied Management, Economics and Finance, Business Academy University. He obtained his degrees at all the levels of studies (BSc appl. in Economics, BSc in Economics, Academic Specialization in the Management of Business Information Systems and PhD in Management and Business) at the Faculty of Management in Zajecar, John Naisbitt University, Belgrade. His current research is focused on the human resource management, management and decision-making theory.