



Some Topological Character of Neutrosophic normed spaces

Vakeel A. Khan¹, Mohammad Daud Khan²

¹ Department of Mathematics, Aligarh Muslim University, Aligarh-202002, India.

Email addresses: vakhanmaths@gmail.com

² Department of Mathematics, Aligarh Muslim University, Aligarh-202002, India.

Email addresses: mhddaudkhan2@gmail.com

Abstract. The concept of neutrosophic normed spaces was introduced by Murat Kirisci* and Necip Simsek [3]. In this paper, by using the compact operator $\psi(z_j)$ and the concept of statistical convergence. we introduce some neutrosophic statistical convergence of sequence spaces defined by compact operator and study of neutrosophic norm and derive the relation between statistical convergence in NNS with the help of compact operator. We also focus on some topological and algebraic properties of these convergent sequence spaces.

Keywords: NNS; t-norm; t-conorm; Statistical convergence; compact linear operator; Bounded linear operator; neutrosophic set.

1. Introduction

The notion of fuzzy set was presented by Zadeh [4], since then several kinds of research have appeared and fuzzication of many classical theories has also been made. Fuzzy sets hypothesis is an amazing handset for demonstrating vulnerability and dubiousness in different problems and issues emerging in field of science and designing. fuzzy topology is perhaps the absolute significant and generally utilized outfits and it ends up being valuable for managing such circumstances where utilization of old style speculations breakdown. [13] he used the concept of fuzzy sets and examine the fuzzy real valued p-absolutely summable multiple sequences in probabilistic normed spaces. Atanassov [29] generalized the fuzzy sets theory and studied the concepts of intuitionistic fuzzy sets (*IFS*). In 2004, [6] Park researched the idea of (*IFS*), further Saadati and Park analyzed this concept in the norm. The idea of *IFNS* and the thought of statistical convergence is a valuable utilitarian apparatus for contemplating the convergence issues of mathematical problems (double sequence) herewith the idea of thickness. [14]. The idea of neutrosophic sets (*NS*) was introduced by Smarandache [23]. This set is an expansion of *IFS* regardless, if the summation of neutrosophic segments is < 1 , or > 1 , or $= 1$. For the situation when the aggregate of the components is 1 (*as in IFS*), in the wake

of satisfying the condition by applying the neutrosophic set operators, different outcomes can be acquired by applying the intuitionistic fuzzy (IF) operators, since the IF operators disregard the (*indeterminacy*), while the NS operators taken into cognizance of the indeterminacy at a similar level as (*truth – membership*) and (*falsehood – nonmembership*) are taken. [26] using the idea of neutrosophic sets defined the notion of Neutrosophic Bipolar Vague Soft Set and Its Application to Decision Making Problems. Further, Smarandache [9, 10, 12, 35] investigated neutroalgebra which is generalization of partial algebra, neutroalgebraic structures and antialgebraic structures. NS is likewise more adaptable and effective in light of the fact that it handles, aside from autonomous (free) components, additionally partially independent and dependent components, while IFS can't manage these cases. Moreover [38] define neutrosophic fuzzy matrices and Some Algebraic Operations. Smarandache [1, 34] analyzed the conflict between neutrosophic rationale, intuitionistic fuzzy rationale, and the comparing NS and IFS . [11] defined Neutrosophic simply soft open set in neutrosophic soft topological space. Moreover, Bera and Mahapatra [24] introduced the neutrosophic soft linear space. Bera and Mahapatra [25] studied convexity, metric, Cauchy sequence, and neutrosophic soft norm linear space (NSNLS). [15] A lot of developments have been made in this areas after the work of Das, S and Pramanik, S defined the Generalized neutrosophic b-open sets in neutrosophic topological space. Further, [21] examine Neutrosophic Multiset Topological Space. [17] another term find Statistically pre-Cauchy fuzzy real-valued sequences defined by Orlicz function. Later on, the concepts of statistical convergence of double sequences have been analyzed in IFNS by Mursaleen and Mohiuddin [14]. Quite recently, Kirisci and Simsek [3] introduced the notion of NNS and statistical convergence. Further [8] examine On almost statistical convergence of new type of generalized difference sequence of fuzzy numbers. Since NNS is a natural generalization of IFNS and statistical convergence. [19] defined a new concept On pointwise statistical convergence of order alpha of sequences of fuzzy mappings. In this paper we aim to define novel statistical convergence of sequence spaces. sequence spaces using neutrosophic norm and using compact operator as a tool and discussed their topological and algebraic properties. We mention the following notions that will be put to use in the paper further.

2. Preliminaries

:

Definition 2.1. A sequence $z = (z_j)$ is called a δ -convergent to the number ξ for each $\varepsilon > 0$, the set $Y(\varepsilon)$ has δ -density zero, where

$$Y_\varepsilon(F) = \{j \in \mathbb{N} : |z_j - \xi| \geq \varepsilon\} \quad (1)$$

we write $S_\delta - \lim z = \xi$ or $z_j \rightarrow \xi(S_\delta)$.

Definition 2.2. [14] A sequence $z = (z_j)$ is called statistically Cauchy sequences if \exists , a number $\mathcal{K} = \mathcal{K}(\epsilon)$ such that for each $\epsilon > 0$

$$\lim_{j \rightarrow \infty} \frac{1}{j} |\{i \leq j : |z_i - (z_N)| \geq \epsilon\}| = 0. \tag{2}$$

Definition 2.3. [7] Let $X \neq \emptyset$, the \mathcal{W} intuitionistic fuzzy set, $\mathcal{W} \subset X$ is defined by

$$\mathcal{W} = \{ \langle z, \mathbf{T}(z), \mathbf{F}(z) \rangle : z \in X \}, \tag{3}$$

where $\mathbf{T}(z), \mathbf{F}(z) : X \rightarrow [0, 1]$, $\mathbf{T}(z) = (\text{Truth})$ and $\mathbf{F}(z) = (\text{Falsehood})$ respectively.

$$0 \leq \mathbf{T}(z) + \mathbf{F}(z) \leq 1$$

Definition 2.4. [31] Let $X \neq \phi$ and $\mathcal{W} \subset X$ Then,

$$\mathcal{W}_{NS} = \{ \langle z, \mathbf{T}(z), \mathbf{I}(z), \mathbf{F}(z) \rangle : z \in X \},$$

where $\mathbf{T}(z), \mathbf{I}(z), \mathbf{F}(z) : X \rightarrow [0, 1]$, $\mathbf{T}(z) = \text{Truth}$, $\mathbf{I}(z) = \text{Indeterminacy}$, and $\mathbf{F}(z) = \text{Falsehood}$ respectively.

$$0 \leq \mathbf{T}(z) + \mathbf{I}(z) + \mathbf{F}(z) \leq 3.$$

The components of neutrosophic are $\mathbf{T}(z), \mathbf{I}(z)$ and $\mathbf{F}(z)$ independent of each other.

Definition 2.5. [1] Suppose $X \neq \phi$, Q and R are neutrosophic sets in X . Then,

- (a) $Q \subset R \iff \mathbf{T}_Q(z) \leq \mathbf{T}_R(z), \mathbf{I}_Q(z) \leq \mathbf{I}_R(z), \mathbf{F}_Q(z) \geq \mathbf{F}_R(z) \forall z \in X$
- (b) $Q = R \iff \mathbf{T}_Q(z) = \mathbf{T}_R(z), \mathbf{I}_Q(z) = \mathbf{I}_R(z), \mathbf{F}_Q(z) = \mathbf{F}_R(z) \forall z \in X$
- (c) $Q \cap R = \{ \langle z, \min(\mathbf{T}_Q(z), \mathbf{T}_R(z)), \min(\mathbf{I}_Q(z), \mathbf{I}_R(z)), \min(\mathbf{F}_Q(z), \mathbf{F}_R(z)) \rangle \mid z \in X \}$
- (d) $Q \cup R = \{ \langle z, \max(\mathbf{T}_Q(z), \mathbf{T}_R(z)), \max(\mathbf{I}_Q(z), \mathbf{I}_R(z)), \max(\mathbf{F}_Q(z), \mathbf{F}_R(z)) \rangle, z \in X \}$
- (e) $Q^c = \{ \langle z, \mathbf{F}_Q(z), 1 - \mathbf{I}_R(z), \mathbf{T}_Q(z) \rangle \mid z \in X \}$
- (f) $Q \setminus R = \{ \langle z, \mathbf{T}_Q(z) \min \mathbf{F}_R(z), \mathbf{I}_Q(z) \min 1 - \mathbf{I}_R(z), \mathbf{F}_Q(z) \max \mathbf{T}_R(z) \rangle \mid z \in X \}.$

Definition 2.6. [40] Consider a binary operation \diamond on the interval. if satisfying following axioms,

$$\diamond : [0, 1]^2 \longrightarrow [0, 1]$$

- (1) $\hat{\alpha} \diamond \hat{\beta} = \hat{\beta} \diamond \hat{\alpha}$
- (2) $(\hat{\alpha} \diamond \hat{\beta}) \diamond \hat{\lambda} = \hat{\alpha} \diamond (\hat{\beta} \diamond \hat{\lambda})$
- (3) $\hat{\alpha} \diamond 1 = \hat{\alpha} \forall \hat{\alpha} \in [0, 1]$,
- (4) $\hat{\alpha} \diamond \hat{\beta} \leq \hat{\lambda} \diamond \hat{\nu}$ whenever $\hat{\alpha} \leq \hat{\lambda}$ and $\hat{\beta} \leq \hat{\nu}$ for each $\hat{\alpha}, \hat{\beta}, \hat{\lambda}, \hat{\nu} \in [0, 1]$.

is called a continuous t -norm.

where $\wedge = \min$ and $\vee = \max$.

Example 2.7. For $\hat{\alpha}, \hat{\beta} \in [0, 1]$, define $\hat{\alpha} \diamond \hat{\beta} = \hat{\alpha}\hat{\beta}$ or $\hat{\alpha} \diamond \hat{\beta} = \wedge\{\hat{\alpha}, \hat{\beta}\}$. then \diamond is continuous t -norm.

Definition 2.8. [40] Consider a binary operation \star on the interval. if satisfying following axioms,

$$\star : [0, 1]^2 \longrightarrow [0, 1]$$

- (1) $\hat{\alpha} \star \hat{\beta} = \hat{\beta} \star \hat{\alpha}$
- (2) $(\hat{\alpha} \star \hat{\beta}) \star \hat{\lambda} = \hat{\alpha} \star (\hat{\beta} \star \hat{\lambda})$
- (3) $\hat{\alpha} \star 0 = \hat{\alpha} \forall \hat{\alpha} \in [0, 1]$,
- (4) $\hat{\alpha} \star \hat{\beta} \leq \hat{\lambda} \star \hat{\nu}$ whenever $\hat{\alpha} \leq \hat{\lambda}$ and $\hat{\beta} \leq \hat{\nu}$ for each $\hat{\alpha}, \hat{\beta}, \hat{\lambda}, \hat{\nu} \in [0, 1]$.

is called a continuous t -conorm.

where $\wedge = \min$ and $\vee = \max$.

Example 2.9. Consider $\hat{\alpha}, \hat{\beta} \in [0, 1]$. define $\hat{\alpha} \star \hat{\beta} = \wedge\{\hat{\alpha} + \hat{\beta}, 1\}$ or $\hat{\alpha} \star \hat{\beta} = \vee\{\hat{\alpha}, \hat{\beta}\}$. Then \star is continuous t -conorm

Definition 2.10. [3] Take X as a vector space and $\mathcal{M} = \{ \langle x, \mathbf{T}(z), \mathbf{I}(z), \mathbf{F}(z) \rangle : z \in X \}$ be a normed space such that $\mathbf{T}(z), \mathbf{I}(z), \mathbf{F}(z) : X \times R^+ \rightarrow [0, 1]$. Assume \diamond show the continuous t -norm and \star be a continuous t -conorm respectively, then $\mathcal{V} = (X, \mathcal{M}, \diamond, \star)$ is called (NNS) . if the subsequent terms holds; $\forall z, a \in X$ and $s, k > 0$

- (i) $0 \leq \mathbf{T}(z, s) \leq 1, 0 \leq \mathbf{I}(z, s) \leq 1, 0 \leq \mathbf{F}(z, s) \leq 1, s \in R^+$,
- (ii) $\mathbf{T}(z, s) + \mathbf{I}(z, s) + \mathbf{F}(z, s) \leq 3$, for $s \in R^+$,
- (iii) $\mathbf{T}(z, s) = 1$ for $s > 0$ iff $z = 0$
- (iv) $\mathbf{T}(\alpha x, s) = \mathbf{T}(z, \frac{s}{|\alpha|})$,
- (v) $\mathbf{T}(z, s) \diamond \mathbf{T}(a, s) \leq \mathbf{T}(z + a, s + k)$,
- (vi) $\mathbf{T}(z, \diamond)$ is continuous non-decreasing function

$$(vii) \lim_{s \rightarrow \infty} \mathbf{T}(z, s) = 1$$

$$(viii) \mathbf{I}(z, s) = 0 \text{ for } s > 0 \text{ iff } z = 0$$

$$(ix) \mathbf{I}(\alpha z, s) = \mathbf{I}(y, \frac{s}{|\alpha|}), \text{ for each } \alpha \neq 0,$$

$$(x) \mathbf{I}(z, s) \star \mathbf{I}(a, k) \geq \mathbf{I}(z + a, s + k),$$

$$(xi) \mathbf{I}(z, \star) \text{ is continuous non-increasing function,}$$

$$(xii) \lim_{s \rightarrow \infty} \mathbf{I}(z, s) = 0,$$

$$(xiii) \mathbf{F}(z, s) = 0 \text{ for } s > 0 \text{ iff } z = 0$$

$$(xiv) \mathbf{F}(\alpha z, s) = \mathbf{F}(z, \frac{s}{|\alpha|}), \text{ for each } \alpha \neq 0,$$

$$(xv) \mathbf{F}(z, s) \star \mathbf{F}(z, k) \geq \mathbf{F}(z + a, s + k),$$

$$(xvi) \mathbf{F}(z, \cdot) \text{ is continuous non-increasing function,}$$

$$(xvii) \lim_{s \rightarrow \infty} \mathbf{F}(z, s) = 0,$$

$$(xviii) \text{ If } s \leq 0, \text{ then } \mathbf{T}(z, s) = 0, \mathbf{I}(z, s) = 1, \mathbf{F}(z, s) = 1.$$

In this case $\mathcal{M} = (\mathbf{T}, \mathbf{I}, \mathbf{F})$ is said to be neutrosophic normed (*NNS*).

Example 2.11. [3] Suppose $(X, \|\cdot\|)$ be a NNS. Give the operations as $TC \ z \diamond a = z + a - za$ and $TN \ z \star a = \min(z, a)$. For $s > \|z\|$,

$$\mathbf{T}_0(z, s) = \frac{s}{s + \|z\|}, \mathbf{I}_0(z, s) = \frac{\|z\|}{s + \|z\|}, \mathbf{F}_0(z, s) = \frac{\|z\|}{s}. \quad (4)$$

for all $z, a \in [0, 1]$ and $s > 0$. If we take $s \leq \|z\|$, and let $\mathbf{T}_0, \mathbf{I}_0$ and \mathbf{F}_0 be neutrosophic sets on $X \times (0, \infty)$ then

$$\mathbf{T}_0(z, s) = 0, \mathbf{I}_0(z, s) = 1 \text{ and } \mathbf{F}_0(z, s) = 1.$$

Then, $(X, \mathcal{M}, \diamond, \star)$ is neutrosophic normed space such that $\mathcal{M} : X \times R^+ \rightarrow [0, 1]$.

Vakeel A. Khan, Mohammad Daud Khan, Some Topological Character of Neutrosophic normed spaces

3. Main Results

In this article, we examine the algebraic and topological properties on the space of neutrosophic statistical convergence given by :

Definition 3.1. [3] Suppose $(X, \mathcal{M}, \star, \diamond)$ be a *NNS* and $z = (z_j)$ is called a Cauchy sequence with respect to \mathcal{M} , if for each $\epsilon > 0$ and $s > 0 \exists, u \in \mathbb{N}$ such that $\mathbf{T}(z_j - a_k, s) > 1 - \epsilon$, $\mathbf{I}(z_j - a_k, s) < \epsilon$ and $\mathbf{F}(z_j - a_k, s) < \epsilon$ for all $j, k \geq u$.

Definition 3.2. Let X be a *NNS*, the sequence $z = (z_j)$ in X is called convergent at $\xi \in X$ iff $\exists, \mathring{N} \in \mathbb{N}$, with respect *NN* ($\mathbf{T}, \mathbf{I}, \mathbf{F}$) if for every $\epsilon > 0, s > 0$

$$\mathbf{T}(\psi(z_j) - \xi, s) > 1 - \epsilon, \mathbf{I}(\psi(z_j) - \xi, s) < \epsilon \text{ and } \mathbf{F}(\psi(z_j) - \xi, s) < \epsilon \quad (5)$$

for all $j \geq \mathring{N}$, i.e.,

$$\lim_{j \rightarrow \infty} \mathbf{T}(\psi(z_j) - \xi, s) = 1, \lim_{j \rightarrow \infty} \mathbf{I}(\psi(z_j) - \xi, s) = 0 \text{ and } \lim_{j \rightarrow \infty} \mathbf{F}(\psi(z_j) - \xi, s) = 0.$$

In such case, we denote $\mathcal{M} - \lim z_j = \xi$.

Definition 3.3. (See [2]). Suppose A and B be two normed linear spaces (*NLS*) and $\psi : \mathcal{D} \rightarrow B$ be a linear operator, where $\mathcal{D}(\psi) \subset A$ Then, the operator ψ is called a bounded, if \exists , a +ve real no c' such that

$$\|\psi z\| \leq c' \|z\|, \forall z \in \mathcal{D}(\psi).$$

The set of all bounded linear operators $\mathbf{B}(A, B)$ [2] is a (*NLS*) normed by

$$\|\psi\| = \sup_{z \in A, \|z\|=1} \|\psi z\|$$

and $\mathbf{B}(A, B)$ is a Banach space if B is a Banach space.

Definition 3.4. Let us Consider two *NLS* A and B

$$\psi : A \rightarrow B$$

is called a compact linear operator, if

(i) ψ is linear

(ii) ψ maps every bounded sequence (z_j) in A on to a sequence $\psi(z_j)$ in B which has a convergent subsequence.

The set of all compact linear operators $\mathcal{C}(A, B)$ is a closed subspace of $\mathbf{B}(A, B)$ and $\mathcal{C}(A, B)$ is Banach space, if B is a Banach space.

Inspire by this [5], we proposed the previous sequence spaces with the assist of compact operator in NNS :

$$\mathcal{S}_{(\mathbf{T}, \mathbf{I}, \mathbf{F})}^N(\psi) = \{(z_j) \in \ell_\infty : \exists k \in \mathbb{N}, \forall j \geq k, \mathbf{T}(\psi(z_j) - \xi, s) > 1 - \epsilon \text{ or } \mathbf{I}(\psi(z_j) - \xi, s) \geq \epsilon \\ \mathbf{F}(\psi(z_j) - \xi, s) \geq \epsilon\}.$$

$$\mathcal{S}_{0(\mathbf{T}, \mathbf{I}, \mathbf{F})}^N(\psi) = \{(z_j) \in \ell_\infty : \exists k \in \mathbb{N}, \forall j \geq k, \mathbf{T}(\psi(z_j) - \xi, s) > 1 - \epsilon \text{ or } \mathbf{I}(\psi(z_j) - \xi, s) \geq \epsilon \\ \mathbf{F}(\psi(z_j) - \xi, s) \geq \epsilon\}.$$

Definition 3.5. [5] Let $\mathcal{V} = (X, M, \diamond, \star)$ is a NNS. For $s > 0$. we define a open ball $(OB) \mathcal{B}(z, r, s)(\psi)$ with centre $z \in X$ and raduis $0 < r < 1$, there exist $k \in \mathbb{N}$ such that for all $j \geq k$

$$\mathcal{B}(z, r, s)(\psi) = \{(a_j) \in \ell_\infty : \mathbf{T}(\psi(z_j) - \psi(a_k), s) < 1 - \epsilon \text{ or } \mathbf{I}(\psi(z_j) - \psi(a_k), s) < \epsilon, \mathbf{F}(\psi(z_j) - \psi(a_k), s) < \epsilon\}.$$

Presently, we are prepared to state and demonstrate our primary outcomes. This hypothesis depends on the linearity of new characterize sequence spaces which is expressed as follows.

Theorem 3.6. $\mathcal{S}_{(\mathbf{T}, \mathbf{I}, \mathbf{F})}^N(\psi)$ and $\mathcal{S}_{0(\mathbf{T}, \mathbf{I}, \mathbf{F})}^N(\psi)$ are linear spaces.

Proof. We shall prove the result for $\mathcal{S}_{(\mathbf{T}, \mathbf{I}, \mathbf{F})}^N(\psi)$ the proof for the different space will take accordingly, let $z = (z_j), a = (a_j) \in \mathcal{S}_{(\mathbf{T}, \mathbf{I}, \mathbf{F})}^N(\psi)$ and \hbar, η be scalars then for a given $\epsilon > 0$, we have

$$\mathcal{W}_1 = \left\{ j \in \mathbb{N} : \mathbf{T}\left(\psi(z_j) - \xi_1, \frac{s}{2|\hbar|}\right) \leq 1 - \epsilon \text{ or } \mathbf{I}\left(\psi(z_j) - \xi_1, \frac{s}{2|\hbar|}\right) \geq \epsilon, \mathbf{F}\left(\psi(z_j) - \xi_1, \frac{s}{2|\hbar|}\right) \geq \epsilon \right\}$$

$$\mathcal{W}_2 = \left\{ j \in \mathbb{N} : \mathbf{T}\left(\psi(a_j) - \xi_2, \frac{s}{2|\eta|}\right) \leq 1 - \epsilon \text{ or } \mathbf{I}\left(\psi(a_j) - \xi_2, \frac{s}{2|\eta|}\right) \geq \epsilon, \mathbf{F}\left(\psi(a_j) - \xi_2, \frac{s}{2|\eta|}\right) \geq \epsilon \right\}$$

$$\mathcal{W}_1^c = \left\{ j \in \mathbb{N} : \mathbf{T}\left(\psi(z_j) - \xi_1, \frac{s}{2|\hbar|}\right) > 1 - \epsilon \text{ or } \mathbf{I}\left(\psi(z_j) - \xi_1, \frac{s}{2|\hbar|}\right) < \epsilon, \mathbf{F}\left(\psi(z_j) - \xi_1, \frac{s}{2|\hbar|}\right) < \epsilon \right\}$$

$$\mathcal{W}_2^c = \left\{ j \in \mathbb{N} : \mathbf{T}\left(\psi(a_j) - \xi_2, \frac{s}{2|\eta|}\right) > 1 - \epsilon \text{ or } \mathbf{I}\left(\psi(a_j) - \xi_2, \frac{s}{2|\eta|}\right) < \epsilon, \mathbf{F}\left(\psi(a_j) - \xi_2, \frac{s}{2|\eta|}\right) < \epsilon \right\}$$

Vakeel A. Khan, Mohammad Daud Khan, Some Topological Character of Neutrosophic normed spaces

Define the set $\mathcal{W}_3 = \mathcal{W}_1 \cup \mathcal{W}_2$ so that $\mathcal{W}_3 \in \mathcal{M}$. It follow that $\mathcal{W}_3^c \neq \phi$. we shall show that for each $(z_j), (a_j) \in \mathcal{S}_{(\mathbf{T}, \mathbf{I}, \mathbf{F})}^N(\psi)$.

$$\mathcal{W}_3^c \subset \left\{ j \in \mathbb{N} : \mathbf{T}\left((\hbar\psi(z_j) + \eta\psi(a_j)) - (\hbar\xi_1 + \eta\xi_2), s\right) > 1 - \epsilon \text{ or } \mathbf{I}\left((\hbar\psi(z_j) + \eta\psi(a_j)) - (\hbar\xi_1 + \eta\xi_2), s\right) < \epsilon, \mathbf{F}\left((\hbar\psi(z_j) + \eta\psi(a_j)) - (\hbar\xi_1 + \eta\xi_2), s\right) < \epsilon \right\}$$

let $m \in \mathcal{W}_3^c$. In this case,

$$\mathbf{T}\left(\psi(z_m) - \xi_1, \frac{s}{2|\hbar|}\right) > 1 - \epsilon \text{ or } \mathbf{I}\left(\psi(z_m) - \xi_1, \frac{s}{2|\hbar|}\right) < \epsilon, \mathbf{F}\left(\psi(z_m) - \xi_1, \frac{s}{2|\hbar|}\right) < \epsilon$$

and

$$\mathbf{T}\left(\psi(a_m) - \xi_2, \frac{s}{2|\eta|}\right) > 1 - \epsilon \text{ or } \mathbf{I}\left(\psi(a_m) - \xi_2, \frac{s}{2|\eta|}\right) < \epsilon, \mathbf{F}\left(\psi(a_m) - \xi_2, \frac{s}{2|\eta|}\right) < \epsilon.$$

We have

$$\begin{aligned} & \mathbf{T}\left((\hbar\psi(z_m) + \eta\psi(a_m)) - (\hbar\xi_1 + \eta\xi_2), s\right) \\ & \geq \mathbf{T}\left(\hbar\psi(z_m) - \hbar\xi_1, \frac{t}{2}\right) \diamond \mathbf{T}\left(\eta\psi(z_m) - \eta\xi_2, \frac{t}{2}\right) \\ & = \mathbf{T}\left(\psi(z_m) - \xi_1, \frac{s}{2|\hbar|}\right) \star \mathbf{T}\left(\psi(z_m) - \xi_2, \frac{s}{2|\eta|}\right) \\ & > (1 - \epsilon) \diamond (1 - \epsilon) = 1 - \epsilon. \end{aligned}$$

In similar way,

$$\begin{aligned} & \mathbf{I}\left((\hbar\psi(z_m) + \eta\psi(a_m)) - (\hbar\xi_1 + \eta\xi_2), s\right) \\ & \leq \mathbf{I}\left(\hbar\psi(z_m) - \hbar\xi_1, \frac{s}{2}\right) \star \mathbf{I}\left(\eta\psi(z_m) - \eta\xi_2, \frac{s}{2}\right) \end{aligned}$$

$$\begin{aligned}
 &= \mathbf{I}\left(\psi(z_m) - \xi_1, \frac{s}{2|\hbar|}\right) \star \mathbf{I}\left(\psi(z_m) - \xi_2, \frac{s}{2|\eta|}\right) \\
 &< \epsilon \star \epsilon < \epsilon.
 \end{aligned}$$

and,

$$\begin{aligned}
 &\mathbf{F}\left((\hbar\psi(z_m) + \eta\psi(a_m)) - (\hbar\xi_1 + \eta\xi_2), s\right) \\
 &\leq \mathbf{F}\left(\hbar\psi(z_m) - \hbar\xi_1, \frac{s}{2}\right) \star \mathbf{F}\left(\eta\psi(z_m) - \eta\xi_2, \frac{s}{2}\right) \\
 &= \mathbf{F}\left(\psi(z_m) - \xi_1, \frac{s}{2|\hbar|}\right) \star \mathbf{F}\left(\psi(z_m) - \xi_2, \frac{s}{2|\eta|}\right) \\
 &< \epsilon \star \epsilon < \epsilon.
 \end{aligned}$$

This implies that,

$$\mathcal{W}_3^c \subset \left\{ j \in \mathbb{N} : \mathbf{T}\left((\hbar\psi(z_j) + \eta\psi(a_j)) - (\hbar\xi_1 + \eta\xi_2), s\right) > 1 - \epsilon \text{ or } \mathbf{I}\left((\hbar\psi(z_j) + \eta\psi(a_j)) - (\hbar\xi_1 + \eta\xi_2), s\right) < \epsilon, \mathbf{F}\left((\hbar\psi(z_j) + \eta\psi(a_j)) - (\hbar\xi_1 + \eta\xi_2), s\right) < \epsilon \right\}.$$

Hence $\mathcal{S}_{(\mathbf{T},\mathbf{I},\mathbf{F})}^N(\psi)$ is a linear space. \square

Remark 3.7. $\mathcal{S}_{(\mathbf{T},\mathbf{I},\mathbf{F})}^N(\psi)$ is an NNS.

Define $\mathcal{T}_{(\mathbf{T},\mathbf{I},\mathbf{F})}^N(\psi) = \{W \subset \mathcal{S}_{(\mathbf{T},\mathbf{I},\mathbf{F})}^N(\psi) : \text{for every } z \in W, \exists, s > 0 \text{ and } r \in (0, 1) \text{ such that } \mathcal{B}(z, r, s)(\psi) \subset W\}$.

Theorem 3.8. *The topology $\mathcal{T}_{(\mathbf{T},\mathbf{I},\mathbf{F})}^N(\psi)$ on $\mathcal{S}_{(\mathbf{T},\mathbf{I},\mathbf{F})}^N(\psi)$ is first countable.*

Proof. $\{B_x(\frac{1}{n}, \frac{1}{n}) : n = 1, 2, 3, \dots\}$ is a local base at x , $\mathcal{T}_{(\mathbf{T},\mathbf{I},\mathbf{F})}^N(\psi)$ on $\mathcal{S}_{(\mathbf{T},\mathbf{I},\mathbf{F})}^N(\psi)$ is first countable. $\mathcal{S}_{(\mathbf{T},\mathbf{I},\mathbf{F})}^N(\psi)$ is Hausdorff. \square

Theorem 3.9. *$\mathcal{S}_{(\mathbf{T},\mathbf{I},\mathbf{F})}^N(\psi)$ is an NNS and $\mathcal{T}_{(\mathbf{T},\mathbf{I},\mathbf{F})}^N(\psi)$ is a topology on $\mathcal{S}_{(\mathbf{T},\mathbf{I},\mathbf{F})}^N(\psi)$ Then a sequence $(z_j) \in \mathcal{S}_{(\mathbf{T},\mathbf{I},\mathbf{F})}^N(\psi)$, $(z_j) \rightarrow z$, if and only if $\mathbf{T}((\psi(z_j) - z), t) \rightarrow 1$, $\mathbf{I}((\psi(z_j) - z), s) \rightarrow 1$ and $\mathbf{F}((\psi(z_j) - z), s) \rightarrow 1$ as $j \rightarrow \infty$.*

Proof. Fix $s_0 > 0$. suppose $\psi(z_j) \rightarrow z$. Then for $r \in (0, 1) \exists, n_0 \in \mathbb{N}$ such that $(\psi(z_j)) \in B_z(r, s)$ for all $j \geq n_0$,

$$B_z(r, s) = \left\{ j \in \mathbb{N} : \mathbf{T}(\psi(z_j) - z, s) \text{ or } \mathbf{I}(\psi(z_j) - z, s) < r, \mathbf{F}(\psi(z_j) - z, s) < r \right\}$$

Such that, $B_z^c(r, s)$ Then,

$$1 - \mathbf{T}((\psi(z_j) - z), s) < r \text{ and } \mathbf{I}((\psi(z_j) - z), s) < r, \mathbf{F}((\psi(z_j) - z), s) < r.$$

Hence,

$$\mathbf{T}((\psi(z_j) - z), s) \rightarrow 1 \text{ and } \mathbf{I}((\psi(z_j) - z), s) \rightarrow 0, \mathbf{F}((\psi(z_j) - z), s) \rightarrow 0 \text{ as } j \rightarrow \infty.$$

\Leftarrow

if for every $s > 0$, $\mathbf{T}((\psi(z_j) - z), s) \rightarrow 1$ and $\mathbf{I}((\psi(z_j) - z), s) \rightarrow 0$, $\mathbf{F}((\psi(z_j) - z), s) \rightarrow 0$ as $j \rightarrow \infty$ then for $r \in (0, 1) \exists, n_0 \in \mathbb{N}$ such that $1 - \mathbf{T}((\psi(z_j) - z), s) < r$ and $\mathbf{I}((\psi(z_j) - z), s) < r$, $\mathbf{F}((\psi(z_j) - z), s) < r$ for all $j \geq n_0$.

It follows that $\mathbf{T}((\psi(z_j) - z), s) > 1 - r$ and $\mathbf{I}((\psi(z_j) - z), s) < r$, $\mathbf{F}((\psi(z_j) - z), s) < r$ for all $j \geq n_0$. Thus $(\psi(z_j)) \in B_z^c(r, s)$ for all $j \geq n_0$ and hence $\rightarrow \psi(z_j) \square$

Theorem 3.10. A sequence $z = (z_k) \in S_{(\mathbf{T}, \mathbf{I}, \mathbf{F})}^N(\psi)$ is \mathcal{M} -convergent iff for each $\epsilon > 0$ and $s > 0 \exists$, a number $\mathcal{K} = \mathcal{K}(z, \epsilon, t)$ such that

$$\left\{ j \in \mathbb{N} : \mathbf{T}\left(\left(\psi(z_j) - \ell\right), \frac{s}{2}\right) > 1 - \epsilon \text{ or } \mathbf{I}\left(\left(\psi(z_j) - \ell\right), \frac{s}{2}\right) < \epsilon \text{ or } \mathbf{F}\left(\left(\psi(z_j) - \ell\right), \frac{s}{2}\right) < \epsilon \right\}$$

Proof. Suppose $\mathcal{M} - \lim x = \ell$. For a given $\epsilon > 0$, Let $s > 0$ such that $(1 - \epsilon) \diamond (1 - \epsilon) > 1 - s$ and $\epsilon \star \epsilon < s, \epsilon \star \epsilon < s$. Then

for every $z \in S_{(\mathbf{T}, \mathbf{I}, \mathbf{F})}^N(\psi)$

$$\mathcal{W} = \left\{ j \in \mathbb{N} : \mathbf{T}\left(\left(\psi(z_j) - \ell\right), \frac{s}{2}\right) \leq 1 - \epsilon \text{ or } \mathbf{I}\left(\left(\psi(z_j) - \ell\right), \frac{s}{2}\right) \geq \epsilon, \mathbf{F}\left(\left(\psi(z_j) - \ell\right), \frac{s}{2}\right) \geq \epsilon \right\}$$

which implies that

$$\mathcal{W}^c = \left\{ j \in \mathbb{N} : \mathbf{T}\left(\left(\psi(z_j) - \ell\right), \frac{s}{2}\right) > 1 - \epsilon \text{ or } \mathbf{I}\left(\left(\psi(z_j) - \ell\right), \frac{s}{2}\right) < \epsilon, \mathbf{F}\left(\left(\psi(z_j) - \ell\right), \frac{s}{2}\right) < \epsilon \right\}$$

Conversely, take $N \in \mathcal{W}$. Then

$$\mathbf{T}\left(\left(\psi(z_j) - \ell\right), \frac{s}{2}\right) > 1 - \epsilon \text{ or } \mathbf{I}\left(\left(\psi(z_j) - \ell\right), \frac{s}{2}\right) < \epsilon, \mathbf{F}\left(\left(\psi(z_j) - \ell\right), \frac{s}{2}\right) < \epsilon.$$

Now we have to prove that \exists , a number $\mathcal{K} = \mathcal{K}(z, \epsilon, t)$ such that,

$$\left\{ j \in \mathbb{N} : \mathbf{T}(\psi(z_j) - \psi(z_N), s) \leq 1 - \gamma \text{ or } \mathbf{I}(\psi(z_k) - \psi(z_N), s) \geq \gamma, \mathbf{F}(\psi(z_j) - \psi(z_N), s) \geq \gamma \right\}$$

for each $z \in S_{(\mathbf{T}, \mathbf{I}, \mathbf{F})}^N(\psi)$

$$B = \left\{ j \in \mathbb{N} : \mathbf{T}(\psi(z_j) - \psi(z_N), s) \leq 1 - \gamma \text{ or } \mathbf{I}(\psi(z_j) - \psi(z_N), s) \geq \gamma, \mathbf{F}(\psi(z_j) - \psi(z_N), s) \geq \gamma \right\}.$$

Now we will prove that $B \subset \mathcal{W}$. consider that $B \subseteq \mathcal{W}$ Then $\exists, n \in B$ and $n \notin \mathcal{W}$ so we get

$$\mathbf{T}(\psi(z_n) - \psi(z_N), s) \leq 1 - s \text{ or } \mathbf{I}(\psi(z_n) - \psi(z_N), \frac{s}{2}) > 1 - \epsilon, \mathbf{F}(\psi(z_n) - \psi(z_N), \frac{s}{2}) > 1 - \epsilon$$

In a different way $\mathbf{T}((\psi(z_N) - \ell), \frac{s}{2}) > 1 - \epsilon$ Therefore we have

$$1 - \gamma \geq \mathbf{T}(\psi(z_j) - \psi(z_N), s) \geq \mathbf{T}((\psi(z_n) - \ell), \frac{s}{2}) \diamond \mathbf{T}((\psi(z_N) - \ell), \frac{s}{2}) \geq (1 - \epsilon) \diamond (1 - \epsilon) > 1 - \gamma,$$

which is impossible. At the same time,

$$\mathbf{I}(\psi(z_n) - \psi(z_N), s) \geq \gamma \text{ or } \mathbf{I}((\psi(z_n) - \ell), \frac{s}{2}) < \epsilon,$$

In a similar way,

$$\mathbf{I}((\psi(z_N) - \ell), \frac{s}{2}) < \epsilon,$$

$$\gamma \leq \mathbf{I}(\psi(z_n) - \psi(z_N), s) \leq \mathbf{I}((\psi(z_n) - \ell), \frac{s}{2}) \star \mathbf{I}((\psi(z_N) - \ell), \frac{s}{2}) \leq \epsilon \star \epsilon < \gamma,$$

which is impossible. At the same time,

$$\mathbf{F}(\psi(z_n) - \psi(z_N), s) \geq \gamma \text{ or } \mathbf{F}((\psi(z_n) - \ell), \frac{s}{2}) < \epsilon,$$

In particular,

$$\mathbf{F}(\psi(z_N) - \ell), \frac{s}{2}) < \epsilon$$

$$\gamma \leq \mathbf{F}(\psi(z_n) - \psi(z_N), s) \leq \mathbf{F}(\psi(z_n) - \ell), \frac{s}{2}) \star \mathbf{F}(\psi(z_N) - \ell), \frac{s}{2}) \leq \epsilon \star \epsilon < \gamma,$$

which is impossible. Therefore $B \subset \mathcal{W}$. Hence $\mathcal{W} \in \mathcal{M}$ implies $B \in \mathcal{M}$. \square

Definition 3.11. Let $\mathcal{V} = (X, M, \diamond, \star)$ is a NNS. For $0 < s < 1$ we fix a closed ball $\mathcal{B}[z, r, s]$ with centre $z \in X$ and radius $r > 0$, there exist $k \in \mathbb{N}$ such that for all $j \geq k$

$$\mathcal{B}[z, r, s] = \{(a_k) \in \ell_\infty : \mathbf{T}(\psi(z_j) - \psi(a_k), s) \geq 1 - r \text{ or } \mathbf{I}(\psi(z_j) - \psi(a_k), s) \leq r, \mathbf{F}(\psi(z_j) - \psi(a_k), s) \leq r\}.$$

Lemma 3.12. Every closed ball $\mathcal{B}[z, r, s](\psi)$ is a closed set (CS).

Proof. Consider $a = (a_j) \in \ell_\infty$ such that $a \in \overline{B[z, r, s](\psi)}$. Since X is first countable, \exists , $a = (a_j) \in \overline{B[z, r, s](\psi)}$ such that $a_j \rightarrow a$ as $j \rightarrow \infty$. implies the set

$$X = \left\{ (a_j) \in \ell_\infty : \mathbf{T}(\psi(z_j) - \psi(a_j), s) \geq 1 - r \text{ or } \mathbf{I}(\psi(z_j) - \psi(a_j), s) \leq r, \mathbf{F}(\psi(z_j) - \psi(a_j), s) \leq r \right\}.$$

Since $a_j \rightarrow a$, $\mathbf{T}(\psi(a_j) - \psi(a), s) \rightarrow 1$ and $\mathbf{I}(\psi(a_j) - \psi(a), s) \rightarrow 0$, $\mathbf{F}(\psi(a_j) - \psi(a), s) \rightarrow 0$, as $j \rightarrow \infty$. for all s . For a given $\epsilon > 0$,

Hence for $j \in X$

$$\mathbf{T}(\psi(z) - \psi(a), s + \epsilon) \geq \lim_{j \rightarrow \infty} \mathbf{T}(\psi(z) - \psi(a_j), s) \diamond \mathbf{T}(\psi(a_j) - \psi(a), \epsilon) \geq 1 \diamond (1 - r) = 1 - r$$

and

$$\mathbf{I}(\psi(z) - \psi(a), s + \epsilon) \leq \lim_{j \rightarrow \infty} \mathbf{I}(\psi(z) - \psi(a_j), s) \star \mathbf{I}(\psi(a_j) - \psi(a), \epsilon) \leq 0 \star r = r.$$

$$\mathbf{F}(\psi(z) - \psi(a), s + \epsilon) \leq \lim_{j \rightarrow \infty} \mathbf{F}(\psi(z) - \psi(a_j), s) \star \mathbf{F}(\psi(a_j) - \psi(a), \epsilon) \leq 0 \star r = r.$$

In particular for $b \in \mathbb{N}$, take $\epsilon = \frac{1}{b}$. Then,

$$\mathbf{T}(\psi(z) - \psi(a), s) = \lim_{b \rightarrow \infty} \mathbf{T}\left(\psi(z) - \psi(a), s + \frac{1}{b}\right) \geq 1 - r.$$

and

$$\mathbf{I}(\psi(z) - \psi(a), s) = \lim_{b \rightarrow \infty} \mathbf{I}\left(\psi(z) - \psi(a), r + \frac{1}{b}\right) \leq r.$$

$$\mathbf{F}(\psi(z) - \psi(a), s) = \lim_{b \rightarrow \infty} \mathbf{F}\left(\psi(z) - \psi(a), s + \frac{1}{b}\right) \leq r.$$

\implies The set

$$\left\{ (a_j) \in \ell_\infty : \mathbf{T}(\psi(z_j) - \psi(a_j), s) \geq 1 - r \text{ or } \mathbf{I}(\psi(z_j) - \psi(a_j), s) \leq r, \mathbf{F}(\psi(z_j) - \psi(a_j), s) \leq r \right\}.$$

$\implies a \in B[z, r, s](\psi)$. Therefore $B[z, r, s](\psi)$ is a closed set. \square

4. Conclusions

In this review, we have studied the concept of statistical convergence using neutrosophic norm space with the help of compact operator, which has an important place in the literature. We have defined the compact operator form statistical convergence of sequence spaces $\mathcal{S}_{(\mathbf{T},\mathbf{I},\mathbf{F})}^N(\psi)$ and $S_{0(\mathbf{T},\mathbf{I},\mathbf{F})}^N(\psi)$ and investigated basic properties. These are illustrated by proper examples.

Acknowledgement

We wish to thank the referee for his/her valuable suggestions and comments which improved the work considerably.

References

1. Smarandache, F. (1999). A unifying field in logics, Neutrosophic Probability, Set and Logic, Rehoboth: American Research Press.
2. Kreyszig E. Introductory Functional Analysis with Application. New York, Chicheste, Brisbane, Toronto: John Wiley and Sons, Inc; 1978
3. M. Kirişci, N. Şimşek, Neutrosophic normed spaces and statistical convergence, The Journal of Analysis, (2020) 1-15. J Anal 28, 1059–1073 (2020). <https://doi.org/10.1007/s41478-020-00234-0>
4. L.A. Zadeh, Fuzzy sets, Inform. Control 8 (1965) 338-353.
5. M. Kirişci, N. Şimşek, Neutrosophic Metric Spaces, arXiv preprint arXiv:1907.00798.
6. J.H. Park, Intuitionistic fuzzy metric spaces, Chaos Solitons Fractals 22 (2004) 1039-1046.
7. R. Saadati, J.H. Park, On the intuitionistic fuzzy topological spaces, Chaos Solitons Fractals 27 (2006) 331-344.
8. B.C. Tripathy, A. Baruah, M. Et and M. Gungor : On almost statistical convergence of new type of generalized difference sequence of fuzzy numbers, Iranian Journal of Science and Technology, Transactions A : Science, 36(2)(2012), 147-155.
9. F. Smarandache, Neutrosophic Set is a Generalization of Intuitionistic Fuzzy Set, Inconsistent Intuitionistic Fuzzy Set (Picture Fuzzy Set, Ternary Fuzzy Set), Pythagorean Fuzzy Set, Spherical Fuzzy Set, and q-Rung Orthopair Fuzzy Set, while Neutrosophication is a Generalization of Regret Theory, Grey System Theory, and Three-Ways Decision (revisited), *Journal of New Theory*, **29** 2019, 1-31.
10. F. Smarandache, NeutroAlgebra is a Generalization of Partial Algebra, *International Journal of Neutrosophic Science (IJNS)*, Volume 2, **2020** .
11. Das, S., and Pramanik, S. (2020). Neutrosophic simply soft open set in neutrosophic soft topological space. *Neutrosophic Sets and Systems*, 38, 235-243.
12. F. Smarandache, NeutroAlgebraic Structures and AntiAlgebraic Structures, in *Advances of Standard and Nonstandard Neutrosophic Theories* , *Pons Publishing House Brussels, Belgium.*, **Chapter No. 6**, pages 240 - 265 , 2019.
13. B.C. Tripathy and R. Goswami: Fuzzy real valued p-absolutely summable multiple sequences in probabilistic normed spaces; *Afrika Matematika*, 26 (7-8), (2015) 1281-1289.
14. M. Mursaleen, S. A. Mohiuddine, Statistical convergence of double sequences in intuitionistic fuzzy normed spaces, *Chaos, Solitons & Fractals* 41(5)(2009) 2414–2421.
15. Das, S. and Pramanik, S. (2020). Generalized neutrosophic b-open sets in neutrosophic topological space, *Neutrosophic Sets and Systems*, 35, 522-530.
16. Steinhaus H (1951) Sur la convergence ordinaire et la convergence asymptotique. *Colloq. Math.* 2:

Vakeel A. Khan, Mohammad Daud Khan, Some Topological Character of Neutrosophic normed spaces

17. B.C. Tripathy and A.J. Dutta, Statistically pre-Cauchy fuzzy real-valued sequences defined by Orlicz function, *Proyecciones Jour Math.*, 33(3), (2014), 235-243.
18. Fast H (1951) Sur la convergence statistique. *Colloq. Math.* 2: 241-244
19. M. Et, B. C. Tripathy and A.J. Dutta : On pointwise statistical convergence of order alpha of sequences of fuzzy mappings; *Kuwait Journal of Science*, 41(3) (2014), 17-30.
20. Karakus S (2007) Statistical convergence on probabilistic normed spaces. *Math. Commun.* 12: 11-23
21. Das, R., and Tripathy, B.C. (2020). Neutrosophic Multiset Topological Space. *Neutrosophic Sets and Systems*, 35, 142-152
22. V. A. Khan, H. Fatima, M. Daud. Khan, Space of neutrosophic lambda- statistical convergence sequence and their properties, Vol. 2021 (23), *jmcS* **2021**, 1-9. <http://dx.doi.org/10.22436/jmcs.023.01.01>
23. F. Smarandache, Neutrosophic set, a generalisation of the intuitionistic fuzzy sets, *Inter. J. Pure Appl. Math.*, 24, (2005), 287–297.
24. T Bera, NK Mahapatra, Neutrosophic Soft Linear Spaces, *Fuzzy Information and Engineering*, 9 (2017) 299-324.
25. T Bera, NK Mahapatra, Neutrosophic Soft Normed Linear Spaces, *neutrosophic Sets and Systems*, 23 (2018) 52-71.
26. Mukherjee, A., and Das, R. (2020). Neutrosophic Bipolar Vague Soft Set and Its Application to Decision Making Problems. *Neutrosophic Sets and Systems*, 32, 410-424.
27. V. A. Khan, M. Ahmad, H. Fatima, Hira and M. F. Khan, On some results in intuitionistic fuzzy ideal convergence double sequence spaces, *Advances in Difference Equations*, 1(2019) 1-10.
28. V. A. Khan, R. K. A. Rababah, H. Fatima, M. Ahmad, Intuitionistic fuzzy I-convergent sequence spaces defined by bounded linear operator, *ICICI Express Letters* 9(2018)955-962.
29. K.T. Atanassov, Intuitionistic fuzzy sets, *Fuzzy Sets and Systems* 20(1) (1986), 87–96.
30. Vakeel A. Khan, Ayhan Esi, Mobeen Ahmad and Mohammad Daud Khan, Continuous and Bounded Linear Operators in Neutrosophic Normed Spaces, *Journal of Intelligent and Fuzzy Systems* 40 (2021) 11063–11070, DOI:10.3233/JIFS-202189 IOS Press.
31. F. Smarandache, Neutrosophic set-a generalization of the intuitionistic fuzzy set, *International journal of pure and applied mathematics*, 24(3) **2005** pages 287.
32. J.A. Fridy, On statistical convergence, *Analysis* 5 (1985),301–313.
33. M. Ilkhan and E.E. Kara, A new type of statistical Cauchy sequence and its relation to Bourbaki completeness, *Cogent Mathematics & Statistics* 5(1) (2018), 1–9. <https://doi.org/10.1080/25742558.2018.1487500>.
34. Florentin Smarandache, A Unifying Field in Logics: Neutrosophic Logic, Multiple-Valued Logic / An International Journal, Vol. 8, No. 3, 385-438, 2002, <http://fs.unm.edu/eBook-Neutrosophics6.pdf>
35. Florentin Smarandache, Neutrosophy. Neutrosophic Probability, Set, and Logic. ProQuest Information and Learning, Ann Arbor, Michigan, USA, 105 p., 1998, 2000, 2002, 2005, 2006.
36. E.E. Kara, Some topological and geometrical properties of new Banach sequence spaces, *J Ineq Appl* 2013(38), (2013), 15. doi: 10.1186/1029-242X-2013-38.
37. V. A. Khan, M. Daud. Khan, M. Ahmad, Some New Type of Lacunary Statistically Convergent Sequences In Neutrosophic Normed Space, Vol. 42, 2021, *Neutrosophic Sets and Systems*, Vol. 42, **2021**, 1-14.
38. Das, R., Smarandache, F., and Tripathy, B.C. (2020). Neutrosophic Fuzzy Matrices and Some Algebraic Operations. *Neutrosophic Sets and Systems*, 32, 401-409.
39. J. Connor, The Statistical and Strong p-Cesaro Convergence of Sequences, *Analysis* 8 (1988) 47–63
40. Schweizer B, Sklar A. and others. Statistical metric spaces. *Pacific J. Math*; 10(1), 313-334, 1960.

Received: Aug 1, 2021. Accepted: Dec 3, 2021