



# VIKOR based MAGDM Strategy with Trapezoidal Neutrosophic Numbers

Surapati Pramanik<sup>1</sup>, Rama Mallick<sup>2</sup>

<sup>1</sup> Department of Mathematics, Nandalal Ghosh B.T. College, Panpur, P.O.-Narayanpur, District –North 24 Parganas, Pin code-743126, West Bengal, India. E-mail: sura\_pati@yahoo.co.in

<sup>2</sup> Department of Mathematics, Umeschandra College, Surya Sen Street ,Kolkata-700012, West Bengal, India .  
E-mail: ramamallick23@gmail.com

**Abstract.** VIKOR strategy is a popular strategy for multi-attribute decision making (MADM). We extend the VIKOR strategy for MAGDM problems in trapezoidal neutrosophic number environment. In decision making situation, single-valued trapezoidal neutrosophic numbers are employed to express the attribute values. Then we develop an extended VIKOR strategy to deal with MAGDM in single-valued trapezoidal neutrosophic number environment. The influence of decision-making mechanism coefficient is presented. To illustrate and validate the proposed VIKOR strategy, an illustrative numerical example of MAGDM problem is solved in trapezoidal neutrosophic number environment.

**Keywords:** Neutrosophic set, Trapezoidal neutrosophic fuzzy number, Multi-attribute decision making, VIKOR strategy..

## 1. Introduction:

Smarandache [1] pioneered the neutrosophic set based on neutrosophy in 1998. In 2010, Wang et al. [2] proposed single valued neutrosophic set (SVNS). SVNS has been successfully applied to solve decision making problems [3-31], image processing [32-35], conflict resolution [36], educational problem [37, 38], social problem [39, 40], etc. Broumi et al. [41] presented an overview neutrosophic sets. Recently, Peng and Dai [42] presented a bibliometric analysis of neutrosophic sets for last two decades.

Single valued trapezoidal neutrosophic number (SVTrNN) [43] is an extension of SVNS. Every element of SVTrNN is a trapezoidal number with three membership degrees namely, truth, indeterminacy and falsity membership degrees. Deli and Subhas [10] developed a ranking strategy of SVTrNNs. Biswas et al. [7] established value and ambiguity based ranking strategy for SVTrNN and employed the strategy to deal with MADM problem. Biswas et al. [44] developed TOPSIS strategy for MADM with trapezoidal neutrosophic numbers (TrNNs). Biswas et al. [45] presented distance measure based MADM strategy with interval trapezoidal neutrosophic numbers (ITrNNs). For simplicity, we call SVTrNN as TrNN.

In 1998, Opricovic [46] first introduced the VIKOR strategy in the literature to deal with multi criteria group decision making (MCGDM) with conflicting objectives[47,48].

Using TOPSIS and VIKOR, Poursmaeil et al.[49] proposed an MAGDM strategy in SVNS environment. Bausys and Zavadskas[50] proposed the VIKOR strategy in interval neutrosophic set (INS) environment. Huang et al.[51] developed a VIKOR strategy in INS environment. Liu and Zhang[52] studied VIKOR strategy in neutrosophic hesitant fuzzy set environment. In 2017, Hu et al.[53] developed a projection based VIKOR strategy for doctor selection problem in INS environment. In 2017, Selvakumari et al. [54] studied VIKOR strategy using octagonal neutrosophic soft matrix. Pramanik et al. [55] developed a VIKOR based MAGDM strategy in bipolar neutrosophic set environment. In 2018, Pramanik et al. [56] introduced a VIKOR strategy for MAGDM in neutrosophic cubic set environment. Dalapati and Pramanik [57] further revisited VIKOR based MAGDM strategy [56] to make it more comprehensive.

VIKOR strategy in trapezoidal neutrosophic number (TrNN) environment is not studied in the literature.

To fill up this research gap, we propose a VIKOR strategy to deal with MAGDM problems in TrNN environment. Also, we solve an MAGDM problem based on VIKOR strategy in trapezoidal neutrosophic number.

The rest of the paper is developed as follows. In section 2, we briefly describe definitions of trapezoidal fuzzy number, TrNN, trapezoidal neutrosophic weighted arithmetic averaging (TrNWAA) operator, Hamming distance between two TrNNs. In section 3, we briefly describe extended VIKOR strategy. Thereafter in section 4, we present a VIKOR strategy in TrNN environment. In section 5, we solve an MAGDM problem using the proposed

VIKOR strategy. In section 6, we present the sensitivity analysis. We represent conclusion and scope of future research in section 7.

**2. Preliminaries**

We present some fundamental definitions of fuzzy sets, neutrosophic set, SVNS, and TrNN.

**Definition 2.1** [58] Let  $\check{Y}$  be a universal set . Then, a fuzzy set F is presented as:

$$F = \{ \langle \dot{y}, \nu_F(\dot{y}) : \dot{y} \in \check{Y} \rangle \} \tag{1}$$

where  $\nu_F(\dot{y})$  is the degree of membership which maps  $\check{Y}$  to  $[0,1]$  or we can express by  $\nu_F : \check{Y} \rightarrow [0,1]$ .

**Definition 2.2**[1]Let  $\check{Y}$  be an universal set . A neutrosophic set N can be presented of the form:

$$N = \{ \langle z : T_N(\dot{y}), I_N(\dot{y}), F_N(\dot{y}) > \dot{y} \in \check{Y} \rangle \} \tag{2}$$

where the functions  $T, I, F : \check{Y} \rightarrow ]^{-}0, 1^{+}[$  define repectively the degree of truth membership, the degree of indeterminacy, and the degree of non-membership or falsity of the component  $\dot{y} \in \check{Y}$  and satisfy the condition,

$$^{-}0 \leq T_N(\dot{y}) + F_N(\dot{y}) + I_N(\dot{y}) \leq 3^{+} \tag{3}$$

**Definition 2.3** [2] Let  $\check{Y}$  be a universal set. An SVNS N in  $\check{Y}$  is described by

$$N = \{ \langle \dot{y} : T_N(\dot{y}), I_N(\dot{y}), F_N(\dot{y}) > \dot{y} \in \check{Y} \rangle \}$$

where  $T_N(\dot{y}) : \check{Y} \rightarrow [0,1]$  ,  $I_N(\dot{y}) : \check{Y} \rightarrow [0,1]$  and  $F_N(\dot{y}) : \check{Y} \rightarrow [0,1]$  with the condition  $0 \leq T_N(\dot{y}) + F_N(\dot{y}) + I_N(\dot{y}) \leq 3$  for all  $\dot{y} \in \check{Y}$  .The functions  $T_N(\dot{y}), I_N(\dot{y})$  and  $F_N(\dot{y})$  are respectively, the truth membership function, the indeterminacy membership function and the falsity membership function of the element  $\dot{y}$  to the set N.

**Definition 2.4**[59] A generalized trapezoidal fuzzy number  $\dot{T}$  denoted by  $\dot{T} = (\dot{b}_1, \dot{b}_2, \dot{b}_3, \dot{b}_4; \nu)$  is described as a fuzzy subset of the real number R with membership function  $\kappa_{\dot{T}}$  which is defined by

$$\kappa_{\dot{T}}(x) = \begin{cases} \frac{(x - \dot{b}_1)\nu}{(\dot{b}_2 - \dot{b}_1)}, \dot{b}_1 \leq x < \dot{b}_2 \\ \nu, \dot{b}_2 \leq x \leq \dot{b}_3 \\ \frac{(\dot{b}_4 - x)\nu}{(\dot{b}_4 - \dot{b}_3)}, \dot{b}_3 < x \leq \dot{b}_4 \\ 0, \text{otherwise} \end{cases} \tag{4}$$

where  $\dot{b}_1, \dot{b}_2, \dot{b}_3, \dot{b}_4$  are real number satisfying  $\dot{b}_1 \leq \dot{b}_2 \leq \dot{b}_3 \leq \dot{b}_4$  and  $\nu$  is the membership degree.

**Definition 2.5**[43, 44] Let  $x$  be a TrNN. Then, its truth membership, indeterminacy membership, and falsity membership functions are presented respectively as:

$$T_x(z) = \begin{cases} \frac{(z - b'_1)t_x}{(b'_2 - b'_1)}, b'_1 \leq z < b'_2 \\ t_x, b'_2 \leq z \leq b'_3 \\ \frac{(b'_4 - z)t_x}{(b'_4 - b'_3)}, b'_3 \leq z \leq b'_4 \\ 0, \text{otherwise} \end{cases} \tag{5}$$

$$I_x(z) = \begin{cases} \frac{(b'_2 - z) + (z - b'_1)i_x}{(b - a)}, b'_1 \leq z < b'_2 \\ i_x, b'_2 \leq z \leq b'_3 \\ \frac{z - b'_3 + (b'_4 - z)i_x}{b'_4 - b'_3}, b'_3 < z \leq b'_4 \\ 0, \text{otherwise} \end{cases} \tag{6}$$

$$F_x(z) = \begin{cases} \frac{b'_2 - z + (z - b'_1)f_x}{b'_2 - b'_1}, & b'_1 \leq z < b'_2 \\ f_x, & b'_2 \leq z \leq b'_3 \\ \frac{z - b'_3 + (b'_4 - z)f_x}{b'_4 - b'_3}, & b'_3 < z \leq b'_4 \\ 0, & \text{otherwise} \end{cases} \tag{7}$$

Here  $0 \leq T_x(z) \leq 1, 0 \leq I_x(z) \leq 1$  and  $0 \leq F_x(z) \leq 1$  and  $0 \leq T_x(z) + I_x(z) + F_x(z) \leq 3; b'_1, b'_2, b'_3, b'_4 \in R$ . Then  $x = ([b'_1, b'_2, b'_3, b'_4]; t_x, i_x, f_x)$  is called a TrNN.

**Definition 2.6** [43] Let  $m_i = \langle (b_{1i}, b_{2i}, b_{3i}, b_{4i}); T_{m_i}, I_{m_i}, F_{m_i} \rangle (i = 1, 2, \dots, n')$  be a group of TrNNs, then a trapezoidal neutrosophic weighted arithmetic averaging (TrNWAA) operator is defined as follows:

$$\text{TrNWAA}(m_1, m_2, \dots, m_n) = \sum_{i=1}^{n'} \dot{w}_i m_i \tag{8}$$

where,  $\dot{w}_i$  is the weight of  $m_i (i = 1, 2, \dots, n)$  such that  $\dot{w}_i > 0$  and  $\sum_{i=1}^{n'} \dot{w}_i = 1$ . Specially, when  $\dot{w}_i = 1/n'$  for  $i=1, 2, \dots, n'$  the TrNWAA operator transform into the trapezoidal neutrosophic arithmetic averaging (TrNAA) operator.

**Definition 2.7**[44] Let  $m_1 = ([\dot{p}_1, \dot{q}_1, \dot{r}_1, \dot{s}_1]; t_{m_1}, i_{m_1}, f_{m_1})$  and  $m_2 = ([\dot{p}_2, \dot{q}_2, \dot{r}_2, \dot{s}_2]; t_{m_2}, i_{m_2}, f_{m_2})$  be any two TrNNs. The normalized Hamming distance between  $m_1$  and  $m_2$  is defined as:

$$d(m_1, m_2) = \frac{1}{12} \left( \left| \dot{p}_1(2+t_{m_1}-i_{m_1}-f_{m_1}) - \dot{p}_2(2+t_{m_2}-i_{m_2}-f_{m_2}) \right| + \left| \dot{q}_1(2+t_{m_1}-i_{m_1}-f_{m_1}) - \dot{q}_2(2+t_{m_2}-i_{m_2}-f_{m_2}) \right| \right. \\ \left. + \left| \dot{r}_1(2+t_{m_1}-i_{m_1}-f_{m_1}) - \dot{r}_2(2+t_{m_2}-i_{m_2}-f_{m_2}) \right| + \left| \dot{s}_1(2+t_{m_1}-i_{m_1}-f_{m_1}) - \dot{s}_2(2+t_{m_2}-i_{m_2}-f_{m_2}) \right| \right) \tag{9}$$

**2.8. Standardize the decision matrix**[44]

Let  $D = (b_{ij})_{p \times n}$  be a neutrosophic matrix, where  $b_{ij} = ([b_{ij}^1, b_{ij}^2, b_{ij}^3, b_{ij}^4]; t_{b_{ij}}, i_{b_{ij}}, f_{b_{ij}})$  is the rating value of the alternative  $X_i$  with respect to attribute  $Y_j$ . To remove the effect of several physical dimensions, we standardize the decision matrix  $(b_{ij})_{p \times n}$  for benefit type and cost type attributes.

We denote the standardized decision matrix by  $D^* = (s_{ij})_{p' \times n'}$

- 1. For benefit type attribute

$$b_{ij}^* = ([\frac{b_{ij}^1}{v_j^+}, \frac{b_{ij}^2}{v_j^+}, \frac{b_{ij}^3}{v_j^+}, \frac{b_{ij}^4}{v_j^+}]; t_{b_{ij}}, i_{b_{ij}}, f_{b_{ij}}) \tag{10}$$

- 2. For cost type attribute:

$$b_{ij}^* = ([\frac{v_j^-}{b_{ij}^4}, \frac{v_j^-}{b_{ij}^3}, \frac{v_j^-}{b_{ij}^2}, \frac{v_j^-}{b_{ij}^1}]; t_{b_{ij}}, i_{b_{ij}}, f_{b_{ij}}) \tag{11}$$

Here  $v_j^+ = \max\{b_{ij}^4 : i = 1, 2, \dots, p'\}$  and  $v_j^- = \min\{b_{ij}^1 : i = 1, 2, \dots, p'\}$  for  $j = 1, 2, \dots, n'$

Hence, we obtain standardized matrix  $D^*$  as:

$$D^* = (s_{ij})_{p' \times n'} = \begin{pmatrix} & b_1 & b_2 & \dots & b_{n'} \\ a_1 & s_{11} & s_{12} & \dots & s_{1n'} \\ a_2 & s_{21} & s_{22} & \dots & s_{2n'} \\ \dots & \dots & \dots & \dots & \dots \\ a_{p'} & s_{p'1} & s_{p'2} & \dots & s_{p'n'} \end{pmatrix} \tag{12}$$

**3. VIKOR Strategy for MADM**

Assume that  $B_1, B_2, \dots, B_s$  are the s alternatives. For the alternative  $B_r$ , assume that the rating of the  $j'$  th criterion is  $h_{rj'}$ , i.e.  $h_{rj'}$  is the value of  $j'$  th criterion for the alternative  $B_i$ ; the number of criteria is assumed to be r. Development of the extended VIKOR strategy is started with the following form of  $L_p$  - metric:

$$L_{iq} = \left\{ \sum_{i=1}^r [(h_i^+ - h_{ij'}) / (h_i^+ - h_i^-)]^q \right\}^{\frac{1}{q}}, 1 \leq q \leq \infty; i' = 1, 2, \dots, s \tag{13}$$

To formulate ranking measure,  $L_{1,i'}$  (as  $S_{i'}$ ) and  $L_{\infty,i'}$  (as  $R_{i'}$ ) are employed. The solution obtained by  $\min S_{i'}$  reflects a maximum group utility (“majority” rule), and the solution obtained by  $\min R_{i'}$  reflects a minimum individual regret of the “opponent”.

VIKOR strategy is presented using the following steps:

- (a) Evaluate the best  $h_j^+$  and the worst  $h_j^-$  values of all criteria  $j' = 1, 2, \dots, n'$ .

$$h_j^+ = \max_{i'} h_{ij'}, h_j^- = \min_{i'} h_{ij'}, \text{for benefit criterion,} \tag{14}$$

$$h_j^+ = \min_{i'} h_{ij'}, h_j^- = \max_{i'} h_{ij'}, \text{for cost criterion} \tag{15}$$

- (a) Calculate the values  $F_{i'}$  and  $G_{i'}$ ;  $i' = 1, 2, \dots, m'$ , by these relations:

$$F_{i'} = \sum_{j'=1}^{n'} \frac{w_{j'}(h_{ij'}^+ - h_{ij'})}{(h_{ij'}^+ - h_{ij}^-)} \tag{16}$$

$$G_{i'} = \max_{j'} \left( \frac{w_{j'}(h_{ij'}^+ - h_{ij'})}{(h_{ij'}^+ - h_{ij}^-)} \right) \tag{17}$$

where  $w_{j'}$  ( $j' = 1, 2, \dots, r$ ) represent the weights of criteria.

- (b) Evaluate the value  $K_{i'}$ ;  $i' = 1, 2, \dots, s$ , using following relation:

$$K_{i'} = \frac{\nu(F_{i'} - F^+)}{(F - F^+)} + \frac{(1-\nu)(G_{i'} - G^+)}{(G^- - G^+)} \tag{18}$$

where

$$F^+ = \min_{i'} F_{i'}, F^- = \max_{i'} F_{i'}$$

$$G^+ = \min_{i'} G_{i'}, G^- = \max_{i'} G_{i'}$$

Here,  $\nu$  indicates the weight of the technique of “the majority of criteria” (or “the maximum group utility”). Here we set  $\nu = 0.5$ .

- (c) Sorting by the values  $F$ ,  $G$  and  $K$  in decreasing order, we rank the alternatives.

- (d) Propose the alternative  $B^1$  as a compromise solution that is ranked the best by the measure  $K$  (minimal) subject to the conditions  $A1$  and  $A2$ :

$A1$ . Acceptable advantage:

$$K(B^2) - K(B^1) \geq DK$$

where  $B^2$  is second alternative in the ranking list by  $K$ ;  $DK = 1/(s-1)$ ;  $s =$  the number of alternatives.

$A2$ . Acceptable stability in decision making:

Using  $F$  or/and  $G$ , we must have alternative  $B^1$  as the best ranked. We say the compromise solution as stable subject to

- i. “voting by majority rule” (when  $\nu > 0.5$  is needed),
- ii. or “by consensus”  $\nu \approx 0.5$ ,
- iii. or “with veto” ( $\nu < 0.5$ ).

$\nu$  reflects the weight of the decision making strategy of “the majority of criteria” (or “the maximum group utility”).

- $B^1$  and  $B^2$  are compromise solutions if  $A2$  is not satisfied, Or
- $B^1, B^2, \dots, B^M$  are compromise solutions if  $A1$  is not satisfied.

Evaluate  $K(B^2) - K(B^1) \geq DK$  to determine  $B^M$  for maximum  $M$  (the positions of these alternatives are “in closeness”).

The minimal value of  $K$  determines the best alternative.

**4.VIKOR strategy for solving MAGDM problem in TrNN environment:**

Consider an MAGDM problem consisting of r alternatives and t attributes. The alternatives and attributes are presented by  $\alpha' = \{\alpha'_1, \alpha'_2, \dots, \alpha'_r\}$  and  $\beta' = \{\beta'_1, \beta'_2, \dots, \beta'_t\}$  respectively. Assume that  $\tau = \{\tau_1, \tau_2, \dots, \tau_t\}$  is the set of weights of the attributes, where  $\tau_{i'} \geq 0$  and  $\sum_{i'=1}^t \tau_{i'} = 1$ . Assume that  $B = \{B_1, B_2, \dots, B_K\}$  be the set of K decision makers and  $\sigma = \{\sigma_1, \sigma_2, \dots, \sigma_K\}$  be the set of weights of the decision makers, where  $\sigma_K \geq 0$  and  $\sum_{i'=1}^K \sigma_{i'} = 1$ . The

rating values offered by the experts are presented in terms of Trnn

The MAGDM strategy is described as follows:

Step-1: Let  $D = (p_{ij}^{N'})$  ( $N' = 1, 2, \dots, s$ ) be the  $N'$ -th decision matrix where  $\alpha'_{i'}$  is alternative with respect to attribute  $\beta'_{j'}$ . The  $N'$ -th decision matrix denoted by  $D^{N'}$  is presented as:

$$D^M = \begin{pmatrix} & \beta'_1 & \beta'_2 & \dots & \beta'_t \\ \alpha'_1 & p_{11}^{N'} & p_{12}^{N'} & \dots & p_{1t}^{N'} \\ \alpha'_2 & p_{21}^{N'} & p_{22}^{N'} & \dots & p_{2t}^{N'} \\ \dots & \dots & \dots & \dots & \dots \\ \alpha'_r & p_{r1}^{N'} & p_{r2}^{N'} & \dots & p_{rt}^{N'} \end{pmatrix} \tag{19}$$

where  $N' = 1, 2, \dots, s$ ;  $i' = 1, 2, \dots, r$ ;  $j' = 1, 2, \dots, t$ .

Step-2: To standardize the benefit criterion, we use the equation (10) and for cost criterion, we use (11). After standardizing, the decision matrix reduces to

$$D^M = \begin{pmatrix} & \beta'_1 & \beta'_2 & \dots & \beta'_t \\ \alpha'_1 & p_{11}^{*N'} & p_{12}^{*N'} & \dots & p_{1t}^{*N'} \\ \alpha'_2 & p_{21}^{*N'} & p_{22}^{*N'} & \dots & p_{2t}^{*N'} \\ \dots & \dots & \dots & \dots & \dots \\ \alpha'_r & p_{r1}^{*N'} & p_{r2}^{*N'} & \dots & p_{rt}^{*N'} \end{pmatrix}$$

$N'=1, 2, \dots, s$ ;  $i'=1, 2, \dots, r$ ;  $j' = 1, 2, \dots, t$ .

Step-3: To obtain aggregate decision matrix, we use trapezoidal neutrosophic weighted arithmetic operator(TrNWAA) which is presented below:

$$p_{ij} = \text{TrNWAA}(p_{ij}^1, p_{ij}^2, \dots, p_{ij}^M) \\ = \sum_{q=1}^M \sigma_q p_{ij}^q \tag{20}$$

Therefore, we obtain the aggregated decision matrix as:

$$D^{N'} = \begin{pmatrix} & \beta'_1 & \beta'_2 & \dots & \beta'_t \\ \alpha'_1 & \check{p}_{11} & \check{p}_{12} & \dots & \check{p}_{1t} \\ \alpha'_2 & \check{p}_{21} & \check{p}_{22} & \dots & \check{p}_{2t} \\ \dots & \dots & \dots & \dots & \dots \\ \alpha'_r & \check{p}_{r1} & \check{p}_{r2} & \dots & \check{p}_{rt} \end{pmatrix}$$

Step -4: Define the positive ideal solution (PIS)  $S^+$  and negative ideal solution (NIS)  $S^-$

$$S^+ = ([\check{b}_1^+, \check{b}_2^+, \check{b}_3^+, \check{b}_4^+]; t_b^+, i_b^-, f_b^-) = ([\check{b}_1^+, \check{b}_2^+, \check{b}_3^+, \check{b}_4^+]; \max t_b, \min i_b, \min f_b) \tag{21}$$

$$S^- = ([\check{b}_1^-, \check{b}_2^-, \check{b}_3^-, \check{b}_4^-]; t_b^-, i_b^+, f_b^+) = ([\check{b}_1^-, \check{b}_2^-, \check{b}_3^-, \check{b}_4^-]; \max t_b, \min i_b, \min f_b) \tag{22}$$

Step 5: Compute

$$\Gamma_m = \sum_{j=1}^t \tau_m \times \frac{d(([\check{b}_1^+, \check{b}_2^+, \check{b}_3^+, \check{b}_4^+]; t_b^+, i_b^-, f_b^-), ([\check{b}_{j1}, \check{b}_{j2}, \check{b}_{j3}, \check{b}_{j4}]; t_{nj}, i_{nj}, f_{nj}))}{d(([\check{b}_1^+, \check{b}_2^+, \check{b}_3^+, \check{b}_4^+]; t_b^+, i_b^-, f_b^-), ([\check{b}_1^-, \check{b}_2^-, \check{b}_3^-, \check{b}_4^-]; t_b^-, i_b^+, f_b^+))} \tag{23}$$

$$Z_m = \max \sum_{j=1}^t \frac{\tau_m \times d(([\dot{b}_1^+, \dot{b}_2^+, \dot{b}_3^+, \dot{b}_4^+]; t_b^+, i_b^-, f_b^-), ([\dot{b}_{j1}, \dot{b}_{j2}, \dot{b}_{j3}, \dot{b}_{j4}]; t_{nj}, i_{nj}, f_{nj}))}{d(([\dot{b}_1^+, \dot{b}_2^+, \dot{b}_3^+, \dot{b}_4^+]; t_b^+, i_b^-, f_b^-), ([\dot{b}_1^-, \dot{b}_2^-, \dot{b}_3^-, \dot{b}_4^-]; t_b^-, i_b^+, f_b^+))} \tag{24}$$

where  $\tau_m$  is the weight of  $\beta'_m$ .

Using equation (9), we obtain

$$d(([\dot{b}_1^+, \dot{b}_2^+, \dot{b}_3^+, \dot{b}_4^+]; t_b^+, i_b^-, f_b^-), ([\dot{b}_{j1}, \dot{b}_{j2}, \dot{b}_{j3}, \dot{b}_{j4}]; t_{nj}, i_{nj}, f_{nj})) = \frac{1}{12} \left( \left| \dot{b}_1^+ (2+t_b^+ - i_b^- - f_b^-) - \dot{b}_{j1} (2+t_{nj} - i_{nj} - f_{nj}) \right| + \left| \dot{b}_2^+ (2+t_b^+ - i_b^- - f_b^-) - \dot{b}_{j2} (2+t_{nj} - i_{nj} - f_{nj}) \right| \right. \\ \left. + \left| \dot{b}_3^+ (2+t_b^+ - i_b^- - f_b^-) - \dot{b}_{j3} (2+t_{nj} - i_{nj} - f_{nj}) \right| + \left| \dot{b}_4^+ (2+t_b^+ - i_b^- - f_b^-) - \dot{b}_{j4} (2+t_{nj} - i_{nj} - f_{nj}) \right| \right)$$

and

$$d(([\dot{b}_1^+, \dot{b}_2^+, \dot{b}_3^+, \dot{b}_4^+]; t_b^+, i_b^-, f_b^-), ([\dot{b}_1^-, \dot{b}_2^-, \dot{b}_3^-, \dot{b}_4^-]; t_b^-, i_b^+, f_b^+)) = \frac{1}{12} \left( \left| \dot{b}_1^+ (2+t_b^+ - i_b^- - f_b^-) - \dot{b}_1^- (2+t_b^- - i_b^+ - f_b^+) \right| + \left| \dot{b}_2^+ (2+t_b^+ - i_b^- - f_b^-) - \dot{b}_2^- (2+t_b^- - i_b^+ - f_b^+) \right| \right. \\ \left. + \left| \dot{b}_3^+ (2+t_b^+ - i_b^- - f_b^-) - \dot{b}_3^- (2+t_b^- - i_b^+ - f_b^+) \right| + \left| \dot{b}_4^+ (2+t_b^+ - i_b^- - f_b^-) - \dot{b}_4^- (2+t_b^- - i_b^+ - f_b^+) \right| \right)$$

Step 6: Compute the  $\Theta$  by the following formula:

$$\Theta_m = \Psi \frac{(\Gamma_m - \Gamma_m'')}{(\Gamma_m^+ - \Gamma_m'')} + (1 - \Psi) \frac{(Z_m - Z_m'')}{(Z_m^+ - Z_m'')} \tag{25}$$

$$\text{where } \Gamma_m'' = \min_m \Gamma_m, \Gamma_m^+ = \max_m \Gamma_m \tag{26}$$

$$Z_m'' = \min_m Z_m, Z_m^+ = \max_m Z_m \tag{27}$$

Here,  $\Psi$  denotes “decision-making mechanism coefficient”.

- i.  $\Theta$  is the minimal if  $\Psi \leq 0.5$
- ii.  $\Theta$  is the “maximum group utility” if  $\Psi \geq 0.5$ ,
- iii.  $\Theta$  is both the minimal and the “maximum group utility” if  $\Psi = 0.5$ .

Step-7: Ranking the alternative by  $\Gamma_m, Z_m$ , and  $\Theta_m$ .

**Step-8: Determine the compromise solution**

Obtain alternative  $\alpha^1$  as a compromise solution, that is ranked as the best by the measure  $\alpha'$  (minimal) if the A1 and A2 are satisfied:

**A1.** Acceptable stability:

$$\Theta(\alpha'^2) - \Theta(\alpha^1) \geq \frac{1}{r-1} \tag{28}$$

where  $\alpha^1, \alpha'^2$  are the alternatives with 1st and 2nd positions in the ranking by  $\Theta$ ;  $r$  = the number of alternatives.

**A2.** Acceptable stability in decision making:

Alternative  $\alpha^1$  must also be the best ranked by  $\Gamma$  or/and  $Z$ . This compromise solution is stable within whole decision making process.

- o  $\alpha^1$  and  $\alpha'^2$  are compromise solutions if A2 is not satisfied, or
- o  $\alpha^1, \alpha'^2, \dots, \alpha'^r$  are compromise solutions if A1 is not satisfied and  $\alpha'^r$  is decided by constraint  $\Theta(\alpha'^2) - \Theta(\alpha^1) \leq \frac{1}{r-1}$  for maximum  $r$ .

The minimal value of  $\Theta$  determines the best alternative.




Figure 1. VIKOR based MAGDM strategy in trapezoidal neutrosophic number environment

### 5. Numerical example

To illustrate the developed VIKOR strategy, we consider an MAGDM problem adapted from [57]. The considered MAGDM problem is described as follows:

An investment company constitutes a decision making board with three experts to invest certain amount of money in the best alternative. The experts evaluate the four alternatives and three attributes which are described below :

Alteranatives:

1. Car company(  $\alpha'_1$  )
2. Food company(  $\alpha'_2$  )
3. Computer company(  $\alpha'_3$  )
4. Arms company(  $\alpha'_4$  )

Attributes:

1. Risk factor(  $\beta'_1$  )
2. Growth factor(  $\beta'_2$  )
3. Environment impact(  $\beta'_3$  )

Suppose,  $\tau = (0.30, 0.42, 0.28)$  be the set of weigts of the decision makers and  $\sigma = (0.33, 0.39, 0.28)$  be the set of weights of the attributes.

Step-1: In this step, we construct the decision matrix in TrNNs form

Decision matrix  $D^1$

$$\left( \begin{array}{ccc} & \beta'_1 & \beta'_2 & \beta'_3 \\ \alpha'_1 & \langle (0.5, 0.6, 0.7, 0.8); 0.1, 0.4, 0.7 \rangle & \langle (0.1, 0.1, 0.2, 0.3); 0.6, 0.7, 0.5 \rangle & \langle (0.1, 0.2, 0.2, 0.3); 0.7, 0.2, 0.4 \rangle \\ \alpha'_2 & \langle (0.3, 0.4, 0.5, 0.5); 0.4, 0.5, 0.2 \rangle & \langle (0.1, 0.2, 0.2, 0.4); 0.1, 0.4, 0.3 \rangle & \langle (0.1, 0.1, 0.2, 0.3); 0.5, 0.3, 0.5 \rangle \\ \alpha'_3 & \langle (0.3, 0.3, 0.30, 3); 0.1, 0.2, 0.3 \rangle & \langle (0.2, 0.3, 0.4, 0.4); 0.8, 0.2, 0.5 \rangle & \langle (0.6, 0.7, 0.8, 0.9); 0.4, 0.3, 0.1 \rangle \\ \alpha'_4 & \langle (0.7, 0.8, 0.8, 0.9); 0.3, 0.3, 0.2 \rangle & \langle (0.1, 0.2, 0.3, 0.3); 0.6, 0.5, 0.2 \rangle & \langle (0.2, 0.2, 0.2, 0.2); 0.5, 0.2, 0.2 \rangle \end{array} \right) \tag{29}$$

Decision matrix  $D^2$

$$\left( \begin{array}{ccc} & \beta'_1 & \beta'_2 & \beta'_3 \\ \alpha'_1 & \langle (0.1, 0.1, 0.2, 0.3); 0.2, 0.5, 0.1 \rangle & \langle (0.2, 0.2, 0.3, 0.4); 0.2, 0.5, 0.1 \rangle & \langle (0.4, 0.5, 0.6, 0.7); 0.5, 0.7, 0.2 \rangle \\ \alpha'_2 & \langle (0.2, 0.3, 0.4, 0.5); 0.3, 0.3, 0.2 \rangle & \langle (0.1, 0.1, 0.2, 0.3); 0.3, 0.3, 0.4 \rangle & \langle (0.2, 0.2, 0.3, 0.3); 0.4, 0.5, 0.2 \rangle \\ \alpha'_3 & \langle (0.1, 0.2, 0.2, 0.3); 0.2, 0.5, 0.6 \rangle & \langle (0.2, 0.3, 0.3, 0.4); 0.2, 0.2, 0.1 \rangle & \langle (0.4, 0.5, 0.6, 0.6); 0.8, 0.1, 0.1 \rangle \\ \alpha'_4 & \langle (0.5, 0.6, 0.7, 0.7); 0.5, 0.2, 0.1 \rangle & \langle (0.2, 0.2, 0.2, 0.2); 0.3, 0.4, 0.5 \rangle & \langle (0.1, 0.1, 0.2, 0.2); 0.3, 0.7, 0.4 \rangle \end{array} \right) \tag{30}$$

Decision matrix  $D^3$

$$\left( \begin{array}{ccc} & \beta'_1 & \beta'_2 & \beta'_3 \\ \alpha'_1 & \langle (0.3, 0.4, 0.4, 0.5); 0.5, 0.1, 0.1 \rangle & \langle (0.1, 0.2, 0.2, 0.3); 0.5, 0.1, 0.1 \rangle & \langle (0.2, 0.2, 0.3, 0.4); 0.6, 0.2, 0.1 \rangle \\ \alpha'_2 & \langle (0.2, 0.2, 0.2, 0.2); 0.3, 0.2, 0.7 \rangle & \langle (0.1, 0.1, 0.1, 0.1); 0.4, 0.4, 0.1 \rangle & \langle (0.6, 0.7, 0.8, 0.8); 0.4, 0.1, 0.1 \rangle \\ \alpha'_3 & \langle (0.2, 0.3, 0.4, 0.5); 0.4, 0.5, 0.3 \rangle & \langle (0.2, 0.3, 0.3, 0.4); 0.5, 0.4, 0.3 \rangle & \langle (0.3, 0.4, 0.4, 0.5); 0.5, 0.2, 0.3 \rangle \\ \alpha'_4 & \langle (0.3, 0.4, 0.4, 0.5); 0.5, 0.2, 0.1 \rangle & \langle (0.1, 0.2, 0.2, 0.3); 0.5, 0.1, 0.1 \rangle & \langle (0.1, 0.2, 0.3, 0.4); 0.2, 0.2, 0.5 \rangle \end{array} \right) \tag{31}$$

Step-2: We do not need to standardize the defining matrix as all the criteria are profit type.

Step-3: Using TrNWAA operator of equation (20), we get aggregate decision matrix of (29), (30), and (31) which is presented below:

$$\left( \begin{array}{ccc} & \beta_1 & \beta_2 & \beta_3 \\ \alpha_1 & \langle (0.276, 0.334, 0.406, 0.506); 0.273, 0.298, 0.179 \rangle & \langle (0.142, 0.17, 0.242, 0.342); .431, 0.352, 0.162 \rangle & \langle (0.254, 0.326, 0.396, 0.496); 0.633, 0.294, 0.203 \rangle \\ \alpha_2 & \langle (0.23, 0.302, 0.374, 0.416); 0.332, 0.312, 0.284 \rangle & \langle (0.1, 0.13, 0.172, 0.274); 0.277, 0.354, 0.249 \rangle & \langle (0.282, 0.31, 0.41, 0.44); 0.564, 0.242, 0.217 \rangle \\ \alpha_3 & \langle (0.188, 0.258, 0.286, 0.356); 0.235, 0.380, 0.401 \rangle & \langle (0.2, 0.3, 0.33, 0.372); 0.537, 0.242, 0.162 \rangle & \langle (0.432, 0.532, 0.604, 0.662); 0.640, 0.169, 0.136 \rangle \\ \alpha_4 & \langle (0.504, 0.604, 0.646, 0.704); 0.447, 0.226, 0.123 \rangle & \langle (0.142, 0.2, 0.23, 0.258); 0.461, 0.290, 0.242 \rangle & \langle (0.13, 0.158, 0.228, 0.256); 0.343, 0.338, 0.346 \rangle \end{array} \right)$$

step-4: Here we define positive ideal soluton and negative solution by employing equations (21) and (22)

The positive ideal solution  $R^+$  is presented as:

$$\langle (0.504, 0.604, 0.646, 0.704); 0.447, 0.226, 0.123 \rangle \quad \langle (0.2, 0.3, 0.33, 0.372); 0.537, 0.242, 0.162 \rangle \quad \langle (0.432, 0.532, 0.604, 0.662); 0.640, 0.169, 0.136 \rangle$$

The negative ideal solution  $R^-$  is presented as:

$$\langle (0.188, 0.258, 0.286, 0.356); 0.235, 0.380, 0.401 \rangle \quad \langle (0.1, 0.13, 0.172, 0.274); 0.277, 0.354, 0.249 \rangle \quad \langle (0.13, 0.158, 0.228, 0.256); 0.343, 0.338, 0.346 \rangle$$

Step-5: Using equations (23) and (24), we compute  $\Gamma_m$  and  $Z_m$  which are presented as:

$$\Gamma_1 = \left(\frac{0.33 \times 0.202}{0.294}\right) + \left(\frac{0.39 \times 0.070}{0.121}\right) + \left(\frac{0.28 \times 0.172}{0.327}\right) = 0.601,$$

$$\Gamma_2 = \left(\frac{0.33 \times 0.238}{0.294}\right) + \left(\frac{0.39 \times 0.119}{0.121}\right) + \left(\frac{0.28 \times 0.181}{0.327}\right) = 0.805,$$

$$\Gamma_3 = \left(\frac{0.33 \times 0.298}{0.294}\right) + \left(\frac{0.39 \times 0}{0.121}\right) + \left(\frac{0.28 \times 0}{0.327}\right) = 0.334,$$

$$\Gamma_4 = \left(\frac{0.33 \times 0}{0.294}\right) + \left(\frac{0.39 \times 0.080}{0.121}\right) + \left(\frac{0.28 \times 0.284}{0.327}\right) = 0.501.$$

Here we use Hamming distance to measure the distance between two TrNN .

And

$$Z_1 = \max \left\{ \left(\frac{0.33 \times 0.202}{0.294}\right) + \left(\frac{0.39 \times 0.070}{0.121}\right) + \left(\frac{0.28 \times 0.172}{0.327}\right) \right\} = 0.227,$$

$$Z_2 = \max \left\{ \left(\frac{0.33 \times 0.238}{0.294}\right) + \left(\frac{0.39 \times 0.119}{0.121}\right) + \left(\frac{0.28 \times 0.181}{0.327}\right) \right\} = 0.383,$$

$$Z_3 = \max \left\{ \left(\frac{0.33 \times 0.298}{0.294}\right) + \left(\frac{0.39 \times 0}{0.121}\right) + \left(\frac{0.28 \times 0}{0.327}\right) \right\} = 0.334,$$

$$Z_4 = \max \left\{ \left(\frac{0.33 \times 0}{0.294}\right) + \left(\frac{0.39 \times 0.080}{0.121}\right) + \left(\frac{0.28 \times 0.284}{0.327}\right) \right\} = 0.258.$$

Step-6: Using (25),(26), and (27) we calculate  $\Theta_i$

$$\Theta_1 = 0.283, \Theta_2 = 1, \Theta_3 = 0.342, \Theta_4 = 0.274$$

Step -7: The ranking order of alternatives is

$$\Theta_4 \leq \Theta_1 \leq \Theta_3 \leq \Theta_2$$

**Table 1.** Preference ranking order and compromise solution based on  $\Gamma$  ,  $Z$  and  $\Theta$

	$\alpha'_1$	$\alpha'_2$	$\alpha'_3$	$\alpha'_4$	Ranking	Compromise solution
$\Gamma$	0.6	0.805	0.334	0.501	$\alpha'_3 < \alpha'_4 < \alpha'_1 < \alpha'_2$	$\alpha'_3$
$Z$	0.228	0.383	0.334	0.258	$\alpha'_1 < \alpha'_3 < \alpha'_3 < \alpha'_2$	$\alpha'_1$
$\Theta(\Psi = 0.5)$	0.282	1	0.342	0.274	$\alpha'_4 < \alpha'_1 < \alpha'_3 < \alpha'_2$	$\alpha'_4$

**Step 8: Determine the compromise solution**

If we rank  $\Theta$  in decreasing order, the best position alternative is  $\alpha'_4$  with  $\Theta(\alpha'_4) = 0.274$ , and the 2nd best position  $\alpha'_1$  with  $\Theta(\alpha'_1) = 0.283$ . Therefore,  $\Theta(\alpha'_1) - \Theta(\alpha'_4) = 0.008 < 0.33$  (since  $r = 4; 1/(r-1) = 0.33$ ), which

does not satisfy the condition  $1(\Theta(\alpha'^2) - \Theta(\alpha'^1)) \geq \frac{1}{r-1}$ .

Here  $\alpha'_4$  is ranked best by  $\Gamma$  and  $Z$  and satisfies the condition 2.

So, the compromise solution as follows:

$$\Theta(\alpha'_1) - \Theta(\alpha'_4) = 0.008 < 0.33,$$

$$\Theta(\alpha'_2) - \Theta(\alpha'_4) = 0.726 > 0.33,$$

$$\Theta(\alpha'_3) - \Theta(\alpha'_4) = 0.05 < 0.33,$$

Therefore,  $\alpha'_1, \alpha'_3$  and  $\alpha'_4$  are compromise solutions.

**6.1 The impact of parameter  $\Psi$**

Table 2 demonstrates how the different values of  $\Psi$  impact the ranking order of the alternatives  $\alpha'_i$ .

**Table 2.** For different values of  $\Psi$ , ranking the value of  $\alpha'_i$  ( $i' = 1, 2, 3, 4$ ).

Values of $\Psi$	Values of $\alpha'_i$	Preference order
$\Psi = 0.1$	$\alpha'_1 = 0.057, \alpha'_2 = 1, \alpha'_3 = 0.615, \alpha'_4 = 0.209$	$\alpha'_1 < \alpha'_4 < \alpha'_3 < \alpha'_2$
$\Psi = 0.2$	$\alpha'_1 = 0.113, \alpha'_2 = 1, \alpha'_3 = 0.547, \alpha'_4 = 0.225$	$\alpha'_1 < \alpha'_4 < \alpha'_3 < \alpha'_2$
$\Psi = 0.3$	$\alpha'_1 = 0.170, \alpha'_2 = 1, \alpha'_3 = 0.479, \alpha'_4 = 0.241$	$\alpha'_1 < \alpha'_4 < \alpha'_3 < \alpha'_2$
$\Psi = 0.4$	$\alpha'_1 = 0.227, \alpha'_2 = 1, \alpha'_3 = 0.410, \alpha'_4 = 0.257$	$\alpha'_1 < \alpha'_4 < \alpha'_3 < \alpha'_2$
$\Psi = 0.5$	$\alpha'_1 = 0.282, \alpha'_2 = 1, \alpha'_3 = 0.342, \alpha'_4 = 0.274$	$\alpha'_4 < \alpha'_1 < \alpha'_3 < \alpha'_2$
$\Psi = 0.6$	$\alpha'_1 = 0.340, \alpha'_2 = 1, \alpha'_3 = 0.274, \alpha'_4 = 0.290$	$\alpha'_3 < \alpha'_4 < \alpha'_1 < \alpha'_2$
$\Psi = 0.7$	$\alpha'_1 = 0.370, \alpha'_2 = 1, \alpha'_3 = 0.205, \alpha'_4 = 0.306$	$\alpha'_3 < \alpha'_4 < \alpha'_1 < \alpha'_2$
$\Psi = 0.8$	$\alpha'_1 = 0.454, \alpha'_2 = 1, \alpha'_3 = 0.137, \alpha'_4 = 0.399$	$\alpha'_3 < \alpha'_4 < \alpha'_1 < \alpha'_2$
$\Psi = 0.9$	$\alpha'_1 = 0.510, \alpha'_2 = 1, \alpha'_3 = 0.068, \alpha'_4 = 0.338$	$\alpha'_3 < \alpha'_4 < \alpha'_1 < \alpha'_2$

**7. Conclusions**

Extended VIKOR strategy for MAGDM in trapezoidal neutrosophic number environment is presented in the paper. TrNWAA operator and Hamming distance are employed to develop the VIKOR strategy for MAGDM. Finally, an MAGDM problem is solved to demonstrate the proposed VIKOR strategy. Here, a sensitivity analysis is performed to demonstrate the impact of different values of the “decision making mechanism coefficient” on ranking system. The proposed extended VIKOR strategy for MAGDM problems can be used to deal with decision making problems such as brick selection [60, 61], logistics center selection [62], teacher selection [63], weaver selection [64], etc.

**References**

[1] F. Smarandache. A unifying field of logics. Neutrosophy: neutrosophic probability, set and logic, American Research Press, Rehoboth, (1998).  
 [2] H. Wang, F. Smarandache, Y. Zhang, and R. Sunderraman. Single valued neutrosophic sets. Multi-space and Multi-structure, 4 (2010), 410-413.  
 [3] P. Biswas, S. Pramanik, and B. C. Giri. Entropy based grey relational analysis method for multi-attribute decision making under single valued neutrosophic assessments. Neutrosophic Sets and Systems, 2 (2014), 102–110.  
 [4] P. Biswas, S. Pramanik, and B. C. Giri. A new methodology for neutrosophic multi-attribute decision making with unknown weight information. Neutrosophic Sets and Systems, 3 (2014), 42–52.

- [5] P. Biswas, S. Pramanik, and B. C. Giri. TOPSIS method for multi-attribute group decision-making under single valued neutrosophic environment. *Neural Computing and Applications*, (2015). doi: 10.1007/s00521-015-1891-2.
- [6] P. Biswas, S. Pramanik, and B. C. Giri. Aggregation of triangular fuzzy neutrosophic set information and its application to multi-attribute decision making. *Neutrosophic Sets and Systems*, 12 (2016), 20-40.
- [7] P. Biswas, S. Pramanik, and B. C. Giri. Value and ambiguity index based ranking method of single-valued trapezoidal neutrosophic numbers and its application to multi-attribute decision making. *Neutrosophic Sets and Systems* 12 (2016), 127-138.
- [8] P. Biswas, S. Pramanik, and B. C. Giri. Multi-attribute group decision making based on expected value of neutrosophic trapezoidal numbers. *New trends in neutrosophic theory and applications-Vol-II*. Pons Editions, Brussels (2018), (pp. 103-124).
- [9] P. Biswas, S. Pramanik, and B. C. Giri. Non-linear programming approach for single-valued neutrosophic TOPSIS method. *New Mathematics and Natural Computation*, (2018). In Press.
- [10] I. Deli, and Y. Subas. A ranking method of single valued neutrosophic numbers and its applications to multi-attribute decision making problems. *International Journal of Machine Learning and Cybernetics*, (2016). doi:10.1007/s13042016-0505-3.
- [11] P. Ji, J. Q. Wang, and H. Y. Zhang. Frank prioritized Bonferroni mean operator with single-valued neutrosophic sets and its application in selecting third-party logistics providers. *Neural Computing and Applications*, (2016). doi:10.1007/s00521-016-2660-6.
- [12] A. Kharal. A neutrosophic multi-criteria decision making method. *New Mathematics and Natural Computation*, 10 (2014), 143–162.
- [13] K. Mondal, and S. Pramanik. Neutrosophic tangent similarity measure and its application to multiple attribute decision making. *Neutrosophic Sets and Systems*, 9(2015), 85-92.
- [14] R. X. Liang, J. Q. Wang, and H. Y. Zhang. A multi-criteria decision-making method based on single-valued trapezoidal neutrosophic preference relations with complete weight information. *Neural Computing and Applications*, (2017). doi: 10.1007/s00521-017-2925-8.
- [15] P. Liu, Y. Chu, Y. Li, and Y. Chen. Some generalized neutrosophic number Hamacher aggregation operators and their application to group decision making. *International Journal of Fuzzy System*, 16 (2) (2014), 242–255.
- [16] P. D. Liu, and H. G. Li. Multiple attribute decision-making method based on some normal neutrosophic Bonferroni mean operators. *Neural Computing and Applications*, 28 (2017), 179–194.
- [17] P. Liu, and Y. Wang. Multiple attribute decision-making method based on single-valued neutrosophic normalized weighted Bonferroni mean. *Neural Computing and Applications*, 25(7) (2014), 2001–2010.
- [18] F. Smarandache, and S. Pramanik (Eds). *New trends in neutrosophic theory and applications*, Vol.2. Brussels: Pons Editions, 2018.
- [19] F. Smarandache, and S. Pramanik (Eds). *New trends in neutrosophic theory and applications*. Brussels: Pons Editions, 2016.
- [20] S. Pramanik, P. Biswas, and B. C. Giri. Hybrid vector similarity measures and their applications to multi-attribute decision making under neutrosophic environment. *Neural Computing and Applications*, 28 (5)(2017), 1163-1176. DOI 10.1007/s00521-015-2125-3.
- [21] R. Sahin, and A. Kucuk. Subsethood measure for single valued neutrosophic sets. *Journal of Intelligent and Fuzzy System*, (2014). doi:10.3233/IFS-141304.
- [22] Biswas, P., Pramanik, S., & Giri, B. C. (2018). Neutrosophic TOPSIS with group decision making. In C. Kahraman & I. Otay (Eds.): *C. Kahraman and I. Otay (eds.), Fuzzy Multicriteria Decision Making Using Neutrosophic Sets*, *Studies in Fuzziness and Soft Computing* 369 (pp. 543-585). doi. 10.1007/978-3-030-00045-5\_21.
- [23] J. Ye. Multicriteria decision-making method using the correlation coefficient under single-valued neutrosophic environment. *International Journal of General Systems*, 42 (2013), 386–394.
- [24] J. Ye. Single valued neutrosophic cross-entropy for multi criteria decision making problems. *Applied Mathematical Modelling*, 38 (3) (2013), 1170–1175.
- [25] J. Ye. Projection and bidirectional projection measures of single valued neutrosophic sets and their decision – making method for mechanical design scheme. *Journal of Experimental and Theoretical Artificial Intelligence*, (2016). doi:10.1080/0952813X.2016.1259263.
- [26] M. Abdel-Basset, M. Mohamed, and F. Smarandache. An extension of neutrosophic AHP–SWOT analysis for strategic planning and decision-making, *Symmetry*, 10( 4) (2018), 116. <https://doi.org/10.3390/sym10040116>
- [27] S. Pramanik, S. Dalapati, S. Alam, F. Smarandache, and T. K. Roy. NS-cross entropy-based MAGDM under single-valued neutrosophic set environment. *Information*, 9(2) (2018), 37. doi:10.3390/info9020037
- [28] S. Pramanik, R. Mallick, and A. Dasgupta, Contributions of selected Indian researchers to multi-attribute decision making in neutrosophic environment. *Neutrosophic Sets and Systems*, 20 (2018), 108-131.
- [29] K. Mondal, S. Pramanik, and B. C. Giri. Single valued neutrosophic hyperbolic sine similarity measure based MADM strategy. *Neutrosophic Sets and Systems*, 20 (2018), 3-11. <http://doi.org/10.5281/zenodo.1235383>.
- [30] K. Mondal, S. Pramanik, and B. C. Giri. Hybrid binary logarithm similarity measure for MAGDM problems under SVNS assessments. *Neutrosophic Sets and Systems*, 20 (2018), 12-25. <http://doi.org/10.5281/zenodo.1235365>.
- [31] K. Mondal, S. Pramanik, and B. C. Giri. Interval neutrosophic tangent similarity measure based MADM strategy and its application to MADM problems. *Neutrosophic Sets and Systems*, 19 (2018), 47-56.

- [32] H. D. Cheng, and Y. Guo. A new neutrosophic approach to image thresholding. *New Mathematics and Natural Computation*, 4 (2008), 291–308.
- [33] Y. Guo, and H. D. Cheng. New neutrosophic approach to image segmentation. *Pattern Recognition*, 42 (2009), 587-595.
- [34] Y. Guo, A. Sengur, and J. Ye. A novel image thresholding algorithm based on neutrosophic similarity score. *Measurement*, 58 (2014), 175–186.
- [35] M. Zhang, L. Zhang, and H. D. Cheng. A neutrosophic approach to image segmentation based on watershed method. *Signal Processing*, 90 (5) (2010), 1510-1517.
- [36] S. Pramanik, and T. K. Roy. Neutrosophic game theoretic approach to Indo-Pak conflict over Jammu-Kashmir. *Neutrosophic Sets and Systems*, 2 (2014), 82-101.
- [37] K. Mondal, and S. Pramanik. Multi-criteria group decision making approach for teacher recruitment in higher education under simplified Neutrosophic environment. *Neutrosophic Sets and Systems*, 6 (2014), 28-34.
- [38] K. Mondal, and S. Pramanik. Neutrosophic decision making model of school choice. *Neutrosophic Sets and Systems*, 7 (2015), 62-68.
- [39] K. Mondal, and S. Pramanik. A study on problems of Hijras in West Bengal based on neutrosophic cognitive maps. *Neutrosophic Sets and Systems*, 5(2014), 21-26.
- [40] S. Pramanik, and S. Chakrabarti. A study on problems of construction workers in West Bengal based on neutrosophic cognitive maps. *International Journal of Innovative Research in Science. Engineering and Technology*, 2(11) (2013), 6387-6394.
- [41] S. Broumi, A. Bakali, M. Talea, F. Smarandache, V. Uluçay, S. Sahin, A. Dey, M. Dhar, R.P. Tan, A. de Oliveira, and S. Pramanik. Neutrosophic sets: An overview. In F. Smarandache, & S. Pramanik (Eds., vol.2), *New trends in neutrosophic theory and applications* (2018)(pp. 403-434). Brussels: Pons Editions.
- [42] X. Peng, and J. Dai. A bibliometric analysis of neutrosophic set: two decades review from 1998 to 2017. - *Artificial Intelligence Review*, 2018. <https://doi.org/10.1007/s10462-018-9652-0>.
- [43] J. Ye. Some weighted aggregation operator of trapezoidal neutrosophic number and their multiple attribute decision making method. *Informatica*, 28 (2) (2017), 387-402.
- [44] P. Biswas, S. Pramanik, and B. C. Giri. TOPSIS strategy for multi-attribute decision making with trapezoidal neutrosophic numbers. *Neutrosophic Sets and Systems*, 19(2018), 29-39.
- [45] P. Biswas, S. Pramanik, and B. C. Giri. Distance measure based MADM strategy with interval trapezoidal neutrosophic numbers. *Neutrosophic Sets and Systems*, 19(2018), 40-46.
- [46] S. Opricovic. *Multicriteria Optimization of Civil Engineering Systems*; Faculty of Civil Engineering: Belgrade, Serbia, 1998.
- [47] S. Opricovic, and G. H. Tzeng. Compromise solution by MCDM methods: A comparative analysis of VIKOR and TOPSIS. *European journal of operational research*, 156 (2004), 445–455.
- [48] S. Opricovic, and G. H. Tzeng. Extended VIKOR method in comparison with outranking methods. *European journal of operational research*, 178 (2) (2007), 514-529.
- [49] H. Pouresmaeil, E. Shivanian, E. Khorram, and H. S. Fathabadi. An extended method using TOPSIS and VIKOR for multiple attribute decision making with multiple decision makers and single valued neutrosophic numbers. *Advances and Applications in Statistics*, 50(4) (2017), 261.
- [50] R. Bausys, and E. K. Zavadskas. Multicriteria decision making approach by VIKOR under interval neutrosophic set environment. *Economic Computation and Economic Cybernetics Studies and Research*, 4 (2015), 33-48.
- [51] Y. Huang, G. Wei, and C. Wei. VIKOR method for interval neutrosophic multiple attribute group decision-making. *Information*, 8 (2017), 144. doi:10.3390/info8040144.
- [52] P. Liu, L. Zhang. The extended VIKOR method for multiple criteria decision making problem based on neutrosophic hesitant fuzzy set, (2014).
- [53] J. Hu, L. Pan, and X. Chen. An interval neutrosophic projection-based VIKOR method for selecting doctors. *Cognitive Computation*, 9(6) (2017), 801-816.
- [54] K. Selvakumari, and M. A. Priyadarshini. VIKOR method for decision making problem using octagonal neutrosophic soft matrix. *International Journal of Latest Engineering Research and Applications*, 2(7) (2017), 41-45.
- [55] S. Pramanik, S. Dalapati, S. Alam, and T.K. Roy. VIKOR based MAGDM strategy under bipolar neutrosophic set environment *Neutrosophic Sets and Systems*, 19(2018), 57-69. <http://doi.org/10.5281/zenodo.1235341>.
- [56] S. Pramanik, S. Dalapati, S. Alam, and T.K. Roy. NC-VIKOR based MAGDM strategy under neutrosophic cubic set environment. *Neutrosophic Sets and Systems*, 20(2018), 95-108. <http://doi.org/10.5281/zenodo.1235367>
- [57] S. Dalapati and S. Pramanik. A revisit to NC-VIKOR based MAGDM strategy in neutrosophic cubic set environment. *Neutrosophic Sets and Systems*, 21(2018), 131-141. <https://doi.org/10.5281/zenodo.1408665>.
- [58] L.A. Zadeh. Fuzzy sets. *Information and Control*, 8 (3) (1965), 338–353.
- [59] A.Kumar. P. Singh, A. Kaur, and P. Kaur. Ranking of generalized trapezoidal fuzzy numbers based on rank, mode, divergence and spread. *Turkish Journal of Fuzzy Systems*, 1(2) (2010), 141-152.
- [60] K. Mondal, and S. Pramanik. Intuitionistic fuzzy multicriteria group decision making approach to quality-brick selection problem *Journal of Applied Quantitative Methods*, 9(2) (2014), 35-50.
- [61] K. Mondal, and S. Pramanik. Neutrosophic decision making model for clay-brick selection in construction field based on grey relational analysis. *Neutrosophic Sets and Systems*, 9 (2015), 64-71.

- [62] S. Pramanik, S. Dalapati, and T. K. Roy. Neutrosophic multi-attribute group decision making strategy for logistic center location selection. In F. Smarandache, M. A. Basset & V. Chang (Eds.), *Neutrosophic Operational Research*, Vol. III. Pons asbl, Brussels, 2018, 13-32.
- [63] S. Pramanik, and D. Mukhopadhyaya. Grey relational analysis based intuitionistic fuzzy multi-criteria group decision-making approach for teacher selection in higher education. *International Journal of Computer Applications*, 34 (2011), 21–29.
- [64] P. P. Dey, S. Pramanik, and B. C. Giri. Multi-criteria group decision making in intuitionistic fuzzy environment based on grey relational analysis for weaver selection in Khadi institution. *Journal of Applied and Quantitative Methods*, 10 (4) (2015), 1-14.

Received: October 1, 2018. Accepted: October 31, 2018